

The value of a single character: the Paleogene European land snail *Ferussina* Grateloup, 1827 is likely a cyclophorid (Gastropoda, Caenogastropoda)

Barna Páll-Gergely¹, Thomas A. Neubauer^{2,3}

1 Plant Protection Institute, Centre for Agricultural Research, Herman Ottó Street 15, Budapest, H-1022, Hungary **2** Department of Animal Ecology & Systematics, Justus Liebig University, Heinrich-Buff-Ring 26–32 IFZ, 35392 Giessen, Germany **3** Naturalis Biodiversity Center, Darwinweg 2, 2333 CR Leiden, The Netherlands

Corresponding author: Barna Páll-Gergely (pall-gergely.barna@agrar.mta.hu; pallgergely2@gmail.com)

Academic editor: M. Haase | Received 14 January 2020 | Accepted 30 January 2020 | Published 12 March 2020

<http://zoobank.org/F93EAB04-79A5-4514-AE6C-1EDCF36DFF2F>

Citation: Páll-Gergely B, Neubauer TA (2020) The value of a single character: the Paleogene European land snail *Ferussina* Grateloup, 1827 is likely a cyclophorid (Gastropoda, Caenogastropoda). ZooKeys 918: 29–40. <https://doi.org/10.3897/zookeys.918.50135>

Abstract

Ferussina Grateloup, 1827 is a European Paleogene land snail genus, which is currently classified in its own family, the Ferussinidae Wenz, 1923 (1915), in the superfamily Cyclophoroidea. The shell of this genus is remarkable by its last quarter whorl turning towards the apex instead of away from it, which is an unusual trait in terrestrial snails. We show, however, that this trait has evolved at least nine times in terrestrial Eupulmonata and Caenogastropoda, and it does not justify distinction at the family level in any of the reported cases. This observation suggests the systematic position of *Ferussina* should not be based on the apexward-turning last quarter whorl alone but instead on the general morphology of the shell. As a result, we re-evaluate the systematic position of the Ferussinidae and treat it as a subfamily of the Cyclophoridae.

Keywords

character evolution, Eocene, Oligocene, parallel evolution, terrestrial Gastropoda, unique trait

Introduction

Ferussina Grateloup, 1827 (and its synonym *Strophostoma* Deshayes, 1828; see Wenz 1923; Kadolsky 2008) is a genus reported from middle Eocene (Lutetian) to upper Oligocene (Chattian) deposits of France, Germany, Italy, and Switzerland (Fig. 1);




Figure 1. Geographic and stratigraphic distribution of *Ferussina* in central and western Europe. Records that are questionable in terms of stratigraphic horizon or species identification are indicated with a question mark (see Suppl. material 1: Table S1 for more information). The map was created with ESRI ArcGIS 10.4.

a dubious record comes from presumably lower Miocene strata of southern France (Degrange-Touzin 1892). It is currently classified in its own family, the Ferussinidae Wenz, 1923 (1915) (Bouchet et al. 2017) in the superfamily Cyclophoroidea Gray, 1847. *Ferussina* is characterized by a relatively large (ca 1–3 cm), depressed-globular shell with an obtusely conical spire, a round aperture, and a last quarter whorl turning towards the apex (Sandberger 1870–1875; Roman 1899; Rey 1968; Kadolsky 2008; Salvador et al. 2016). As a result, the aperture opens in the adapical direction of the shell, orientating the umbilicus of the shell upwards while the animal was crawling.

In his catalogue of fossil non-marine gastropods, Wenz (1923) included seven species in the genus *Ferussina*, i.e. *F. anomphalus* (Sandberger, 1871), *F. anostomaeformis* Grateloup, 1827 (the type species by monotypy), *F. globosa* Dumas, 1876, *F. lapicida* Leufroy, 1828, *F. praeglobosa* (Roman, 1904), *F. striata* (Deshayes, 1828), and *F. tricarinata* (Braun, 1838). These are distinguished by the relative height of the spire, presence of an inflation on the last whorl, presence of a keel or angulation on the last whorl, and




Figure 2. Representatives of the genus *Ferussina*. **A–C** *Ferussina anostomaeformis* Grateloup, 1827, Gaas (Larrat), France, lower Oligocene; syntype, UBRG, Grateloup collection, no. 65-2-150 **D** *Ferussina anostomaeformis*, St-Paul-lès-Dax (Abesse, “Château”), France, upper Oligocene; MNHN.FA72133 **E–H** *Ferussina anomphalus capellinii* (Sandberger, 1873), Blaustein (Arnegg), Germany, lower Oligocene; syntype, SMNS 22180 **I–L** *Ferussina tricarinata* (Braun, 1838), Hochheim, Germany, upper Oligocene (Hochheim Formation, “Landschneckenkalk”); NHMW 75000/E/1778. Photos: Laurent Charles (**A–C**), Pierre Lozouet (**D**), Rodrigo Salvador (**E–H**), Barna Páll-Gergely (**I–L**). Scale bars: 1 cm.

presence and width of an umbilicus, as well as surface ornamentation (Fig. 2; compare also Deshayes 1828; Sandberger 1870–1875; Roman 1899, 1904; Kadolsky 2008). Sculpture ranges from fine to distinct, riblet-like growth lines and, in the case of *F. tricarinata*, narrow spiral keels on base and periphery.

An “upright” turning last whorl (termed “anostomy” by Nordsieck 1986) is unusual in terrestrial snails but has repeatedly evolved in both the Eupulmonata (“pulmonates”) (at least six times) and Caenogastropoda (at least three times) (Schileyko 1998, 1999, 2000; Egorov 2009, 2013), and even in a Devonian marine gastropod (Braun 1838). This trait has not been considered a justification for the distinction at the family level in

any of the reported cases. This observation suggests the systematic position of *Ferussina* should not be based on the apexward-turning last quarter whorl alone but instead on the general morphology of the shell. As a result, we re-evaluate the systematic position of the Ferussinidae and treat it as a subfamily of the Cyclophoridae Gray, 1847.

Abbreviations used:

MNHN – Muséum National d’Histoire Naturelle, Paris; NHMW – Natural History Museum Vienna; SMNS – State Museum of Natural History Stuttgart; UBRG – Université du Bordeaux.

Results

While the apexward-turning last whorls are unique among fossil European land snails, we found this trait in a number of unrelated extant and fossil clades, including 12 pulmonate (Wenz 1940; Schileyko 1998, 1999, 2000) and four caenogastropod genera (Egorov 2009, 2013), representing at least nine independent events. Table 1 summarises the key information.

Table 1. Summary of key information of extant land snail genera with apexward turning body whorl. In addition, we provide information on shell shape of relatives within the same family to assess the relevance of shape traits for systematic placement. Information derives from Wenz (1940) and Schileyko (1998, 1999, 2000).

Genera	Size (mm)	Shell shape without body whorl	Habitat	Geographic region	Shell shape of relatives
<i>Boysia</i> , <i>Gyliotrachela</i> , <i>Hypselostoma</i>	2–4	ovoid, conic	rock surfaces	Southeast Asia to Australia	ranging from ovoid and conic to lenticular and globular
<i>Campolaemus</i>	2	ovoid	not rock-dwelling	Saint Helena	unknown (might be ovoid or depressed)
<i>Anostoma</i> , <i>Clinispira</i> , <i>Ringicella</i>	14–16	obesely lenticular (depressed-globular), elongate-conical	not rock-dwelling (caves, under stones/logs)	Brazil	mostly high-spined
<i>Hendersoniella</i>	11–13	flat	rock surfaces	Mexico	all high-spined
<i>Tonkinia</i>	4.3–5	elliptical	not rock-dwelling	Vietnam	mostly high-spined
<i>Anostomopsis</i> , <i>Enneopsis</i> , <i>Srophostomella</i>	11–18	cup-shaped, ovoid, depressed globular	unknown	Austria, Hungary, France	planispiral, lenticular, ovoid
<i>Anosycolus</i>	12	conical	unknown, probably not rock-dwelling	Madagascar	conic, ovoid, high-spined
<i>Laotia</i>	2.2–4.4	depressed globular	unknown, probably not rock-dwelling	Laos and Vietnam	mostly low-spined to conical
<i>Opisthostoma</i> , <i>Plectostoma</i>	1.0–3.7	ovoid to depressed-globular	rock surfaces	Southeast Asia	ovoid to conic

Eupulmonata

(1) Genera *Boysia* Pfeiffer, 1849, *Gyliotrachela* Tomlin, 1930, *Hypselostoma* Benson, 1856

Remarks. These genera were included in the family Hypselostomatidae by Schileyko (1998), which was recognized as a subfamily of Gastrocoptidae Pilsbry, 1918 by Bouchet et al. (2017). Other genera of the same (sub)family are variable in shape, ranging from ovoid and conic to lenticular and globular. The direction of the aperture is variable even in the same genus. Some *Hypselostoma* and *Gyliotrachela* species have even normally coiled shells. The shells are small (2–4 mm). All the species with detached last whorl inhabit limestone rock areas and spend a considerable time of their life tightly attached to rock surfaces (Panha and Burch 2005).

(2) Genus *Campolaemus* Pilsbry, 1892

Remarks. This genus was classified in the Hypselostomatidae by Schileyko (1998). However, this species more probably belongs to the Streptaxidae (Páll-Gergely 2020). Nevertheless, its position within that family is questionable. Shell height is ca 2 mm. No information on its habitat preference is known. However, it is probably not a rock-dwelling species, because streptaxids typically occur among leaf litter, in decaying plant material, and under logs and stones (Páll-Gergely pers. obs.).

(3) Genera *Anostoma* Fischer von Waldheim, 1807, *Clinispira* Simone & Casati, 2013, *Ringicella* Gray, 1847

Remarks. *Anostoma* was classified in the tribe Odontostomini (Bulimulidae, Bulimulinae) by Schileyko (1999), which was recognized as a distinct family by Bouchet et al. (2017). According to Schileyko (1999), there are 11 high-spired genera and 3 low-spired/globular genera in the Odontostomini, all of which comprise relatively large snails (30–45 mm in shell diameter). *Anostoma* inhabit the semi-arid biomes of Brazil (the Cerrado and Caatinga ecoregions), and living specimens are typically found under stones (Rodrigo Salvador, pers. comm.). The genus *Ringicella* (treated as a genus of its own by Simone 2006 and as a subgenus of *Anostoma* by Schileyko 1999) is known from the Amazon region, and animals have been found living under decaying logs (Rodrigo Salvador pers. comm.). *Clinispira* Simone & Casati, 2013 was collected in caves in the semi-dry environment of the Caatinga ecoregion (Simone and Casati 2013). Inferring from the flat profile of the peristome, *Clinispira* might live attached to rock surfaces.

(4) Genus *Hendersoniella* Dall, 1905

Remarks. This genus was classified in the Urocoptidae, Holospirinae (Schileyko 1999), where many high-spired genera belong. Shell diameter is 11–13 mm. *Hendersoniella*

are obligate rock-dwelling, as the other members of the family (“live snails were found under limestone slabs that were spalding from the underlying rock”; Thompson and Correa 1991: 15).

(5) Genus *Tonkinia* Mabilie, 1887

Remarks. This genus was classified as a member of the Streptaxidae, Streptaxinae by Schileyko (2000), and in the Diapheridae in MolluscaBase (2020) following Dance (1970), who mentioned that *Tonkinia* and its probably closest relative, *Platycochlium* Laidlaw, 1950, are most similar to juvenile shells of *Diaphera* Albers, 1850 and *Sinoenena* Kobelt, 1904. With the exception of *Platycochlium* and *Tonkinia*, all other diapherids are high-spired. The shell is 4.3–5 mm wide (Schileyko 2000). We have not found any published information about its habitat preference, but it probably lives among decaying plant material and under logs and stones as other Diapheridae.

(6) Genera *Anostomopsis* Sandberger, 1871, *Enneopsis* Wenz, 1940, *Strophostomella* Fischer, 1883

Remarks. The three genera derive from upper Cretaceous (Coniacian–Maastrichtian) strata of Europe (Austria, Hungary, and France) and are currently classified in the fossil family Anostomopsidae with uncertain position in the Stylommatophora (Nordsieck 2014, 2017). *Strophostomella* has a depressed-globular shell similar to that of *Ferussina* (Tausch 1886, there as “*Strophostoma*”), *Anostomopsis* has a peculiarly cup-shaped morphology with flat apical side and narrow, tube-like aperture (Sandberger 1870–1875), and *Enneopsis* is characterized by an ovoid shape (Roule 1886, as “*Anostomopsis*”). All share a complex system of internal plicae (Nordsieck 2014; see also Wenz 1940).

Caenogastropoda

(1) Genus *Anosycolus* Fischer-Piette, C.P. Blanc, F. Blanc & Salvat, 1993

Remarks. This taxon was classified in the Hainesiidae by Egorov (2009) and in the Cyclophoridae in MolluscaBase (2020). However, a current investigation suggests it is a relative of *Boucardicus*, which includes conical-globular and high-spired species and may deserve its own family within Cyclophoroidea (Páll-Gergely unpublished information). Shell does not exceed 12 mm in maximum diameter.

(2) Genus *Laotia* Saurin, 1953

Remarks. This genus was classified in the Diplommatinidae by Egorov (2013) and in the Alycaecidae in Do et al. (2015). Recent investigations corroborate placement in Alycaecidae, where it will be classified in a separate new subfamily together with *Messageria*

Bavay & Dautzenberg, 1904 (Páll-Gergely unpublished information). Shell diameter is 2.2–4.4 mm (Páll-Gergely 2014). Nothing is known about its habitat preference, but *Laotia* is probably not an obligate rock-dwelling genus, since the aperture is not flat in front profile to allow attachment to rock surfaces.

(3) Genera *Opisthostoma* W.T. Blanford & H.F. Blanford, 1860, *Plectostoma* Adams, 1865

Remarks. Both are members of the Diplommatinidae (Webster et al. 2012; Egorov 2013) together with a number of other genera usually possessing high-spired and ovoid shells. *Plectostoma* is 1.0–3.7 mm in shell height, *Opisthostoma* is less than 1.3 mm in largest measurement, and both are obligate limestone-dwelling genera (Liew et al. 2014; Vermeulen 1991).

Discussion

The list above shows that shells with the last whorl turned apexward are present in numerous unrelated lineages of pulmonate and operculate terrestrial snails. In all cases, the species and genera with this peculiar shape have normally coiled relatives. Similarly, the fossil *Ferussina* certainly evolved from normally coiled ancestors, and we should not give too great importance to this trait when determining its systematic position. Moreover, the closest relatives of these genera are often species with high-spired shells. This suggests that we cannot exclude high-spired cyclophoroideans from the possible relatives of *Ferussina*.

We can exclude the Pomatiidae as possible relatives, as members of this family have calcareous opercula that are often found as fossils. No such opercula have been documented for *Ferussina*. The Cochlostomatinae, also with numerous extant and fossil members, are smaller than *Ferussina* and are characterized by high, conical shells, and some members have calcareous opercula (Fehér 2004; Zallot et al. 2015). The cyclophoroid family Craspedopomatidae, represented by several fossil species in Europe, comprises only very small, globular forms of only a few millimetres in diameter (Wenz 1923; Harzhauser and Neubauer 2018).

The most probable group of relatives is the Cyclophoridae. Most members of this family have broadly conical shells similar to that of *Ferussina*, except for the apexwards turn of the last quarter whorl. Extant Cyclophoridae have non-calcareous opercula, which are not preserved as fossils. So far, 14 species of Cyclophoridae are known from the Cenozoic sedimentary record of Europe (Wenz 1923; Steklov 1966; Schütt 1991, 1997; Stworzewicz 1995). The oldest records derive from the upper Paleocene (Thanetian) of France. Earlier mentions of European cyclophoroids from the Jurassic and Cretaceous belong to the families Diplommatinidae, Megalostomatidae, and Pupiniidae, or are unassigned cyclophoroids (Hrubesch 1965; Bandel 1991, 1993; Neubauer et al. 2019). The genus *Ventriculus* Wenz in Fischer & Wenz, 1914 was classified in the

family Cyclophoridae, subfamily Pupinellinae by Wenz (1923), a group now included in Pupinidae (Bouchet et al. 2017). The Pupinidae presently inhabits Asia from India to the oceanic islands (Egorov 2013).

Cyclophoridae are otherwise mostly restricted to south-eastern Asia, and the European fossils represent a rare exception of biogeographic affinity between both regions. Only a few other taxa that are widespread in East Asia today are also found in the European Cenozoic fossil record, such as Diplommatinidae, Strobilopsidae, and Pupinidae (e.g. Wenz 1923; Manganelli et al. 2008; Páll-Gergely et al. 2015; Harzhauser and Neubauer 2018).

In summary, we suggest a revised systematic position of the genus *Ferussina* in the Cyclophoridae. Given the distinct biogeographic and stratigraphic setting and morphological differences to extant Cyclophoridae, we suggest to maintain the genus in a distinct subfamily, Ferussininae.

The extant genera with apexward-turning body whorl listed above inhabit various habitats, with about half of them being obligatory rock-dwellers, indicating that this peculiar trait can be developed under various environmental conditions. *Ferussina* lived in a period when the regions it occurred in central and western Europe (France, Switzerland, Germany, Austria, and northern Italy) were dominated by warm-temperate to subtropical evergreen forests (Pound and Salzmann 2017). The Late Oligocene *Ferussina tricarinata* was thriving in semiarid conditions in the Mainz Basin on the shores of a brackish to hypersaline lake (Kadolsky 1989). Other *Ferussina* species may have dwelled in more humid climates among leaf litter and under decaying logs, but we have insufficient data about the taphonomy and paleoecology of their occurrences.

Acknowledgements

We are grateful to Bruno Cahuzac, Laurent Charles, Anita Eschner, Pierre Lozouet, Hartmut Nordsieck, Emmanuel Robert, and Rodrigo Salvador for sending photos and/or providing valuable literature and information. Dietrich Kadolsky and Mathias Harzhauser are thanked for their constructive reviews. We are indebted to The Biodiversity Heritage Library for the multitude of rare literature made available to us (<http://www.biodiversitylibrary.org>). This study was supported by the MTA (Hungarian Academy of Sciences) Premium Post Doctorate Research Program granted to B. Páll-Gergely. T.A. Neubauer was supported by a DFG grant (no. NE 2268/2-1).

References

- Bandel K (1991) Gastropods from brackish and fresh water of the Jurassic–Cretaceous transition (a systematic reevaluation). *Berliner Geowissenschaftliche Abhandlungen*, A 134: 9–55.
- Bandel K (1993) Caenogastropoda during Mesozoic times. *Scripta Geologica*, Special Issue 2: 7–56.

- Bouchet P, Rocroi J-P, Hausdorf B, Kaim A, Kano Y, Nützel A, Parkhaev P, Schrödl M, Strong EE (2017) Revised classification, nomenclator and typification of gastropod and monoplacophoran families. *Malacologia* 61 (1–2): 1–526. <https://doi.org/10.4002/040.061.0201>
- Braun M (1838) Über eine neue Art von *Strophostoma* und ein neues Genus *Scoliostoma* mit ähnlicher Bildung des Gehäuses. *Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde* 1838: 291–298. <https://www.biodiversitylibrary.org/page/35870875>
- Dance SP (1970) Non-marine molluscs of Borneo 1. Streptaxacea: Streptaxidae. *Journal of Conchology* 27: 149–162.
- Degrange-Touzain A (1892) Étude sur la faune terrestre, lacustre et fluviatile de l'Oligocène supérieur et du Miocène dans le Sud-Ouest de la France et principalement dans la Gironde. *Actes de la Société Linnéenne de Bordeaux* 45: 125–230. <https://biodiversitylibrary.org/page/33424078>
- Deshayes G-P (1828) Mémoire sur le strophostome, nouveau genre de coquilles fossiles de la famille des hélices. *Annales des Sciences Naturelles* 13: 282–287. <https://biodiversitylibrary.org/page/5978285>
- Do DS, Nguyen THT, Do VN (2015) A checklist and classification of terrestrial prosobranch snails from Son La, north-western Vietnam. *Ruthenica* 25(4): 117–132. <https://www.biotaxa.org/Ruthenica/article/view/16805>
- Egorov RV (2009) *Treasure of Russian Shells; Supplement 3, A Review of the Genera of the Recent Terrestrial Pectinibranch Molluscs (Synopsis Mainly Based on Published Data); Part II, Littoriniformes: Hainesiidae, Aciculidae, Cyclophoridae, Craspedopomatidae*. Colus-Doverie, Moscow, 57 pp.
- Egorov RV (2013) *Treasure of Russian Shells. Supplement 3. A Review of the Genera of the Terrestrial Pectinibranch Molluscs (Synopsis Mainly Based on Published Data). Part III. Littoriniformes: Liareidae, Pupinidae, Diplommatinidae, Alycaeidae, Cochlostomidae*. Roman Egorov, Moscow.
- Fehér Z (2004) A revision of the genus *Cochlostoma*, subgenus *Titanopoma* (Gastropoda, Caenogastropoda, Cochlostomatidae), in particular the forms occurring in Albania. *Basteria* 68(1–3): 25–44.
- Grateloup J-PS de (1827) Description de plusieurs espèces de coquilles fossiles des environs de Dax (Landes). *Bulletin d'Histoire Naturelle de la Société Linnéenne de Bordeaux* 2(7): 3–26. <https://biodiversitylibrary.org/page/35729870>
- Harzhauser M, Neubauer TA (2018) Opole (Poland) – a historical key locality for middle Miocene terrestrial mollusc faunas. *Bulletin of Geosciences* 93(1): 71–146. <https://doi.org/10.3140/bull.geosci.1692>
- Hrubesch K (1965) Die santone Landschneckenfauna von Glanegg bei Salzburg, Österreich. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und historischen Geologie* 5: 83–120. https://www.zobodat.at/pdf/Mitt-Bayer-Staatsslg-Pal-hist-Geol_5_0083-0120.pdf
- Kadolsky D (1989) Stratigraphie und Molluskenfaunen von “Landschneckenkalk” und “Cerithienschichten” im Mainzer Becken (Oberoligozän bis Untermiozän?). *Stratigraphische, paläogeographische und paläoökologische Ergebnisse. Geologisches Jahrbuch, Reihe A* 110: 69–133.

- Kadolsky D (2008) Mollusks from the Late Oligocene of Oberleichtersbach (Rhön Mountains, Germany). Part 2: Gastropoda: Neritimorpha and Caenogastropoda. *Courier Forschungsinstitut Senckenberg* 260: 103–137.
- Liew T-S, Vermeulen JJ, bin Marzuki ME, Schilthuizen M (2014) A cybertaxonomic revision of the microlandsnail genus *Plectostoma* Adam (Mollusca, Caenogastropoda, Diplommatinidae), from Peninsular Malaysia, Sumatra and Indochina. *ZooKeys* 393: 1–107. <https://doi.org/10.3897/zookeys.393.6717>
- Manganelli G, Aldinucci M, Capezzuoli E, Benocci A (2008) Strobilopsid land snails (Gastropoda: Pulmonata, Strobilopsidae) from the late Messinian–Early Pliocene (?) of Balze di Caspreno (central Italy). *Bollettino della Società Paleontologica Italiana* 46 (2–3): 101–106.
- MolluscaBase (2020) MolluscaBase. <http://www.molluscabase.org> [Accessed on: 2020-01-06]
- Neubauer TA, Páll-Gergely B, Jochum A, Harzhauser M (2019) Striking case of convergence – alleged marine gastropods in Cretaceous Burmese amber are terrestrial cyclophoroids. Comment on Yu et al. *Palaeoworld* 28(4): 572–575. <https://doi.org/10.1016/j.palwor.2019.05.015>
- Nordsieck H (1986) The system of the Stylommatophora (Gastropoda), with special regard to the systematic position of the Clausiliidae, II. Importance of the shell and distribution. *Archiv für Molluskenkunde* 117(1–3): 93–116.
- Nordsieck H (2014) Annotated check-list of the genera of fossil land snails (Gastropoda: Stylommatophora) of western and central Europe (Cretaceous–Pliocene), with description of new taxa. *Archiv für Molluskenkunde* 143(2): 153–185. <https://doi.org/10.1127/arch.moll/1869-0963/143/153-185>
- Nordsieck H (2017) Pulmonata, Stylommatophora, Helicoidea: Systematics with Comments. *ConchBooks*, Harxheim, 98 pp.
- Páll-Gergely B (2014) Description of the second *Laotia* Saurin, 1953; a genus new to the fauna of Vietnam (Gastropoda: Cyclophoroidea). *Folia Malacologica* 22(4): 289–292. <https://doi.org/10.12657/folmal.022.025>
- Páll-Gergely B, Hunyadi A, Asami T (2015) A new Chinese species of *Eostrobilops* Pilsbry, 1927 with a checklist of *Eostrobilops* and *Enteroplax* Gude, 1897 species (Gastropoda, Pulmonata, Strobilopsidae). *ZooKeys* 508: 85–95. <https://doi.org/10.3897/zookeys.508.10004>
- Páll-Gergely B (2020) *Campolaemus* Pilsbry, 1892 is not a hypselostomatid but a streptaxid (Gastropoda: Eupulmonata). *Ruthenica* 30(1): 69–73.
- Panha S, Burch JB (2005) An introduction to the microsnailed Thailand. *Malacological Review* 37/38: 1–155.
- Pound MJ, Salzmann U (2017) Heterogeneity in global vegetation and terrestrial climate change during the late Eocene to early Oligocene transition. *Scientific Reports* 7: 43386. <https://doi.org/10.1038/srep43386>
- Rey R (1968) L'Oligocène a partir des mollusques continentaux. *Annales Guébard* 44: 162–196.
- Roman F (1899) Monographie de la faune lacustre de l'Éocène moyen. *Annales de l'Université de Lyon, Nouvelle Série, I. Sciences, Médecine* 1: 1–41. <https://doi.org/10.5962/bhl.title.42238>
- Roman F (1904) Contributions à l'étude des bassins lacustres de l'Éocène et de l'Oligocène du Languedoc. *Bulletin de la Société Géologique de France, quatrième série* 3: 546–616.

http://jubilotheque.upmc.fr/img-viewer/fonds-bulsgf/GB_000078_001/Contenu/JPEG_HD/viewer.html?ns=GB_000078_001_J3_0590.jpg

- Roule L (1886) Nouvelle recherches sur les mollusques du terrain lacustre inférieur de Provence. *Annales de Malacologie* 2: 195–228. <http://www.biodiversitylibrary.org/page/16301303>
- Sandberger CLF (1870–1875) Die Land- und Süßwasser-Conchylien der Vorwelt. C. W. Kreidel, Wiesbaden, livr. 1–3: pp. 1–96, pl. 1–12 (1870); livr. 4–5: pp. 97–160, pl. 13–20 (1871); livr. 6–8: pp. 161–256, pl. 21–32 (1872); livr. 9–10: pp. 257–352, pl. 33–36 (1873); livr. 11: pp. i–viii + 353–1000 (1875). <https://doi.org/10.5962/bhl.title.112296>
- Salvador RB, Höltke O, Rasser MW, Kadolsky D (2016) Annotated type catalogue of the continental fossil gastropods in the Staatliches Museum für Naturkunde Stuttgart, Germany. *Palaeodiversity* 9: 15–70.
- Schileyko AA (1998) Treatise on Recent terrestrial pulmonate molluscs. Part 2. Gastrocoptidae, Hypselostomatidae, Vertiginidae, Truncatellinidae, Pachnodidae, Enidae, Sagdidae. *Ruthenica*, Supplement 2: 129–261.
- Schileyko AA (1999) Treatise on Recent terrestrial pulmonate Molluscs, Part 3: Partulidae, Aillyidae, Bulimulidae, Orthalicidae, Megaspiridae, Urocoptidae. *Ruthenica*, Supplement 2: 263–436.
- Schileyko AA (2000) Treatise on Recent terrestrial pulmonate molluscs. Part 6. Rhytididae, Chlamydephoridae, Systrophiidae, Haplotrematidae, Streptaxidae, Spiraxidae, Oleaciniidae, Testacellidae. *Ruthenica*, Supplement 2: 731–880.
- Schütt H (1991) Fossile Mollusken dreier anatolischer Ovas. *Archiv für Molluskenkunde* 120 (4/6): 131–147. <https://doi.org/10.1127/arch.moll/120/1991/131>
- Schütt H (1997) Altpleistozäne Mollusken im Becken von Pasinler bei Erzurum, Türkei. *Geologica et Palaeontologica* 31: 275–284.
- Simone LRL (2006) Land and Freshwater Molluscs of Brazil. EGB, Fapesp, São Paulo, 390 pp.
- Simone LRL, Casati R (2013) New land mollusk fauna from Serra da Capivara, Piauí, Brazil, with a new genus and five new species (Gastropoda: Orthalicoidea, Streptaxidae, Subuliniidae). *Zootaxa* 3683 (2): 145–158. <https://doi.org/10.11646/zootaxa.3683.2.4>
- Steklov AA (1966) Nazemnyye mollyuski neogena Predkavkaz'ya i ikh stratigraficheskoye znachenije. Akademia Nauka S.S.S.R., Moskva, 262 pp.
- Stworzewicz E (1995) Miocene land snails from Belchatów (Central Poland), I. Cyclophoridae, Pomatiasidae (Gastropoda Prosobranchia). *Paläontologische Zeitschrift* 69 (1/2): 19–30. <https://doi.org/10.1007/BF02985971>
- Tausch Lv (1886) Über die Fauna der nicht-marinen Ablagerungen der oberen Kreide des Csingerthales bei Ajka im Bakony (Veszprimer Comitatus, Ungarn) und über einige Conchylien der Gosaumergel von Aigen bei Salzburg. *Abhandlungen der k. k. Geologischen Reichsanstalt* 12: 1–32. http://opac.geologie.ac.at/wwwopacx/wwwopac.ashx?command=getcontent&server=images&value=Abh_Bd_12_1.pdf
- Thompson FG, Correa S (1991) Mexican land snails of the genus *Hendersoniella*. *Bulletin of the Florida Museum of Natural History* 36 (1): 1–23. <https://ufdc.ufl.edu/UF00095826/00001>
- Uit de Weerd DR, Gittenberger E (2013) Phylogeny of the land snail family Clausiliidae (Gastropoda: Pulmonata). *Molecular Phylogenetics and Evolution* 67: 201–216. <https://doi.org/10.1016/j.ympev.2013.01.011>

- Vermeulen JJ (1991) Notes on the non-marine molluscs of the island of Borneo 2. The genus *Opisthostoma* (Gastropoda Prosobranchia: Diplommatinidae), part 1. *Basteria* 55: 139–163.
- Webster NB, Van Dooren TJ, Schilthuizen M (2012) Phylogenetic reconstruction and shell evolution of the Diplommatinidae (Gastropoda: Caenogastropoda). *Molecular Phylogenetics and Evolution* 63(3): 625–638. <https://doi.org/10.1016/j.ympev.2012.02.004>
- Wenz W (1923–1930) Fossilium Catalogus I: Animalia. Gastropoda extramarina tertiaria. W. Junk, Berlin. Vol. I: 1–352 pp. (1923), Vol. II: 353–736 pp. (1923), Vol. III: 737–1068 pp. (1923), Vol. IV: 1069–1420 pp. (1923), Vol. V: 1421–1734 pp. (1923), Vol. VI: 1735–1862 pp. (1923), Vol. VII: 1863–2230 pp. (1926), Vol. VIII: 2231–2502 pp. (1928), Vol. IX: 2503–2886 pp. (1929), Vol. X: 2887–3014 pp. (1929), Vol. XI: 3015–3387 pp. (1930).
- Wenz W (1938–1944) Gastropoda. Teil 1: Allgemeiner Teil und Prosobranchia. xii + 1639 pp. In: Schindewolf OH (Ed.) *Handbuch der Paläozoologie*, Band 6. Bornträger, Berlin. Lief. 1, 1–240 [March 1938]; 3, 241–480 [October 1938]; 4, 481–720 [July 1939]; 6, 721–960 [August 1940]; 7, 961–1200 [October 1941]; 8, 1201–1506 [October 1943]; 9, 1507–1639, i–xii. [November 1944]
- Wenz W (1940) Die ältesten Stylommatophoren des europäischen Raumes. *Archiv für Molluskenkunde* 72(5/6): 129–144. https://www.zobodat.at/pdf/Archiv-fuer-Molluskenkunde_72_0129-0144.pdf
- Zallot E, Groenenberg DSJ, De Mattia W, Fehér Z, Gittenberger E (2015) Genera, subgenera and species of the Cochlostomatidae (Gastropoda, Caenogastropoda, Cochlostomatidae). *Basteria* 78 (4–6): 63–88.

Supplementary material I

Table S1

Authors: Barna Páll-Gergely, Thomas A. Neubauer

Data type: occurrence data

Explanation note: Geographic and stratigraphic occurrence data for species of *Ferussina*.

Provided are approximate GPS coordinates of the localities, alternative spellings of locality names, indications of type localities and literature.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.918.50135.suppl1>