

RESEARCH

Open Access

Modulation of respiratory dendritic cells during *Klebsiella pneumoniae* infection

Holger Hackstein^{1*}, Sabine Kranz¹, Anne Lippitsch¹, Andreas Wachtendorf¹, Olivia Kershaw², Achim D Gruber², Gabriela Michel¹, Jürgen Lohmeyer³, Gregor Bein¹, Nelli Baal¹ and Susanne Herold³

Abstract

Background: *Klebsiella pneumoniae* is a leading cause of severe hospital-acquired respiratory tract infections and death but little is known regarding the modulation of respiratory dendritic cell (DC) subsets. Plasmacytoid DC (pDC) are specialized type 1 interferon producing cells and considered to be classical mediators of antiviral immunity.

Method: By using multiparameter flow cytometry analysis we have analysed the modulation of respiratory DC subsets after intratracheal *Klebsiella pneumoniae* infection.

Results: Data indicate that pDCs and MoDC were markedly elevated in the post acute pneumonia phase when compared to mock-infected controls. Analysis of draining mediastinal lymph nodes revealed a rapid increase of activated CD103⁺ DC, CD11b⁺ DC and MoDC within 48 h post infection. Lung pDC identification during bacterial pneumonia was confirmed by extended phenotyping for 120G8, mPDCA-1 and Siglec-H expression and by demonstration of high Interferon-alpha producing capacity after cell sorting. Cytokine expression analysis of *ex vivo*-sorted respiratory DC subpopulations from infected animals revealed elevated Interferon-alpha in pDC, elevated IFN-gamma, IL-4 and IL-13 in CD103⁺ DC and IL-19 and IL-12p35 in CD11b⁺ DC subsets in comparison to CD11c⁺ MHC-class II^{low} cells indicating distinct functional roles. Antigen-specific naive CD4⁺ T cell stimulatory capacity of purified respiratory DC subsets was analysed in a model system with purified ovalbumin T cell receptor transgenic naive CD4⁺ responder T cells and respiratory DC subsets, pulsed with ovalbumin and matured with *Klebsiella pneumoniae* lysate. CD103⁺ DC and CD11b⁺ DC subsets represented the most potent naive CD4⁺ T helper cell activators.

Conclusion: These results provide novel insight into the activation of respiratory DC subsets during *Klebsiella pneumoniae* infection. The detection of increased respiratory pDC numbers in bacterial pneumonia may indicate possible novel pDC functions with respect to lung repair and regeneration.

Keywords: *Klebsiella pneumoniae*, Pneumonia, Plasmacytoid dendritic cells

Introduction

Klebsiella spp. are gram-negative bacteria and important opportunistic pathogens causing life-threatening nosocomial infections [1,2]. Typical clinical presentations of *Klebsiella pneumoniae* are nosocomial respiratory tract infections, urinary tract infections, infections of the bloodstream and premature infant intensive care unit infections [1,3-6]. Multidrug resistant *Klebsiella pneumoniae* strains are becoming an increasingly relevant

medical problem worldwide with limited clinical treatment options [7-11]. Therefore, there is a medical need for the development of novel therapeutic strategies in addition to classical antibiotic therapies. However, despite the clinical significance of *Klebsiella pneumoniae*, little is known regarding the modulation of respiratory dendritic cells (DC) subsets during lower respiratory tract infection with *Klebsiella spp.*

DC are central instigators of innate and adaptive immunity regulating both inflammation and activation of antigen-specific lymphocytes [12-14]. Based on the expression of different surface markers respiratory DC can be separated in at least four subsets, plasmacytoid DC

* Correspondence: Holger.Hackstein@immunologie.med.uni-giessen.de

¹Institute for Clinical Immunology and Transfusion Medicine, Justus-Liebig-University Giessen, Member of the German Center for Lung Research (DZL), Langhansstr. 7, D-35392, Giessen, Germany
Full list of author information is available at the end of the article

(pDC), CD103⁺ DC, CD11b^{hi} DC and monocytic DC (MoDC) [15-19]. pDC represent the most potent producers of Interferon-alpha and therefore play critical roles in antiviral immunity [20-23]. Additionally, it has been reported that pDC sense skin injury and promote wound healing through type I interferon suggesting a novel role for these professional antigen presenting cells [24].

Respiratory CD103⁺ DC have been reported to be involved in the development of airway hyperresponsiveness and to play important roles in naive CD8⁺ T cell activation and antiviral respiratory immunity [25,26]. With respect to bacterial respiratory tract infections, CD103⁺ DC have been shown to play a key role for α -Galactosylceramide-mediated protection against lethal *Streptococcus pneumoniae* infection [27]. Together with CD11b^{hi} DC and MoDC, CD103⁺ DC have been suggested to represent the major migratory DC subsets promoting naive CD4⁺ and CD8⁺ T cell activation in the draining lymph nodes [15,26,28-30].

We have used multicolor flow cytometry to dissect and quantitate respiratory DC subsets during murine *Klebsiella pneumoniae*. Additionally, respiratory DC subsets were purified by cell sorting from *Klebsiella*-infected animals and analysed for cytokine mRNA expression. The functional CD4⁺ naive T cell stimulatory capacity of purified respiratory DC subsets was determined *in vitro* after *Klebsiella pneumoniae* lysate stimulation of ovalbumin-pulsed DCs and ovalbumin (OVA) T-cell receptor (TCR)-transgenic naive CD4⁺ responder T cells.

Materials and methods

Mice and treatment protocol

Specific-pathogen-free C57BL/6 (C57BL/6NCr) and OVA TCR transgenic C57BL/6-Tg(TcraTcrb)425Cbn/J mice (18-21 g) were purchased from Charles River, Sulzfeld, Germany and maintained under specific-pathogen-free conditions. The transgenic mice express the TCR that pairs with the CD4 coreceptor and is specific for chicken OVA 323-339 in the context of MHC-class-II (I-A^b). Wild-type mice were infected intratracheally with *Klebsiella pneumoniae* serotype 2 (1×10^4 in sterile PBS in 50 μ l) as described [31] and analysed at the indicated timepoints. Mock-infected controls were treated identical but received sterile NaCl instead of *Klebsiella pneumoniae*. Analyses were performed after approval of the regional authority board City of Giessen (#71/2009 and # A25/2009).

Lung and mediastinal lymph node preparation

Lung single cell suspension were prepared after enzymatic digestion as described in detail elsewhere [16,17]. Absolute respiratory cell counts were enumerated with the trucount method (BD Biosciences, Germany) as described [16,17]. Trucount tubes contain a known number of fluorescent beads allowing the flow cytometer

software to calculate absolute cell counts. Single cell suspension from mediastinal lymph nodes (MLN) were minced and digested in RPMI 1640/10% FCS (PAA laboratories, Germany) with 10 U/ml DNase and 1 mg/ml Collagenase A (Roche, Germany) for 30 min at 37 C, resuspended with a 20 G 1/2 canule (0.9 \times 40 mm, BD, Germany), mashed through a 70 μ m cell strainer and washed two times with HBSS (7 min, 400 g; PAA laboratories, Germany).

Flow cytometry and fluorescent activated cell sorting

Cellular phenotyping was performed on a BD CantoII flow cytometer and fluorescent activated cell sorting was performed on a BD ARIA3 cell sorter (Becton Dickinson, San Jose, CA, USA). The following fluorochrome-labelled monoclonal antibodies conjugated to FITC, PE, PeCy7, PerCPCy5.5, APC, APC-Cy7, Pacific Blue and appropriate isotype controls were used for surface staining according to the manufacturer's instructions: CD11b, CD11c, CD45, CD64, CD86, CD103, CD274, I-A^b, GR-1, Ly-6G, F4/80, NK1.1, Fc ϵ R1 α (MAR-1), Siglec-H, Siglec-F (BD Biosciences, Germany), mPDCA-1 (MiltenyiBiotec, Germany), 120G8 (Dendritics, France). Surface mAb or isotype staining time was 30 min on ice and cells were washed with staining buffer (1 \times HBSS, PAA, Germany) at 400 g, 5 min, room temperature (RT) before analysis. The number of acquired events was $\geq 500,000$ after surface stainings.

Highly purified naive OVA TCR transgenic CD4⁺ T cells were prepared from spleen cell suspensions. First, MHC-class-II and CD19 positive cells were depleted with magnetic-beads (MiltenyiBiotec) on an AutoMACS magnetic-bead sorter (MiltenyiBiotec) and subsequently naive CD4⁺ T cells were sorted on the ARIA3 sorter after surface staining with a lineage cocktail (CD8a, CD25, CD11b, CD11c, CD45R, CD49b) and CD62L mAbs to identify naive T cells. The post sort purity of CD4⁺ CD62L⁺ CD44^{dim} cells was $>98\%$. Respiratory DC subsets were sorted on the ARIA3 cell sorter (purity $>98\%$; Additional file 1: Figure S1). Viability of cells was $>90\%$ as indicated by sytox blue (Life Technologies, Germany) staining. All mabs were ordered from Biolegend, Germany unless indicated otherwise.

Gating strategy for respiratory leukocyte subset discrimination

The gating strategy for the respiratory subsets, including respiratory DC subsets has been described recently in detail [17,19]. Briefly, respiratory leukocytes were identified by CD45 expression. Out of the CD45⁺ cells, neutrophils were identified by GR1^{bright}CD11b^{bright} expression. Subsequently, out of the neutrophil negative fraction, macrophages were identified as SiglecF⁺F4/80⁺ double positive cells. Respiratory dendritic cells (DC) were identified according to CD11c⁺Siglec-F^{neg} NK1.1^{neg} expression to

exclude autofluorescent macrophages and NK cells and further dissected into pDC ($120\text{ g}8^+ \text{ CD}11\text{b}^{\text{neg}}$), CD103 DC ($\text{CD}103^+ \text{ CD}11\text{b}^{\text{neg}}$) and CD11b DC ($\text{CD}11\text{b}^+ \text{ CD}103^{\text{neg}}$). MoDC were discriminated from CD11b DC based on CD64 and MAR1 expression as described recently [19]. In MLN, cells were additionally stained for CCR7 to identify migrating CD103 DC, CD11b DC and MoDC (Additional file 2: Figure S2). Fluorescence minus one (FMO) isotype controls were used regularly for proper identification of marker positive subsets.

Naive CD4+ T cell proliferation assay

Purified DC subsets (1×10^4 /well) were incubated in culture medium ($\pm 100\ \mu\text{g/ml}$ endotoxin-free OVA; Hyglos, Germany) with carboxyfluorescein diacetate succinimidyl ester (CFSE) labeled purified naive CD4⁺ OVA TCR transgenic responder cells (1×10^5 /well) in 96 well round bottom wells (Greiner, Germany) for five days. *Klebsiella pneumoniae* lysate was prepared as described [16] and added at a concentration of 10 ul/ml (1:100) 4 h after OVA for DC maturation. DC maturation with different *Klebsiella pneumoniae* lysate dilutions (1:10–1:500) was controlled by flow cytometry analysis of CD86 and MHC-class II expression on respiratory DC (Additional file 3: Figure S3). Culture medium consisted of RPMI 1640 with L-glutamine (PAA Laboratories), penicillin/streptomycin (PAN Biotech), sodium-pyruvate (Gibco), nonessential aminoacids (Sigma), Hepes buffer (Gibco), and 10% heat inactivated FCS (PAA Laboratories). CFSE labeling concentration was 1 μM and was performed according to the manufacturer instructions (Vybrant CFDA-Cell Tracer kit, Molecular probes, Eugene, USA). Positive controls consisted of *in vitro* bone-marrow-derived DC (expanded as described [32]) with OVA and responder cells. Negative controls consisted of responder cells cultured with OVA in the absence of stimulator cells and responder cells cultured with stimulator cells in the absence of ovalbumin. Proliferation was quantitated by measurement of CFSE-dilution in responder cells by flow cytometry as described [33]. Positive bone-marrow-derived control DC pulsed with OVA induced naive CD4⁺ responder proliferation rates of >80%. Proliferation rates in negative controls were < 5%.

IFN- α induction and quantitation

IFN- α production was stimulated with CpG ODN2216 (6 $\mu\text{g/ml}$) in sorted cell subsets after overnight culture in RPMI 1640 with 10% FCS (PAA Laboratories). IFN- α was quantitated in cellular supernatants on the FACS CantoII flow cytometer in parallel to recombinant cytokine standards by using the cytometric bead assay technology according to the manufacturer instructions (Flowcytomix, eBioscience, Germany).

Cytokine mRNA quantification

Cytokine mRNA expression was determined in sorted respiratory DC subsets from *Klebsiella*-infected animals by quantitative RT-PCR on an ABI TaqMan Step-One plus real time PCR system (Applied Biosystems, Germany) with the RT² profiler PCR assay (Qiagen, Germany). RNA was isolated with RNeasy Mini Kit (Qiagen) and transcribed into cDNA with the RT² Pre-amp cDNA Synthesis Kit (Qiagen) according to the manufacturer instructions.

Statistical analyses

The significance of differences between groups were analysed by one-way ANOVA and Tukey post-test for multiple comparisons. A p-value < 0.05 was considered significant. Statistical analyses were performed with Prism 5.02 software (Graphpad software, Inc., USA).


Results

Respiratory dendritic cell subsets and *Klebsiella pneumoniae* infection

By using multiparameter flow cytometry we analysed absolute and relative changes of major respiratory leukocyte populations including neutrophils, macrophages and DC during acute (48 h p.i.) and post acute (d5 p.i.) respiratory *Klebsiella pneumoniae* infection (Figure 1A). Total DC were further dissected by flow cytometry into four major subsets, MoDC, CD11b⁺ DC, CD103⁺ DC and pDC and relative DC subset frequencies and absolute DC subset numbers were calculated (Figure 1B,C; Figure 2). Results indicated that total respiratory DCs in general were increased both at 48 h p.i. and d5 p.i. but the dissection of DC subsets revealed different patterns for each subpopulation (Figure 1B,C). Early after infection (48 h p.i.), relative CD103⁺ DC numbers were significantly decreased in accordance with their reported function to rapidly transport antigen to regional lymph nodes [34]. Unexpectedly, with respect to pDC representing the rarest subset, we found a marked relative and absolute elevation of respiratory pDC numbers in the post acute phase of *Klebsiella pneumoniae* (d5 p.i.) when compared to mock-infected controls ($p < 0.01$; Figure 1B,C). Additionally, absolute numbers of respiratory CD11b⁺ DC and MoDC were markedly increased at 48 h and day 5 p.i. (Figure 1C). The massive leukocyte subset changes detected by flow cytometry correlated well with histopathology exhibiting extensive cellular infiltrates and cellular necrosis indicating severe bacterial pneumonia (Figure 3).

Confirmation of functional respiratory pDC during *Klebsiella pneumoniae* infection by extended phenotyping and cell sorting



Given the fact, that *Klebsiella pneumoniae* infection induced major respiratory inflammation we discussed the



possibility that pDC identification by flow cytometry might have been confounded by unspecific surface marker upregulation due to the inflammatory process. Accordingly, we extended pDC surface marker analysis on Klebsiella-infected animals (d5 p.i.) and included Siglec-H and mPDCA-1 mAb surface staining (Figure 4A). These experiments confirmed expression of Siglec-H and mPDCA-1 on respiratory pDC during Klebsiella pneumonia. Additionally, in order to use an independent functional method we sorted lung 120G8⁺ CD11b⁻ CD11c⁺ CD45⁺ putative respiratory pDC (d5 p.i.) and tested their capacity to produce IFN- α . These experiments confirmed high IFN- α producing capacity in respiratory pDC from Klebsiella-infected animals in comparison to non-pDC control fractions (Figure 4B). These data demonstrated that the immunophenotypically identified pDC during bacterial pneumonia, represented viable and functional respiratory pDC.

Distinct cytokine expression of purified respiratory DC subsets during Klebsiella pneumonia infection


To determine the cytokine expression of respiratory DC subsets during Klebsiella pneumonia we sorted pDC, CD103⁺ DC, CD11b⁺ DC and CD11c⁺ MHC-class II^{low} cells from Klebsiella-infected animals (d5 p.i.) to high purity (>98%) and analysed mRNA expression by quantitative real time RT-PCR. Controls included genomic DNA controls, reverse transcriptase controls and positive PCR controls. Relative cytokine expression of pDC, CD103⁺ DC and CD11b⁺ DC in comparison to CD11c⁺ MHC-class II^{low} cells was calculated by the $2^{-\Delta\Delta CT}$ method to indicate differences between the subsets (Figure 5). CD11c⁺ MHC-class II^{low} cells were selected as common reference because we wanted to display relative cytokine expression differences in-between pDC, CD103⁺ DC and CD11b⁺ DC. Respiratory pDC exhibited elevated IFN- α expression indicating activation during bacterial pneumonia. In contrast, CD103⁺ DC



exhibited increased IFN- γ , IL-4 and IL-13 expression, whereas CD11b DC showed elevated expression of IL-12p35 and IL-19 in comparison to CD11c⁺ MHC-class II^{low} cells.

Different modulation of MHC class II, CD86 and CD274 expression on respiratory DC subsets during acute *Klebsiella pneumoniae* infection

Expression of MHC-class II, CD86 and CD274 represent functionally important molecules of antigen presenting cells to regulate CD4⁺ T responder cell activation. Accordingly, we quantitated MHC-class II, CD86 and CD274 surface expression on total respiratory DC and DC subsets to determine the modulation during acute and post-acute respiratory *Klebsiella pneumoniae* infection (Figure 6A,B). Results demonstrated that at 48 h p.i. and at d5 p.i. respiratory MoDC and to a lesser extent CD11b⁺ DC from *Klebsiella pneumoniae*-infected animals exhibited reduced MHC-class II expression. In contrast, costimulatory CD86 expression was markedly increased at 48 h p.i. on respiratory CD103⁺ DC and CD11b⁺ DC of infected animals when compared to mock-infected-controls. pDC and MoDC did not exhibit major CD86 expression changes at 48 h p.i. Moreover, MoDC exhibited at d5 p.i. reduced CD86 expression





which may be related to the differentiation of newly arrived DC. The immunoregulatory molecule CD274 was most markedly induced on CD11b⁺ DC and MoDC both at 48 h p.i. and d5 p.i. whereas expression on CD103⁺ DC and pDC was less affected. These results indicated, that respiratory DC subsets exhibited major heterogeneity with respect to MHC-class II, CD86 and CD274 expression during respiratory Klebsiella infection suggesting different functional capacity to activate responder T cells. These differences would have been masked, if respiratory DC were analysed as a single

CD11c⁺ Siglec F^{neg} MHC-II⁺ population without subset discrimination.

Klebsiella pneumoniae-activated respiratory DC subsets exhibit different naive CD4⁺ T cell stimulatory capacity

In order to determine the naive CD4⁺ T cell stimulatory capacity of purified respiratory DC subsets activated by Klebsiella pneumonia we used an *in vitro* model system with CFSE-labelled OVA TCR transgenic CD4⁺ T helper cells as antigen-specific responder cells. Stimulatory respiratory DC subsets were sorted to high purity (>98% for





each subset) labelled with the model antigen OVA and matured with Klebsiella pneumonia lysate. Additional controls included sorted respiratory macrophages and granulocytes and *in vitro* bone-marrow-derived DC without OVA (negative control). Untouched naive CD4⁺ TCR transgenic responder cells were magnetic-bead sorted and the phenotype after sorting was controlled by surface CD4, CD62L and CD44 staining (purity >98%). Results indicated that Klebsiella pneumonia-activated CD103⁺ DC and CD11b⁺ DC subsets represented the most efficient stimulators of purified naive CD4 T responder cells (Figure 7; mean proliferation rates 95% and 64%, respectively). In contrast, Klebsiella pneumonia-activated pDC and CD11c⁺ MHC-class II^{low} cells, as well as granulocytes and macrophages induced no or only minor naive CD4 T cell proliferation (< 5%). Interestingly, sorted DC subsets that were not activated with Klebsiella pneumonia induced comparable CD4⁺ T cell proliferation indicating that Klebsiella may induce both activatory and inhibitory effects on DC. Control experiments with cultures of DC + CD4⁺ T cells without OVA and CD4⁺ T cells + OVA without DC showed no responder cell proliferation and confirmed the antigen-specificity of the assay (Figure 7A,B). Naive CD4⁺ T cell stimulatory capacity of purified, Klebsiella activated respiratory DC subsets correlated closely with MHC-class II and CD86 expression of respiratory DC

subsets from Klebsiella-infected animals (Figure 7) underlining the functional importance of MHC-class II and CD86 expression. Overall, these results indicated marked differences of Klebsiella-activated respiratory DC subsets to activate naive CD4⁺ T cells.

Rapid accumulation of activated CD103⁺ DC, CD11b⁺ DC and MoDC in the MLN

Analysis of DC subset changes in the draining MLN revealed a marked increase of relative and absolute CCR7⁺ CD103⁺DC, CD11b⁺ DC and MoDC numbers within 48 h p.i. whereas absolute pDC numbers were elevated on D5 p.i. (Figure 8A,B). Phenotypic analysis of activatory CD86 expression indicated significant upregulation on CD103⁺ DC and CD11b⁺ DC in contrast to pDC and MoDC (Figure 8 C). Interestingly, MHC-class II expression was impaired to different extent on all DC subsets in comparison to mock-infected controls again suggesting that Klebsiella pneumonia infection may induce both activatory and inhibitory effects on antigen presenting cells.

Discussion

Klebsiella pneumoniae is a frequent multidrug-resistant emerging pathogen playing an increasingly important role in severe nosocomial infections [7,8,10,11]. DC are central regulators of innate and adaptive immunity against





Figure 7 Different capacity of purified *Klebsiella pneumoniae*-stimulated respiratory DC subsets to activate OVA TCR-transgenic naive CD4⁺ responder T cells in a CFSE-dilution assay. Respiratory DC subsets were sorted to high purity (>98%), pulsed with the model antigen OVA and matured with *Klebsiella pneumoniae* lysate (Kp). Naive CD4⁺ responder T cells were sorted from OVA TCR transgenic animals and labeled with CFSE. Negative controls included DC without *Klebsiella* lysate (control), responder T cells cultured without DC (no DC) and DC-T cell co-cultures without OVA (no OVA). Proliferation was quantitated by CFSE-dilution of responder T cells (A,B). Mean ± SEM; n ≥ 3.

different kinds of pathogens [35,36,37]. Based on different developmental pathways, surface markers and tissue localizations, DC can be divided into distinct subsets exhibiting specific functions [38]. By using multiparameter flow cytometry and cell sorting we have determined the number and activation of respiratory DC subsets during experimental *Klebsiella pneumoniae* infection.

Subset-specific respiratory DC quantification of *Klebsiella*-infected animals showed that the rare pDC population were markedly elevated in the post acute disease phase. We confirmed the identity of functional respiratory pDC during bacterial pneumonia by extended phenotyping and demonstration of large IFN- α production capacity after cell sorting. Moreover, elevated IFN- α mRNA expression was demonstrated in purified respiratory pDC sorted from *Klebsiella*-infected

animals. Unexpectedly, cytokine mRNA analysis indicated IL-4 and IL-13 expression in CD103 DC and IL-19 expression in CD11b DC. Due to the rather low cytokine mRNA expression, marked pulmonary inflammation and the technical possibility of contaminating innate lymphoid cells, further detailed studies using combinations of different methods are required to dissect the cytokine producing capacity of respiratory DC subsets during gramnegative bacterial pneumonia.

The accumulation of functional respiratory pDC during bacterial pneumonia has to the best of our knowledge not been reported so far and raises the question regarding potential novel functions of pDC in bacterial pneumonia. Respiratory pDC may play a role in tissue repair. Gregorio et al. recently reported, that pDC accumulate in large numbers in injured skin areas and express



IFN- α [24]. The expression of IFN- α was shown to be related to the detection of self nucleic-acids that were released due to the skin injury. Additionally, it was demonstrated that pDC depletion significantly delayed wound healing suggesting a functional role for these rare DC subset in tissue repair. Alternatively, accumulation of pDC in the post acute bacterial pneumonia phase may be related to immunoregulatory functions of these cells. pDC have the capacity to control both CD4⁺ and CD8⁺ T cell activation [39]. Antigen presentation to CD4 T cells via pDC has been described to inhibit T helper 1 and T helper 17 immune responses, suggesting a possible role to prevent T helper-dependent automimmunity [40]. In a different study using a conditional plasmacytoid depletion model, pDC were demonstrated to play an important role in the stimulation of antiviral CD8⁺ T cell responses [41].

Our results indicate that Klebsiella pneumonia may exhibit additionally inhibitory effects on DC. The *ex vivo* analysis of MHC-class II expression on lung DC subsets of Klebsiella pneumonia infected animals consistently indicated decreased MHC-class II expression in comparison to mock-infected controls. This may be related to the recruitment or differentiation of newly arrived DC.

However, analysis of CCR7 migrating DC in the draining MLN during infection additionally indicated decreased MHC-class II expression suggesting Klebsiella pneumonia-mediated suppression. Additionally, analysis of the inhibitory molecule CD274 on lung DC during Klebsiella infection revealed a massive upregulation on CD11b DC and MoDC and to a lesser extent on CD103 DC suggesting another pathway of Klebsiella pneumonia-mediated suppression. Klebsiella-mediated induction of inhibitory molecule CD274 on lung DC may be important in limiting antibacterial effector T cell responses and promoting infection-associated immunosuppression [42,43]. In this context it would be interesting to establish technologies allowing precise DC subset specific enumeration of bacterial load. Sorting strategies are limited due to a number of technical factors, such as passive bacteria attachment, complex DC phenotype, low DC frequency. Therefore, development of other technologies, such as combination of specific staining of bacterial-related gene products with DC subset markers might be more successful [44].

Our results using ovalbumin-pulsed, Klebsiella pneumoniae-stimulated purified pDC as stimulators of TCR transgenic naive CD4⁺ T cells confirmed the finding, that pDC

are not efficient CD4⁺ T cell stimulators. In direct comparison to other respiratory DC subsets, Klebsiella pneumonia-stimulated CD103⁺ DC and CD11b DC represented the most efficient naïve CD4⁺ T cell activators. Moreover, MHC-class II and CD86 surface expression, representing important functional molecules required for MHC-class II restricted CD4 activation were markedly reduced on respiratory pDC from Klebsiella-infected animals compared to either CD103⁺ DC and CD11b DC. Additionally, analysis of draining MLN during infection revealed rapid relative and absolute accumulation of CD103 DC, CD11b DC and MoDC numbers within 48 h p.i. whereas pDC numbers did exhibit only an absolute increase at d5 p.i. indicating a different role for T cell activation.

In summary, the present study has revealed marked quantitative and qualitative differences for respiratory DC subsets during sublethal respiratory Klebsiella pneumonia infection. Both the *in vivo* as well as the *in vitro* data underline the importance of differentiating respiratory DC subsets during Klebsiella pneumonia. The analysis of lung DC as a single cell population during infection would have masked many of these differences. Our study highlights that these lung subsets exhibit different migration kinetics, different activatory/regulatory surface molecule expression and different functional CD4 activatory capacity. With respect to the novel finding of increased pDC during Klebsiella infection, future studies are necessary to address the functional role of pDC during bacterial pneumonia. The use of *in vivo* DC subset depletion models in future studies will provide additional insight into the relevance of DC subsets for CD4 and CD8 T effector cell expansion during Klebsiella pneumonia infection.

Additional files

Additional file 1: Figure S1. Gating strategy and post-sort controls for pDC, MHC-II^{low} CD11c⁺ cells, CD103 DC and CD11b DC. Dotplot staining of sorting gates (A) and post-sort controls with % cell purity (B) for pDC, MHC-II^{low} CD11c⁺ cells, CD103 DC and CD11b DC.

Additional file 2: Figure S2. Dissection of DC subsets in the MLN during Klebsiella pneumonia infection. Dotplot staining and gating of MLN CD103⁺ DC, CD11b⁺ DC, MoDC and pDC subsets in Klebsiella-infected animals and mock-infected controls. Migratory CD11c⁺ CD103⁺ DC, CD11b⁺ DC and MoDC were identified as MHC-class II⁺ CCR7⁺ cells and discriminated according to differential CD103, CD11b, MAR1 and CD64 expression. MLN pDC were identified as CD11c⁺ 120G8⁺ CD11b^{neg} cells. Fluorescence minus one controls (FMO) were used for proper identification of CCR7⁺ and CD64⁺ populations.

Additional file 3: Figure S3. Maturation control for Klebsiella pneumonia lysate. Lung homogenate was stimulated with indicated concentrations of Kp lysate and median fluorescence intensity (MFI) for CD86 and MHC-II expression was quantitated after 24 h on CD11c⁺ Siglec F^{neg} NK1.1^{neg} DC by flow cytometry. Mean ± SEM; n ≥ 3.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

HH, NB, JL, GB and SH designed the study. HH drafted the manuscript and performed the statistical analysis. NB, AW and SK performed animal experiments. GM, AL and NB performed *in vitro* immunoassays and expression analysis. OK and AG performed the histopathological analysis. All authors read and approve the final manuscript.

Acknowledgements

The work was supported by the SFB Transregio 84, German Science Foundation, Innate Immunity of the Lung: Mechanisms of Pathogen Attack and Host Defence in Pneumonia (SFB TR84, Project B3 HH). Additional support was provided by UGMLC (University Gießen and Marburg Lung Center) and Excellence Cluster Cardiopulmonary System (ECCPS). We thank Ms Rebecca Winkler for excellent technical support.

Author details

¹Institute for Clinical Immunology and Transfusion Medicine, Justus-Liebig-University Giessen, Member of the German Center for Lung Research (DZL), Langhansstr. 7, D-35392, Giessen, Germany. ²Department of Veterinary Pathology, Freie Universität Berlin, Robert-von-Ostertag-Str. 15, 14163, Berlin, Germany. ³Department of Internal Medicine II, Klinikstrasse 33, Berlin, Germany.

Received: 4 February 2013 Accepted: 13 September 2013

Published: 17 September 2013

References

- Ahmad TA, El-Sayed LH, Haroun M, Hussein AA, El Ashry ESH: Development of immunization trials against Klebsiella pneumoniae. *Vaccine* 2012, **30**:2411–2420.
- Livermore DM: Current epidemiology and growing resistance of gram-negative pathogens. *Korean J Intern Med* 2012, **27**:128–142.
- Gupta A: Hospital-acquired infections in the neonatal intensive care unit—Klebsiella pneumoniae. *Semin Perinatol* 2002, **26**:340–345.
- Matsen JM: The sources of hospital infection. *Medicine (Baltimore)* 1973, **52**:271–277.
- Jarvis WR, Munn VP, Highsmith AK, Culver DH, Hughes JM: The epidemiology of nosocomial infections caused by Klebsiella pneumoniae. *Infect Control* 1985, **6**:68–74.
- Yankov IV, Shmylev TI: Ventilator-associated pneumonias in children (I)—diagnostic criteria, etiology and pathogenesis. *Folia Med (Plovdiv)* 2012, **54**:5–11.
- Arnold RS, Thom KA, Sharma S, Phillips M, Kristie Johnson J, Morgan DJ: Emergence of Klebsiella pneumoniae carbapenemase-producing bacteria. *South Med J* 2011, **104**:40–45.
- Rapp RP, Urban C: Klebsiella pneumoniae carbapenemases in Enterobacteriaceae: history, evolution, and microbiology concerns. *Pharmacotherapy* 2012, **32**:399–407.
- da Silva RM, Traebert J, Galato D: Klebsiella pneumoniae carbapenemase (KPC)-producing Klebsiella pneumoniae: a review of epidemiological and clinical aspects. *Expert Opin Biol Ther* 2012, **12**:663–671.
- Chong Y, Ito Y, Kamimura T: Genetic evolution and clinical impact in extended-spectrum beta-lactamase-producing Escherichia coli and Klebsiella pneumoniae. *Infect Genet Evol* 2011, **11**:1499–1504.
- Nordmann P, Cuzon G, Naas T: The real threat of Klebsiella pneumoniae carbapenemase-producing bacteria. *Lancet Infect Dis* 2009, **9**:228–236.
- Banchereau J, Steinman RM: Dendritic cells and the control of immunity. *Nature* 1998, **392**:245–252.
- Lambrecht BN, Hammad H: Lung dendritic cells in respiratory viral infection and asthma: from protection to immunopathology. *Annu Rev Immunol* 2012, **30**:243–270.
- Guilliams M, Lambrecht BN, Hammad H: Division of labor between lung dendritic cells and macrophages in the defense against pulmonary infections. *Mucosal Immunol* 2013, **6**:464–473.
- Kim T, Braciale T: Respiratory dendritic cell subsets differ in their capacity to support the induction of virus-specific cytotoxic CD8⁺ T cell responses. *PLoS One* 2009, **4**:e4204.
- Hackstein H, Hagel N, Knoche A, Kranz S, Lohmeyer J, Wulffen W, von Kershaw O, Gruber AD, Bein G, Baal N: Skin TLR7 triggering promotes accumulation of respiratory dendritic cells and natural killer cells. *PLoS One* 2012, **7**:e43320.

17. Hackstein H, Wachtendorf A, Kranz S, Lohmeyer J, Bein G, Baal N: **Heterogeneity of respiratory dendritic cell subsets and lymphocyte populations in inbred mouse strains.** *Respir Res* 2012, **13**:94.
18. Unkel B, Hoegner K, Clausen BE, Lewe-Schlösser P, Bodner J, Gattenloehner S, Janssen H, Seeger W, Lohmeyer J, Herold S: **Alveolar epithelial cells orchestrate DC function in murine viral pneumonia.** *J Clin Invest* 2012, **122**:3652–3664.
19. Plantinga M, Guillems M, Vanheerswyngheles M, Deswarte K, Branco-Madeira F, Toussaint W, Vanhoutte L, Neyt K, Killeen N, Malissen B, Hammad H, Lambrecht BN: **Conventional and monocyte-derived CD11b(+) dendritic cells initiate and maintain T helper 2 cell-mediated immunity to house dust mite allergen.** *Immunity* 2013, **38**:322–335.
20. Siegal FP, Kadovaki N, Shodell M, Fitzgerald-Bocarsly PA, Shah K, Ho S, Antonenko S, Liu YJ: **The nature of the principal type 1 interferon-producing cells in human blood.** *Science* 1999, **284**:1835–1837.
21. Gilliet M, Cao W, Liu Y: **Plasmacytoid dendritic cells: sensing nucleic acids in viral infection and autoimmune diseases.** *Nat Rev Immunol* 2008, **8**:594–606.
22. Szabo G, Dolganiuc A: **The role of plasmacytoid dendritic cell-derived IFN alpha in antiviral immunity.** *Crit Rev Immunol* 2008, **28**:61–94.
23. Bjorck P: **The multifaceted murine plasmacytoid dendritic cell.** *Hum Immunol* 2002, **63**:1094–1102.
24. Gregorio J, Meller S, Conrad C, Di Nardo A, Homey B, Lauerma A, Arai N, Gallo RL, Digiovanni J, Gilliet M: **Plasmacytoid dendritic cells sense skin injury and promote wound healing through type I interferons.** *J Exp Med* 2010, **207**:2921–2930.
25. Sung S, Fu S, Rose C, Gaskin F, Ju S, Beaty S: **A major lung CD103 (alphaE)-beta7 integrin-positive epithelial dendritic cell population expressing Langerin and tight junction proteins.** *J Immunol* 2006, **176**:2161–2172.
26. Ho AWS, Prabhu N, Betts RJ, Ge MQ, Dai X, Hutchinson PE, Lew FC, Wong KL, Hanson BJ, Macary PA, Kemeny DM: **Lung CD103+ dendritic cells efficiently transport influenza virus to the lymph node and load viral antigen onto MHC class I for presentation to CD8 T cells.** *J Immunol* 2011, **187**:6011–6021.
27. Ivanov S, Fontaine J, Paget C, Macho Fernandez E, van Maele L, Renneson J, Maillet I, Wolf NM, Rial A, Leger H, Ryffel B, Frisch B, Chabalgoity JA, Sirard JC, Benecke A, Faveeuw C, Trottein F: **Key role for respiratory CD103(+) dendritic cells, IFN-gamma, and IL-17 in protection against Streptococcus pneumoniae infection in response to alpha-galactosylceramide.** *J Infect Dis* 2012, **206**:723–734.
28. Le B, Etchart N, Goubier A, Lira S, Sirard J, van Rooijen N, Caux C, Ait-Yahia S, Vicari A, Kaiserlian D, Dubois B: **Dendritic cells rapidly recruited into epithelial tissues via CCR6/CCL20 are responsible for CD8+ T cell crosspriming in vivo.** *Immunity* 2006, **24**:191–201.
29. Leon B, Lopez-Bravo M, Ardavin C: **Monocyte-derived dendritic cells formed at the infection site control the induction of protective T helper 1 responses against Leishmania.** *Immunity* 2007, **26**:519–531.
30. Furuhashi K, Suda T, Hasegawa H, Suzuki Y, Hashimoto D, Enomoto N, Fujisawa T, Nakamura Y, Inui N, Shibata K, Nakamura H, Chida K: **Mouse Lung CD103+ and CD11bhigh dendritic cells preferentially induce distinct CD4+ T cell responses.** *Am J Respir Cell Mol Biol* 2012, **46**:165–172.
31. Herold S, Tabar TS, Janssen H, Hoegner K, Cabanski M, Lewe-Schlösser P, Albrecht J, Driever F, Vadasz I, Seeger W, Steinmueller M, Lohmeyer J: **Exudate macrophages attenuate lung injury by the release of IL-1 receptor antagonist in gram-negative pneumonia.** *Am J Respir Crit Care Med* 2011, **183**:1380–1390.
32. Hackstein H, Taner T, Zahorchak AF, Morelli AE, Logar AJ, Gessner A, Thomson AW: **Rapamycin inhibits IL-4-induced dendritic cell maturation in vitro and dendritic cell mobilization and function in vivo.** *Blood* 2003, **101**:4457–4463.
33. Hackstein H, Knoche A, Nockher A, Poeling J, Kubin T, Jurk M, Vollmer J, Bein G: **The TLR7/8 ligand resiquimod targets monocyte-derived dendritic cell differentiation via TLR8 and augments functional dendritic cell generation.** *Cell Immunol* 2011, **271**:401–412.
34. Ho AWS, Prabhu N, Betts RJ, Ge MQ, Dai X, Hutchinson PE, Lew FC, Wong KL, Hanson BJ, Macary PA, Kemeny DM: **Lung CD103+ dendritic cells efficiently transport influenza virus to the lymph node and load viral antigen onto MHC class I for presentation to CD8 T cells.** *J Immunol* 2011, **187**:6011–6021.
35. Terrazas CA, Terrazas LI, Gomez-Garcia L: **Modulation of dendritic cell responses by parasites: a common strategy to survive.** *J Biomed Biotechnol* 2010, **2010**:357106.
36. Bedoui S, Kupz A, Wijburg OL, Walduck AK, Rescigno M, Strugnell RA: **Different bacterial pathogens, different strategies, yet the aim is the same: evasion of intestinal dendritic cell recognition.** *J Immunol* 2010, **184**:2237–2242.
37. Liu B, Woltman AM, Janssen HLA, Boonstra A: **Modulation of dendritic cell function by persistent viruses.** *J Leukoc Biol* 2009, **85**:205–214.
38. Kushwah R, Hu J: **Complexity of dendritic cell subsets and their function in the host immune system.** *Immunology* 2011, **133**:409–419.
39. Takagi H, Fukaya T, Eizumi K, Sato Y, Sato K, Shibasaki A, Otsuka H, Hijikata A, Watanabe T, Ohara O, Kaisho T, Malissen B, Sato K: **Plasmacytoid dendritic cells are crucial for the initiation of inflammation and T cell immunity in vivo.** *Immunity* 2011, **35**:958–971.
40. Loschko J, Heink S, Hackl D, Dudziak D, Reindl W, Korn T, Krug AB: **Antigen targeting to plasmacytoid dendritic cells via Siglec-H inhibits Th cell-dependent autoimmunity.** *J Immunol* 2011, **187**:6346–6356.
41. Swiecki M, Gilfillan S, Vermi W, Wang Y, Colonna M: **Plasmacytoid dendritic cell ablation impacts early interferon responses and antiviral NK and CD8(+) T cell accrual.** *Immunity* 2010, **33**:955–966.
42. Hotchkiss RS, Monneret G, Payen D: **Immunosuppression in sepsis: a novel understanding of the disorder and a new therapeutic approach.** *Lancet Infect Dis* 2013, **13**:260–268.
43. Wolfe SJ, Strebowski J, Bartz H, Sahr A, Arnold C, Kaiser C, Dalpke AH, Heeg K: **PD-L1 expression on tolerogenic APCs is controlled by STAT-3.** *Eur J Immunol* 2011, **41**:413–424.
44. Sha J, Rosenzweig JA, Kirtley ML, van Lier CJ, Fitts EC, Kozlova EV, Erova TE, Tiner BL, Chopra AK: **A non-invasive in vivo imaging system to study dissemination of bioluminescent Yersinia pestis CO92 in a mouse model of pneumonic plague.** *Microb Pathog* 2013, **55**:39–50.

doi:10.1186/1465-9921-14-91

Cite this article as: Hackstein et al.: Modulation of respiratory dendritic cells during Klebsiella pneumonia infection. *Respiratory Research* 2013 **14**:91.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

