



Holophytochrome-Interacting Proteins in *Physcomitrella*: Putative Actors in Phytochrome Cytoplasmic Signaling

Anna Lena Ermert, Katharina Mailliet and Jon Hughes*

Institute for Plant Physiology, Justus Liebig University, Giessen, Germany

Phytochromes are the principle photoreceptors in light-regulated plant development, primarily acting via translocation of the light-activated photoreceptor into the nucleus and subsequent gene regulation. However, several independent lines of evidence indicate unambiguously that an additional cytoplasmic signaling mechanism must exist. Directional responses in filament tip cells of the moss *Physcomitrella patens* are steered by phy4 which has been shown to interact physically with the blue light receptor phototropin at the plasma membrane. This complex might perceive and transduce vectorial information leading to cytoskeleton reorganization and finally a directional growth response. We developed yeast two-hybrid procedures using photochemically functional, full-length phy4 as bait in *Physcomitrella* cDNA library screens and growth assays under different light conditions, revealing Pfr-dependent interactions possibly associated with phytochrome cytoplasmic signaling. Candidate proteins were then expressed *in planta* with fluorescent protein tags to determine their intracellular localization in darkness and red light. Of 14 candidates, 12 were confirmed to interact with phy4 *in planta* using bimolecular fluorescence complementation. We also used database information to study their expression patterns relative to those of phy4. We discuss the likely functional characteristics of these holophytochrome-interacting proteins (HIP's) and their possible roles in signaling.

Keywords: phytochrome, signal transduction, phototropin, *Physcomitrella patens*, yeast two-hybrid, split-YFP, bimolecular fluorescence complementation, protein-protein interaction

OPEN ACCESS

Edited by:

Enamul Huq,

The University of Texas at Austin, USA

Reviewed by:

Andreas Hiltbrunner,

University of Freiburg, Germany

Margaret Ahmad,

University of Paris, France

*Correspondence:

Jon Hughes

jon.hughes@uni-giessen.de

Specialty section:

This article was submitted to

Plant Cell Biology,

a section of the journal

Frontiers in Plant Science

Received: 01 January 2016

Accepted: 21 April 2016

Published: 12 May 2016

Citation:

Ermert AL, Mailliet K and Hughes J (2016) Holophytochrome-Interacting Proteins in *Physcomitrella*: Putative Actors in Phytochrome Cytoplasmic Signaling. *Front. Plant Sci.* 7:613. doi: 10.3389/fpls.2016.00613

Abbreviations: AD, GAL4 transcription factor activation domain; AFR, ATTENUATED FAR RED RESPONSE; B, blue; BD, GAL4 transcription factor DNA binding domain; BLAST, basic local alignment search tool; CDPK, calcium-dependent protein kinase; CDS, coding DNA sequence; CFK1, COP9 INTERACTING F-BOX KELCH 1; CFP, cyan fluorescent protein; CHUP1, CHLOROPLAST UNUSUAL POSITIONING; CO, CONSTANS; COP, CONSTITUTIVELY PHOTOMORPHOGENIC; D, dark; DSD, double synthetic dropout; EF1 α , translation elongation factor 1 α ; eIF-5A, eukaryotic translation initiation factor 5A; FBK, F-box/kelch; FHY1, FAR-RED-ELONGATED HYPOCOTYL 1; FKF1, FLAVIN-BINDING KELCH-REPEAT F-BOX 1; FP, fluorescent protein; FR, far-red; FT, FLOWERING LOCUS T (florigen); GFP, green fluorescent protein; GPA1, G-protein α -subunit; HIP, holophytochrome-interacting protein; HT1, HIGH LEAF TEMPERATURE 1; LKP2, LOV-KELCH PROTEIN 2; NES, nuclear export signal/sequence; NLS, nuclear localization signal/sequence; NPH3, NON-PHOTOTROPIC HYPOCOTYL 3; PCB, phycocyanobilin; Pfr, far-red absorbing form of phytochrome; PHL, PHYTOCHROME-DEPENDENT LATE-FLOWERING; phot, phototropin holoprotein; phy, phytochrome holoprotein; PIE, phytochrome-interacting factor; PKS1, phytochrome kinase substrate 1; PORA, protochlorophyllide oxidoreductase A; PP2A, PROTEIN PHOSPHATASE 2A; Pr, red absorbing form of phytochrome; PRL1, PLEIOTROPIC REGULATORY LOCUS 1; PRN1, pirin 1; QSD, quadruple synthetic dropout; R, Rc/Rp, red, continuous red, pulsed red; RBL, rhomboid-like protein; RFI2, RED AND FAR RED INSENSITIVE 2; RFP, red fluorescent protein; RPT2, ROOT PHOTOTROPISM 2; SCF, SKP1p-Cdc53p (cullin)-RBX1-F-box; SnRK, SNF1-related protein kinase; SOT, sulfotransferase; TSD, triple synthetic dropout; VOZ1, VASCULAR PLANT ONE ZF 1; W, white; Y2H, yeast two-hybrid; YFP, YFP_N/YFP_C, yellow fluorescent protein, N-terminal region, C-terminal region; YPDA, yeast peptone dextrose adenine; ZF, zinc finger.

INTRODUCTION

Plant phytochromes absorb predominantly red (R; ~660 nm) and far-red (FR; ~710–730 nm) light, thereby steering physiological responses including seed germination, de-etiolation, shade avoidance, and flowering. The R-absorbing Pr dark state is restricted to the cytoplasm and is physiologically inactive. In light it is converted to the physiologically active FR-absorbing Pfr state, which is then translocated from the cytoplasm to the nucleus where it regulates the transcription of numerous genes via direct interaction with master transcription factors such as the PIFs, leading to the long-term responses mentioned above. However, several physiological responses cannot be explained by gene regulation. Firstly, responses like R/FR-dependent ion fluxes (Tanada, 1968) or cytoplasmic streaming (Takagi et al., 2003) occur within seconds, far too fast for such a mechanism. Secondly, the directional responses in higher plants mediated by the blue light (B) absorbing photoreceptor phototropin are modulated by phytochromes: although this might derive from phytochrome effects on gene expression, several of these responses are still seen in mutants in which nuclear translocation is defective, implying that Pfr is also physiologically active in the cytoplasm (Rösler et al., 2007; Kami et al., 2012). Indeed, *Arabidopsis* phyA and phyB Pfr bind and phosphorylate PKS1, a cytoplasmic protein, whereas phyB regulates translation of PORA by Pfr-dependent binding to PENTA1, also a cytoplasmic protein (Fankhauser et al., 1999; Paik et al., 2012). The fact that phytochromes can connect to signal transduction systems independently of transcription/translation is even more obvious in bryophytes. Phototropic growth of moss filament tip cells is steered by R in a FR-reversible manner, indicating that the photoreceptor is phytochrome. This cannot result from transcription/translation regulation because directional information would inevitably be lost in the process. Furthermore, gene targeting in *Physcomitrella* and *Ceratodon* showed that the response is steered by a specific phytochrome, phy4 (Mittmann et al., 2004, 2009). Thus phy4 must be associated with a signaling mechanism which faithfully transmits the vectorial information given by the incident light. Indeed, we found that phy4 interacts with phototropin at the plasma membrane (Jaedicke et al., 2012), suggesting a physical link to the machinery controlling cytoskeletal organization and tip growth (Meske and Hartmann, 1995; Meske et al., 1996). Chloroplast translocation too is regulated vectorially by phy4 (Kadota et al., 2000; Mittmann et al., 2004). Interestingly, in both the fern *Adiantum* and the alga *Mougeotia* chloroplast movements are also steered vectorially in a R/FR-reversible manner, indicative of phytochrome action – but in these cases via neochrome, a chimeric photoreceptor comprising a phytochrome sensory module attached to a phototropin. Remarkably, transgenic neochrome in *Arabidopsis* rescues phototropism in phototropin-deficient mutants – but the response is then seen in both B and R (Nozue et al., 1998; Kanegae et al., 2006; Kanegae and Kimura, 2015). The field of phytochrome cytoplasmic signaling has been reviewed recently (Hughes, 2013).

It remains unclear, however, how phytochrome cytoplasmic signals are transmitted. Even assuming that the phytochrome

simply hijacks the phototropin-associated machinery is futile as little is known about phototropin signaling beyond NPH3 (Roberts et al., 2011). We therefore decided to search for *Physcomitrella* phy4 partners directly, describing our initial results here. We developed Y2H procedures using full-length, photochemically functional, R-activated phy4 holophytochrome as bait in cDNA library screens: to our knowledge this is the first report of such a procedure. The apparent interaction was checked with the full-length CDS model and R/FR reversibility tested. Candidate interactors were then investigated *in planta* regarding their intracellular localization in darkness (D) and R by fusing fluorescent tags to the N- and C-termini of the respective gene products. Finally *in planta* interaction with phy4 was investigated using bimolecular fluorescence complementation (split-YFP) methods. We thereby identified 14 putative holophytochrome interacting proteins (HIP's), 12 of which were confirmed in split-YFP. Their possible roles in phytochrome cytoplasmic signaling are discussed.

MATERIALS AND METHODS

Cloning and Y2H Procedures

The MatchMaker GAL4 Two-Hybrid System 3 (Clontech) was used to screen for potential cytoplasmic signaling partners of *Physcomitrella* phy4 (Pp3c27_7830V1.1). Full-length phy4 bait was cloned from first strand *Physcomitrella* cDNA into pGBKT7 and pBAC, a derivative of pBRIDGE (both Clontech), to yield hybrid proteins with the BD attached to the N- and C-termini (BD:phy4 and phy4:BD), respectively. A pre-transformed oligo dT-primed cDNA library in the prey vector pGADT7 was kindly provided by Hans Sommer (MPI Cologne) in yeast strain AH109. Single, double and screening transformations were performed as described (Agatep et al., 1998; Gietz and Woods, 2002). The library was either transformed with 40 µg BD:phy4 in a 40x scale or mated to Y187 yeast pre-transformed with phy4:BD. For mating, the bait strain was grown overnight, harvested and mixed in a 2:1 ratio with cells of the thawed library aliquot in 2x YPDA medium and incubated for 24 h at 40 rpm and 30°C. Transformation and mating mixtures were plated either with water or 0.5x YPDA on -Trp/-Leu/-His TSD medium with 2.5 mM 3-AT or with 0.25 mM 3-AT and 30 µM PCB, respectively. PCB was extracted from *Spirulina* and purified as described (Jaedicke et al., 2012). Plates were incubated for 14 days at 30°C in D or in 0.7 µmol m⁻² s⁻¹ Rc (660 nm LEDs). Fresh PCB was added 2–3 times during the incubation period. Positives were picked and transferred to a -Trp/-Leu DSD (double SD) masterplate and also further selected on TSD with 3-AT and on -Trp/-Leu/-His/-Ade QSD (quadruple SD) in the case of double transformation and TSD with PCB in case of library mating. DNA extracts were transformed into *E. coli*, the cloned pGADT7 DNA extracted, the inserts sequenced and finally identified by BLAST searches at NCBI¹ and Cosmoss². Based on EST and our own

¹www.ncbi.nlm.nih.gov

²www.cosmoss.org

sequencing data, the likely best gene model was selected and the corresponding amino acid sequences were BLASTed against *Arabidopsis thaliana* and *Viridiplantae* non-redundant protein databases at NCBI in order to identify putative homologs in higher plants.

AD:HIP4 and AD:HIP6 constructs were cloned by Phusion amplification of cDNA, subcloning into pCR bluntII TOPO (Invitrogen) and ClaI/SacI cloning into pGADT7. All other constructs of candidate phy4-interacting partners were obtained by Gateway cloning. Full length entry clones were generated by Phusion amplification of cDNA in a 2-step PCR whereby attB1 and attB2 sites were attached to the product. For N-terminal and C-terminal tags, the CDS was amplified with and without stop codons, respectively. Gel-purified PCR products were cloned into pDONR207 with BP-clonase (Invitrogen). PRL1 and PLP entry clones were created by TA-cloning of PCR products into pCR8GW TOPO (Invitrogen). The resulting full length entry clones were used to create expression constructs for Y2H assays and cytological analyses via the LR reaction. Inserts were recombined into AD: and :AD prey vectors pGADT7g (Uetz and Grigoriev, 2005) and pGADCg (Stellberger et al., 2010), respectively, and transformed into yeast together with BD:phy4 and phy4:BD holophytochrome baits for semi-quantitative growth assays in order to verify the interaction and its potential light-dependency. For this, 2×10^5 doubly transformed yeast cells were spotted on TSD medium with 3-AT and 30 μM PCB and incubated for 5 days in continuous 0.7 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Rc) or 12 min/h red light (660 nm LEDs, 4 $\mu\text{mol m}^{-2} \text{s}^{-1}$) pulses (Rp) or red followed by 12 min/h far-red (740 nm LEDs, 5 $\mu\text{mol m}^{-2} \text{s}^{-1}$) pulses (Rp+FRp) or darkness (D). Various controls were included to ensure that appropriate interactions were being monitored (see also Jaedicke et al., 2012). Constitutive dimerization was demonstrated by BD:phy4 and phy4:BD baits combined with the AD:phy4 prey. As negative controls and for the adjustment of appropriate 3-AT concentrations, BD:phy4 and phy4:BD baits were combined with the empty pGADT7 vector (from this 2.5 mM or 1 mM 3-AT were used for doubly transformed yeast growth assays). As a control for successful holophytochrome formation, BD:phyA (in pGBKT7) and phyA:BD (in pBAC) baits were combined with AD:FHY1 (1 mM 3-AT) as this Y2H interaction is Pfr-dependent (Hiltbrunner et al., 2005).

For intracellular localization studies, each partner CDS was Gateway-cloned into modified destination vectors p2CGW7-2 \times 35S, p2GWC7-2 \times 35S, p2FGW7-2 \times 35S, and p2GWF7-2 \times 35S (Karimi et al., 2002), creating N- or C-terminal CFP- or GFP-fusions, respectively. For *in planta* interaction studies, the CDS was recombined into pSAT4-DEST-n(1-174)eYFP-CI(YFP_N:) and pSAT4(A)-DEST-n(1-174)eYFP-N1(:YFP_N) or in pSAT5-DEST-C(175-end)eYFP-CI (YFP_C:) and pSAT5(A)-DEST-c(175-end)EYFP-N1 (:YFP_C) destination vectors (Citovsky et al., 2006). phy4 pSAT constructs were created as described (Jaedicke et al., 2012). Negative controls for interaction were provided by empty YFP_C and YFP_N vectors in combination with the HIP pSAT expression plasmids.

In Planta Localization and Interaction Studies

Biolistic transfection of *Physcomitrella* filaments was performed as described (Jaedicke et al., 2012). For subcellular protein localization studies, transfected filaments were incubated for 2 days in D and observed via fluorescence microscopy without pre-treatment and following 1 h R pre-treatment (660 nm LEDs, 3 $\mu\text{mol m}^{-2} \text{s}^{-1}$). For split-YFP studies, plant material was incubated for 3 days in D and then subjected to confocal microscopy with and without analogous pre-treatments.

Transformed cells were identified by R fluorescence of the co-transfected nuclear marker mCherry:VirD2NLS using a Leica Z16 apo fluorescence microscope. For localization and split-YFP studies, an automated DM6000b fluorescence microscope and a TCS SP2 AOBs confocal laser scanning microscope (both Leica) were used, respectively. For filter cube and look-up table details, see Jaedicke et al. (2012). Single scans of YFP fluorescence were recorded with 4x line and 6x frame average. In order to distinguish YFP signals from possible mCherry bleed-through, YFP emission detection was narrowed to 525–535 nm. The red mCherry- and chlorophyll autofluorescence and the transmission images were taken with 4x line and 2x frame average. Adobe Photoshop CS5 was used for image processing (enhancement of brightness and contrast).

In Silico Analyses

The protein sequences derived from the relevant HIP gene models were further analyzed to identify domains and thus potential functions. Co- and possible synexpression with phy4 was investigated using data from eFP, Phytozome and Genevestigator with the help of ClustVis^{3,4,5,6}.

RESULTS AND DISCUSSION

Y2H Library Screens Revealed 14 Putative HIP's

We developed Y2H methods to screen for phy4-interacting proteins that might be involved in the phy4 cytoplasmic signaling pathway. Importantly, instead of photochemically inactivated fragments, we established the use of functional, full-length holophytochrome as bait by feeding the transgenic yeast cells with PCB to allow auto assembly *in vivo*, as evidenced by R/FR reversible interaction of *Arabidopsis* phyA with FHY1 (Supplementary Figure S1) and, subsequently, of phy4 with various putative partners. We generated bait constructs with the BD N- and C-terminally fused to full-length phy4, carefully characterizing their homodimerization and autoactivation properties. The N-terminal fusion thereby showed a higher autoactivation of the *HIS3*-reporter than the C-terminally fused bait. Accordingly, 3-AT (a competitive inhibitor of the *HIS3*

³bar.utoronto.ca/efp_physcomitrella/cgi-bin/efpWeb.cgi

⁴phytozome.jgi.doe.gov

⁵www.genevestigator.com

⁶biit.cs.ut.ee/clustvis/

gene product) was included at 2.5 and 1 mM, respectively. Both phy4 baits showed constitutive homodimerization together with the phy4 prey fused N-terminally to the AD (Supplementary Figure S1). We screened the pre-transformed cDNA library with both apo-BD:phy4 and holo-phy4:BD in Rc via sequential double transformation and mating, respectively. Functionality of the mating protocol for holophytochrome formation was confirmed by mating Y187 phyA:BD to AH109 AD:FHY1 and selecting on PCB supplemented medium in Rc (Supplementary Figure S1). The yeast mating protocol evoked substantially less autoactivation than the transformation protocol, necessitating minimal 3-AT (0.25 mM only for phy4:BD). Therefore we expected also weak or temporary interactions with phy4 Pfr to be identifiable.

The 4.2×10^5 clones were screened with phy4:BD bait, corresponding to about fourfold coverage of the *Physcomitrella* transcriptome. 1×10^4 clones were additionally screened with BD:phy4 bait. The 69 and 75 clones, respectively, survived two rounds of selection and were sequenced, following which 54 and 23 yeast clones excluding duplicates were subjected to BLAST analyses at Cosmoss to identify the corresponding loci in the *Physcomitrella* genome (see Supplementary Tables S1 and S2). Annotation was improved by domain analysis and by BLAST searches of the respective protein sequences against the *Viridiplantae* and *Arabidopsis* non-redundant protein sequence databases. On this basis, genes which were considered unlikely to be involved in signaling were not considered further, leaving candidates putatively involved in signaling processes, posttranslational modification, transcription/translation and likely membrane, transport- or cytoskeleton-associated proteins. We subsequently analyzed 11 and 4 candidates, respectively, by cloning the full-length cDNA into AD: and :AD prey plasmids and combining them in yeast with full-length photoactive BD:phy4 and phy4:BD bait in semiquantitative growth assays under different light conditions (Supplementary Figure S2). Only one now failed to interact with BD:phy4 and phy4:BD (data not shown), perhaps because a serendipitous binding site shown by the partial cDNA expression product was lost in the full length configuration. The remaining 14 candidates showed interaction with BD:phy4 and/or phy4:BD full-length holophytochrome bait either constitutively or, more commonly, in a Rp-enhanced (FRp-reversible) manner. These were thus provisionally termed holophytochrome-interacting proteins (HIP1–14).

Verification and Characterization of HIP's

The cellular localization of each putative HIP was subsequently studied *in planta* using *Physcomitrella* filaments transfected with appropriate fluorescent protein hybrid constructs in D and after R pre-treatment. As a rule the same cells were observed before and after illumination. Finally, interaction *in planta* between each putative HIP and phy4 was studied using bimolecular fluorescence complementation (in D and after R pre-treatment) with split-YFP hybrid proteins, whereby the cellular localization of the interaction itself was documented. In all cases, hybrids in which the additional domains were fused either to the N- or the C-termini of the putative HIP and phy4 were generated,

thereby accounting both for possible functional artifacts arising from the fusion and for topological differences that might affect the interaction readout. Respective split-YFP negative controls are shown in Supplementary Figure S3. Twelve out of the 14 putative HIP's were thereby proven to interact *in planta*. The two putative HIP's which failed to show an *in planta* split-YFP signal might interact too weakly for YFP reconstitution or represent false positives from Y2H. The first of these, Pp3c19_20830V1.1 (initially named HIP2) represents a likely pirin-related protein (see HIP1 below). Also depending on the configuration, this protein localized to the cytoplasm or to cytoplasm and nucleus, respectively (see Supplementary Figure S5). The second, Pp3c1_11190C1.1 (Cosmoss Version 3 accession number; initially named HIP10), encodes a protein with an N-terminal ankyrin- and a C-terminal BTB/POZ (Boad-complex, Tamtrac, Bric a brac/POx virus and ZF)-domain (see Supplementary Figure S4). CFP-fusions were localized either in cytosolic speckles or uniformly in the cytoplasm and nucleus, depending on the FP-configuration. Our results for the 12 HIP's confirmed *in planta* are described below.

HIP1 (Pp3c2_10320V1.1)

Independent library screens revealed two different members of the **pirin** iron-containing subgroup of the **cupin superfamily**, (HIP1, 38.7 kDa) and Pp3c19_20830V1.1 (initially HIP2, 33.9 kDa). Semiquantitative studies of full length CDS models showed strong and weaker binding to phy4 holophytochrome in Y2H, respectively. Indeed except in the case of HIP1–BD:phy4, the interaction was R/FR reversible (Supplementary Figure S2). However, as Pp3c19_20830 showed insignificant split-YFP-signals (Supplementary Figure S5B) it was excluded as a genuine HIP. In *Physcomitrella* HIP1 was distributed in the cytoplasm and nucleus (**Figure 1A**). *In planta* interaction was verified for HIP1 in the phy4:YFP_C–YFP_N:HIP1 configuration (**Figure 1B**) although the result was ambiguous because the signal was detectable either in cytoplasm and nucleus or in cytosolic speckles and nucleus in different cells both in D and following R pre-treatment.

HIP1 is apparently expressed under all conditions and in all cell types although weekly in leaflets (see Supplementary Materials). The data provide little evidence for synexpression with phy4.

Holophytochrome-interacting protein 1 (like Pp3c19_20830) harbors an N-terminal cupin domain including the potential metal-coordinating residues H116, H118, H160, and E162 corresponding to H57, H59, H101, and E103 in the human sequence (Pang et al., 2004; Adams and Jia, 2005). The pirin-typical C-terminal domain (which resembles the cupin domain while lacking the metal-coordinating residues) is also present in HIP1 (but not Pp3c19_20830). Conserved in mammals, fungi, plants and prokaryotes, four and three genes encoding pirin-like proteins exist in *Arabidopsis* and *Physcomitrella*, respectively (see Summary Table 1 in the Supplementary Material).

Pirin functions are quite diverse. Human, *Arabidopsis* and *E. coli* pirins have been shown to possess quercetinase activity (Adams and Jia, 2005). Quercetin – amongst other things – plays a role in UV acclimation (Hectors et al., 2014) and has




FIGURE 1 | HIP1 (Pp3c2_10320V1.1) intracellular localization (A) and split-YFP-studies of HIP1–phy4 interaction (B) each without (D) and with red pre-treatment (R) using fluorescence- and confocal microscopy, respectively. (A) Green fluorescence of N- and C-terminal HIP1:GFP fusions monitored by GFP (column 1) and was visible in nucleus and cytoplasm in both cases; Red fluorescence of the co-transfected mCherry:NLS nuclear marker (RFP) is shown in column 2; Chlorophyll autofluorescence (chlorophyll) and a differential interference contrast image (DIC) is shown in columns 3 and 4, respectively. For each FP configuration cells were photographed before (D) and after red light pre-treatment (R). The apparent “hole” in the nucleus of the HIP1:GFP fluorescence images and the corresponding black dot in the DIC images represents a large gold particle. **(B)** All possible split-YFP configurations were analyzed without (D) and with red pre-treatments (R) via confocal microscopy as shown in column 1: YFP_C:phy4–YFP_N:HIP1 (rows 1+2), YFP_C:phy4–HIP1:YFP_N (rows 3 and 4), phy4:YFP_C–YFP_N:HIP1 (rows 5–8) and phy4:YFP_C–HIP1:YFP_N (rows 9 and 10). Analogous to panel (A), columns depict (from left to right) YFP fluorescence, RFP fluorescence from mCherry:NLS, chlorophyll autofluorescence and a transmission image. Two different signal patterns were observed for phy4:YFP_C–YFP_N:HIP1, thus representative images are shown for each. Other configurations yielded only background signals alongside channel bleed-through. Scale bars 30 μm.

been shown to inhibit polar auxin transport (Fischer et al., 1997; Kuhn et al., 2011; Buer et al., 2013) and thus might play a role in tropic responses (Lewis et al., 2011). Indeed, light-grown *prn1⁻* *Arabidopsis* seedlings showed disoriented hypocotyl growth, whereas etiolated mutant seedlings showed enhanced UV-induced quercetin levels (Orozco-Nunnally et al., 2014). In *Arabidopsis* quercetin is localized in and around the nucleus, in the endomembrane system and at the plasma membrane (Peer et al., 2001). Although this would correlate with the phy4–phototropin complex at the plasma membrane (Jaedicke et al., 2012) our data do not imply that HIP1 is localized at the plasma membrane (**Figure 1**). Diverse physiological effects have been associated with pirins in plants, including pathogen susceptibility, apoptosis, ABA-suppression of seed germination and B-induced *LHC* expression, possibly in association with G-proteins (Orzaez et al., 2001; Lapik and Kaufman, 2003; Warpeha et al., 2007; Zhang et al., 2014). Phytochrome signaling via G-proteins is controversial, however (Neuhaus et al., 1993; Okamoto et al., 2001; Jones et al., 2003; Urano et al., 2013). The *Physcomitrella* genome codes for at least one G-protein but this has not been studied to-date.

HIP3 (Pp3c10_4820V1.1)

Holophytochrome-interacting protein 3 (37.4 kDa) carries CHY, CTCHY, and C3HC4 (referring to a specialized cysteine-histidine pattern) RING-type ZF domains. Y2H analysis with the full-length CDS showed weak but R-enhanced interaction with phy4 in all conformations. phy4:BD–ADg:HIP3 showed interaction in Rc but not in Rp (Supplementary Figure S2). *In planta*, although no particular changes in localization were seen upon irradiation, HIP3 showed cytosolic or nuclear and cytosolic localization in approximately half of the observed cells, respectively (CFP:HIP3) or cytosolic and nuclear (HIP3:CFP) localization, depending on the position of the CFP tag (see **Figure 2A**). Cytosolic localization is interesting because both fusion proteins might be small enough (37 kDa + 27 kDa tag) to enter the nucleus passively, at least as monomers. Conventional NLS and NES motifs are not apparent in the HIP3 sequence. *In planta* interaction was seen for phy4:YFP_N–HIP3:YFP_C in the nucleus and the cytoplasm, consistent with the localization data, the other combinations yielding only background signals alongside channel bleed-through (see **Figure 2B**). Interestingly, the split-YFP signal only appeared after R pre-irradiation, implying Pfr-dependent interaction *in planta* as in Y2H.

Holophytochrome-interacting protein 3 is apparently expressed under all conditions and in all cell types except in dehydrated material (see Supplementary Materials). The data provide little evidence for synexpression with phy4.

Zinc finger-proteins are known to bind diverse ligands including DNA, RNA, lipids, and proteins (Laity et al., 2001; Matthews and Sunde, 2002), thereby executing functions in a variety of cellular processes such as transcription, translation, cytoskeleton organization, protein folding, and others. The CHY-ZF is named due to its conserved motif CxHY, the function of which is still unknown (Leng et al., 2003). The 40–60 residues comprising RING-type ZF's, however, are considered to bind two zinc atoms and to mediate protein–protein interactions

(Freemont, 1993; Borden and Freemont, 1996; Saurin et al., 1996). RING-type ZFs can also serve as E3-ubiquitin ligases in protein degradation and are thus commonly involved in gene regulation via repressor destruction in plants. BLAST searches indeed revealed high similarity between HIP3 and 4–6 other *Physcomitrella* and at least seven *Arabidopsis* proteins. For example, *Arabidopsis* RING FINGER AND CHY ZINC FINGER DOMAIN-CONTAINING PROTEIN 1 (AT5G22920) shows 63% identity in the 250 residue region of similarity. This nuclear-localized E3 ubiquitin ligase is involved in the regulation of stomatal opening, its activity being regulated via phosphorylation by SnRK2.6 (Ding et al., 2015). CHY and CTCHY and RING-type ZF protein (AT5G25560), a PGPD 14-like protein (Kosarev et al., 2002) of unknown function, shows 57% identity except at the N-terminus. Similarly, MYB30-INTERACTING E3 LIGASE 1 (AT5G18650) with 59% identity also except in the 64 residue N-terminal region) degrades the MYB30 transcription factor, a negative regulator of defense responses in *Arabidopsis* (Marino et al., 2013). Several known *Arabidopsis* phytochrome-interacting and/or signaling molecules are known to possess RING-ZF domains, for example RFI2, which plays a role in phytochrome-controlled seedling de-etiolation (Chen and Ni, 2006) or VOZ1, a phyB partner controlling flowering (Yasui et al., 2012). The central regulator of photomorphogenesis, the E3 ubiquitin ligase COP1 also contains a C3HC4-type RING-ZF motif. Moreover, the PENTA protein involved in translational regulation of PORA (the key light-regulated enzyme in chlorophyll biosynthesis) by interaction with phyB is also a C3H-type ZF protein (Paik et al., 2012).

HIP4/PRL1 (Pp3c16_8560E1.1)

Holophytochrome-interacting protein 4 (53.9 kDa) is encoded by *PRL1* sharing 62% sequence identity with its homolog in *Arabidopsis*. In Y2H studies full length HIP4/PRL1 interacted with holophy4 in all four bait–prey configurations. Three interactions were R-enhanced and partially FR-reversible whereas BD:phy4–HIP4:ADg interacted constitutively (Supplementary Figure S2). Independently of the light treatment, HIP4:GFP localized to both nucleus and cytoplasm while CFP:HIP4 was concentrated in the nucleus (**Figure 3A**). Constitutive interaction was confirmed *in planta* in all combinations except phy4:YFP_C–HIP4:YFP_N (**Figure 3B**), the reconstituted split-YFP signal being apparent in both nucleus and cytoplasm for phy4:YFP_C–YFP_N:HIP4 and, less clearly, YFP_C:phy4–HIP4:YFP_N, corresponding to the localization of HIP4:GFP. YFP_C:phy4–YFP_N:HIP4 interacted in the cytoplasm.

Holophytochrome-interacting protein 4 is apparently expressed under all conditions and in all cell types (see Supplementary Materials). The Genevestigator data for both treatments and anatomy provide some evidence for synexpression with phy4. This is not apparent for the other data, however.

PLEIOTROPIC REGULATORY LOCUS 1 is well-conserved amongst land plants, fungi, and animals including mammals. Two homologs are present in both *Physcomitrella* and *Arabidopsis* (HIP4/PRL1 & Pp3c5_15520V1.1 and PRL1 & PRL2, respectively). *Arabidopsis* PRL1 is an α -importin-interacting




FIGURE 2 | HIP3 (Pp3c10_4820V1.1) intracellular localization (A) and split-YFP-studies of HIP3–phy4 interaction (B) each without (D) and with red pre-treatment (R) using fluorescence- and confocal microscopy, respectively. Overall figure structure is analogous to **Figure 1**. **(A)** CFP:HIP3 (rows 1–4) and HIP3:CFP (rows 5 and 6). CFP:HIP3 localization was ambiguous, being observable either in the nucleus and cytoplasm (rows 2 and 4) or in the cytoplasm only (rows 1 and 3) in both conditions, whereas HIP3:CFP showed nuclear and cytoplasmic localization (column 1, CFP). **(B)** Split-YFP analyses of YFP_N:phy4–YFP_C:HIP3 (rows 1 and 2), YFP_N:phy4–HIP3:YFP_C (rows 3 and 4) and phy4:YFP_N–HIP3:YFP_C (rows 5 and 6) configurations. Significant nucleo-cytoplasmic signals were only seen for phy4:YFP_N–HIP3:YFP_C after R pre-treatment (column 1, YFP; D and R here from separate experiments). All other configurations/conditions yielded only background signals alongside channel bleed-through. Scale bars 30 μm.




FIGURE 3 | HIP4 (Pp3c16_8560E1.1) intracellular localization (A) and split-YFP-studies of HIP4–phy4 interaction (B) each without (D) and with red pre-treatment (R) using fluorescence- and confocal microscopy, respectively. Overall figure structure is analogous to **Figure 1**. **(A)** CFP::HIP4 (rows 1 and 2) was mostly nuclear and HIP4::GFP (rows 3 and 4) was distributed in nucleus and cytoplasm (column 1, FP). **(B)** All possible split-YFP configurations were analyzed in D and R, respectively: YFP_C:phy4–YFP_N:HIP4 (rows 1 and 2), YFP_C:phy4–HIP4:YFP_N (rows 3 and 4), phy4:YFP_C–YFP_N:HIP4 (rows 5 and 6) and phy4:YFP_C–HIP4:YFP_N (rows 7 and 8). Significant signals were observed for YFP_C:phy4–YFP_N:HIP4 (cytoplasmic signal), phy4:YFP_C–YFP_N:HIP4 (nucleo-cytoplasmic signal), and weakly for YFP_C:phy4–HIP4:YFP_N (maybe nucleo-cytoplasmic signal) in both D and R (column 1, YFP). phy4:YFP_C–HIP4:YFP_N yielded only background signals alongside channel bleed-through. Scale bars 30 μm.

protein localizing both to the nucleus and associating with the nuclear envelope- and ER membranes (Nemeth et al., 1998). It harbors seven WD40 domains and a variant of the SV40-type NLS in the C-terminal region (residues 238–241), which is not constitutively recognized but rather dependent on phosphorylation of the NLS or the importin (Nemeth et al., 1998). Our localization studies of HIP4/PRL1 indicate that at least R does not affect recognition of the NLS and subsequent nuclear import since we do not observe any change in localization upon light treatment (Figure 3A). Although N-terminally fused CFP:HIP4 was exclusively nuclear, HIP4:GFP was seen in both the nucleus and cytoplasm, whereas YFP_C:phy4–YFP_N:HIP4 was exclusively cytoplasmic (Figure 3B). Perhaps phy4 binds the C-terminus of HIP4/PRL1, thereby hampering NLS recognition and effecting cytoplasmic localization.

The PRL1 WD40 domains probably constitute β -propellers, together forming a platform for protein–protein interactions. Such platforms are often involved in signal transduction and transcriptional regulation, as in the case of COP1 in repression of photomorphogenic development. *Arabidopsis* PRL1 seems to be involved in numerous responses, as the name implies. Initially reported to play a role in cell polarity determination by acting on the assembly of the actin cytoskeleton in fission yeast (Xia et al., 1996), PRL1 was later suggested to play a central role in integrating light signals with cytokinin- and sugar response pathways in plants (Nemeth et al., 1998). Indeed *prl1*[−] hypocotyls are shorter than the wild type in darkness but not in light, representing a weak COP phenotype, consistent with PRL1 acting as a positive regulator of brassinosteroid synthesis (Szekeres et al., 1996; Nemeth et al., 1998). It was shown to promote BR-biosynthesis by interacting with different SnRK's (see also HIP3 above) in light and dark, possibly by targeting them for degradation (Bhalerao et al., 1999; Lee et al., 2008; Flores-Perez et al., 2010). PRL1 is also involved in root meristem regulation possibly by restricting transcription of the homeobox transcription factor WOX5 to the quiescent center (Ji et al., 2015). Thus PRL1 in *Arabidopsis* might either represent a convergence center for integrating light, sugar, hormone, and other responses or be involved in distinct signaling pathways in different cells. SnRK photoregulation would be a possible function of HIP4/PRL1 in *Physcomitrella*, as SNF1 homologs are present (Thelander et al., 2004, 2007). Light, sugar and the phytohormones cytokinin and auxin are all closely involved in the developmental switches in *Physcomitrella* whereby filament cells develop into chloronemata, caulonemata, or gametophore buds (Thelander et al., 2004, 2007; Decker et al., 2006; von Schwartzenberg, 2006).

HIP5 (Pp3c7_3040V1.1)

Holophytochrome-interacting protein 5 (54.4 kDa) bears four **kelch repeats** toward the C-terminus. It has one homolog in *Physcomitrella* (Pp3c11_19970V1.1, see Summary Table 1 in the Supplementary Material). The semi-quantitative Y2H interaction assay confirmed R-enhanced and FR-reversible interaction in all phy4–HIP5 configurations tested with the exception of BD:phy4–HIP5:ADg which showed no interaction (Supplementary Figure S2). HIP5 seems to be constitutively present in nucleus and

cytoplasm irrespective of the position of the fluorescent tag (Figure 4A). Interestingly, an older and N-terminally shorter version of the HIP5 gene model showed exclusively cytoplasmic localization as N-terminal GFP-fusion, implying that the putative NLS resides at the very N-terminus (in the first 129 amino acids; not shown). phy4–HIP5 interaction was confirmed *in planta*, but in contrast to the Y2H data, interaction was not affected by light treatments (Figure 4B). Analogously to the HIP5 localization data, nuclear and cytoplasmic distribution was seen for phy4:YFP_C–HIP5:YFP_N and perhaps also for YFP_C:phy4–YFP_N:HIP5 (although this fluorescence might represent RFP bleed-through in the YFP channel as the signal was very weak). Interestingly, the YFP-signal from the reconstituted phy4:YFP_C–YFP_N:HIP5 located exclusively to the cytoplasm, although GFP:HIP5 was apparent in both cytoplasm and nucleus.

Holophytochrome-interacting protein 5 is apparently expressed under all conditions and in all cell types (see Supplementary Materials). The Genevestigator data for treatments, anatomy and development provide some evidence for synexpression with phy4. This is not apparent for the eFP and Phytozome data, however.

The ~50-residue kelch motif forms a blade-like fold comprising 4–6 anti-parallel β -sheets, often repeated to form a β -propeller structure frequently involved in protein–protein interactions. In fungi, various kelch repeat proteins associate with microtubules and actin nucleating proteins at cell termini, thereby establishing cell polarity (Mata and Nurse, 1997; Feierbach et al., 2004) whereas others are involved in polarized growth (Takeshita et al., 2008). We speculate therefore that HIP5 might play a similar role in phy4-directed vectorial growth. Indeed, amongst both fungal and *Arabidopsis* BLAST hits was a scruin – an actin crosslinker, implying a role in cytoskeleton organization consistent with the role of microfilaments in filament tip cell phototropism (Meske and Hartmann, 1995; Meske et al., 1996). On the other hand, HIP5 shares ~35% sequence identity with many *Arabidopsis* F-box/kelch-repeat (FBK) proteins such as galactose oxidase/kelch repeat superfamily protein (AT1G14330), MIR2111-5p target protein (AT3G27150) and SKIP11 (AT2G02870), although these sequences rather align with the C-terminal region and thus the kelch repeats of HIP5 (64–71% of the query), whereas the first 135–175 amino acids of HIP5 are dissimilar to the *Arabidopsis* proteins. All three *Arabidopsis* FBK proteins are chloroplast- and/or nuclear-localized, interacting with SKP1-like proteins (Schumann et al., 2011). FBK kelch repeats for example are responsible for binding of target proteins to the SCF ubiquitin ligase complex in the degradation pathway (Cardozo and Pagano, 2004; Hua and Vierstra, 2011). The *Arabidopsis* and *Physcomitrella* genomes contain ca. 100 and 71 FBK genes, respectively (Schumann et al., 2011), several of which are known to be involved in photoresponses. The PAS/LOV-domain-containing ZEITLUPE, FKF1, LKP2 FBK's are themselves photoreceptors (Sawa et al., 2007; Demarsy and Fankhauser, 2009; Takase et al., 2011), whereas AFR and CFK1 are positive regulators downstream of phyA and phyB, respectively (Wei et al., 1994a,b; Harmon and Kay, 2003; Franciosini et al., 2013). CFK1 seems to be absent from *Physcomitrella* although the




FIGURE 4 | HIP5 (Pp3c7_3040V1.1) intracellular localization (A) and split-YFP-studies of HIP5–phy4 interaction (B) each without (D) and with red pre-treatment (R) using fluorescence- and confocal microscopy, respectively. Overall figure structure is analogous to **Figure 1**. **(A)** GFP::HIP5 (rows 1 and 2) and HIP5::GFP (rows 3 and 4) were distributed in the cytoplasm and in the nucleus (column 1, GFP). **(B)** Three split-YFP configurations were examined and showed different signal patterns/strengths (column 1, YFP): YFP_C:phy4–YFP_N:HIP5 (rows 1 and 2; maybe nucleo-cytoplasmic distribution), phy4:YFP_C–YFP_N:HIP5 (rows 3 and 4; cytoplasmic distribution) and phy4:YFP_C–HIP5:YFP_N (rows 5 and 6; nucleo-cytoplasmic distribution). The apparent YFP_C:phy4–YFP_N:HIP5 signal might also represent RFP bleed-through in the YFP channel since the signal was very weak. Scale bars 30 μm.

COP9 signalosome with which CFK1 interacts is thought to be universal in eukaryotes (Wei et al., 1998), thus HIP5 might be involved in a similar regulatory system. Most FBK's are nuclear although some are located in both cytoplasm and nucleus – as is the 26S proteasome (Schumann et al., 2011). Pp3c2_13250, Pp3c1_22300, Pp3c10_4990, and Pp3c14_4360, on the other hand, might represent *Arabidopsis* AFR homologs with 29–26% identity covering 72–100% of the query sequence.

HIP6 (Pp3c6_4480V1.1)

Holophytochrome-interacting protein 6 (43.7 kDa) harbors a predicted transmembrane domain and a **p-loop containing sulfotransferase** domain at the N- and C-terminus, respectively. Interaction of HIP6 with phy4 in Y2H was constitutive in the BD:phy4-AD:HIP6 and phy4:BD-AD:HIP6 configurations, whereas phy4:BD-HIP6:ADg showed R/FR-reversible interaction (Supplementary Figure S2). *In planta* both N- and C-terminal fusion configurations localized exclusively to the cytoplasm, with HIP6:GFP concentrating in discrete regions (cytoplasmic foci; **Figure 5A**). This distribution was also observed for phy4:YFP_C-YFP_N:HIP6 interaction, whereas other split-YFP configurations (phy4:YFP_C-HIP6:YFP_N and perhaps YFP_C:phy4-YFP_N:HIP6) rather showed cytoplasmic and weaker nuclear localization (**Figure 5B**; although narrower emission detection implied in the latter case that the apparent YFP nuclear signal might have arisen from RFP bleed-through; data not shown). Light pre-treatments had little effect on *in planta* interaction (also in the case of phy4:YFP_C-HIP6:YFP_N corresponding to phy4:BD-HIP6:ADg in Y2H). In neither Y2H nor split-YFP *in planta* was significant interaction seen between N-terminally tagged phy4 and C-terminally tagged HIP6.

Holophytochrome-interacting protein 6 is apparently expressed under all conditions and in all cell types (see Supplementary Materials). The Genevestigator and Phytozome data provide some evidence for synexpression with phy4.

The p-loop domain is widespread in nature, using conserved motifs GXXXXGK(T/S) and ZZZ(D/E) (where X and Z represent arbitrary and hydrophobic residues, respectively) with a β -sheet followed by a glycine-rich loop and an α -helix to co-ordinate the ATP or GTP β - γ -phosphate moiety (Walker et al., 1982; Saraste et al., 1990; Milner-White et al., 1991; Ramakrishnan et al., 2002). Thus, HIP6 belongs to the superfamily of p-loop-containing nucleoside triphosphate hydrolases. This large group is involved in processes like replication, transcription, DNA repair, translation, signal transduction, protein localization, and -transport, signal-sequence recognition, membrane transport, chromosome partitioning, and activation of metabolites (Saraste et al., 1990; Vetter and Wittinghofer, 1999; Koonin et al., 2000). Apart from the common motifs and the p-loop structure itself, the sequences (even the core of the p-loop) and structures are extremely diverse, hampering (phylogenetic and) functional classification (Pathak et al., 2014). So far, only a few plant p-loop NTPases have been functionally characterized. Some are thought to be involved in pathogen defense responses and others in salinity stress tolerance (Fenyk et al., 2012; Cheung et al., 2013).

Plant sulfotransferases (SOT's) are either soluble or transmembrane proteins that transfer the sulfate group from

a 3'-phosphoadenosine-5'-phosphosulfate donor to proteins or smaller substrates including brassinosteroids, jasmonates, flavonoids, or salicylic acid (Klein and Papenbrock, 2004; Hirschmann et al., 2014). Intriguingly, plant SOT's sulfate the flavonol quercetin (Varin et al., 1992), discussed above in connection with pirins. Quercetin and quercetin sulfates are thought to regulate root development by altering basipetal auxin transport, elongation growth and gravitropism (Faulkner and Rubery, 1992; Lewis et al., 2011). Whereas three HIP6 homologs are apparent in *Physcomitrella* (see Summary Table 1 in the Supplementary Material), 22 SOT's with different sulfotransfer-domain-types and diverse expression patterns have been identified in *Arabidopsis*. The hydroxyl-jasmonate sulfating SOT15 is transcriptionally controlled by the circadian clock and PIF4 and -5 (Yamashino et al., 2013) whereas SOT18 seems to be regulated by the phytochrome-controlled transcription factor HY5 (Huseby et al., 2013). Various *Arabidopsis* SOT's are thought to be functional in such diverse processes as defense and stress responses, regulation of hormone synthesis or activity, signaling and developmental regulation (Rouleau et al., 1999; Gidda et al., 2003; Marsolais et al., 2007; Hirschmann et al., 2014). Mutation of *Arabidopsis* tyrosyl-protein sulfotransferase has pleiotropic defects such as in cotyledon and leaf size and pigmentation, flower number and root development (Komori et al., 2009) the latter being associated with disturbed auxin distribution due to altered PIN and auxin-biosynthesis gene expression (Zhou et al., 2010). The function of HIP6 is therefore unclear, especially as it shares highest similarity to the newly identified *Arabidopsis* SOT's 19–21 (see Supplementary Materials) whose substrate specificity is unknown (Klein and Papenbrock, 2009; Hirschmann et al., 2014).

HIP7 (Pp3c17_9390V1.1)

The singleton HIP7 (31.5 kDa) is likely to represent a **Ser/Thr protein kinase**. In Y2H studies full length HIP7 interaction with phy4 was comparatively weak but nevertheless R-enhanced (Supplementary Figure S2), strongest growth being observed for the phy4:BD-ADg:HIP7 configuration. HIP7 fusions with CFP were intracellularly distributed in the nucleus and the cytoplasm (**Figure 6A**) with CFP:HIP7 in particular showing cytosolic foci in ~40% of the cells observed (data not shown). In split-YFP, nuclear and cytoplasmic signals were seen for YFP_C:HIP7 in combination with both YFP_N:phy4 and phy4:YFP_N, in accordance with the localization data (**Figure 6B**). Since the split-YFP signals were significantly weaker than those from the RFP:NLS nuclear marker, we checked for RFP bleed-through by narrowing detection to 10 nm around the emission maximum of YFP. As the fluorescence pattern was unchanged, we consider the signal to represent a *bona fide* nuclear interaction. Interestingly, the phy4:YFP_N-YFP_C:HIP7-interaction appeared to be R-dependent, the signal only being observable after illumination, implying Pfr-dependent binding *in planta*.

Holophytochrome-interacting protein 7 is apparently expressed under all conditions and in all cell types (see Supplementary Materials). Some Genevestigator data groups imply synexpression with phy4 as does the Phytozome data.




FIGURE 5 | HIP6 (Pp3c6_4480V1.1) intracellular localization (A) and split-YFP-studies of HIP6–phy4 interaction (B) each without (D) and with red pre-treatment (R) using fluorescence- and confocal microscopy, respectively. Overall figure structure is analogous to **Figure 1**. **(A)** Intracellular localizations of CFP::HIP6 (rows 1 and 2) and HIP6::GFP (rows 3 and 4) were both localized in the cytoplasm, with HIP6::GFP showing subcellular foci (column 1, FP). **(B)** All four split-YFP configurations were investigated: YFP_C::phy4–YFP_N::HIP6 (rows 1 and 2), YFP_C::phy4–HIP6::YFP_N (rows 3 and 4), phy4::YFP_C–YFP_N::HIP6 (rows 5 and 6), and phy4::YFP_C–HIP6::YFP_N (rows 7 and 8). Significant YFP signals were apparent only for phy4::YFP_C–YFP_N::HIP6 (cytoplasm and cytoplasmic foci) and phy4::YFP_C–HIP6::YFP_N (cytoplasm and nucleus; column 1, YFP), whereas apparent YFP_C::phy4–YFP_N::HIP6 nuclear fluorescence might represent RFP bleed-through and YFP_C::phy4–HIP6::YFP_N only shows chloroplast bleed-through. Scale bars 30 μ m.




FIGURE 6 | HIP7 (Pp3c17_9390V1.1) intracellular localization (A) and split-YFP-studies of HIP7-phy4 interaction (B) each without (D) and with R pre-treatment (R) using fluorescence- and confocal microscopy, respectively. Overall figure structure is analogous to **Figure 1**. **(A)** Intracellular localization of CFP:HIP7- (rows 1 and 2) and HIP7:CFP-fusions (rows 3 and 4) was observed, showing nuclear and cytoplasmic signal patterns (column 1, CFP). CFP:HIP7 in addition showed cytosolic foci in ~40% of the cells observed. **(B)** Three of four possible split-YFP configurations were analyzed: YFP_N:phy4-YFP_C:HIP7 (rows 1 and 2), YFP_N:phy4-HIP7:YFP_C (rows 3 and 4) and phy4:YFP_N-YFP_C:HIP7 (rows 5 and 6). YFP_N:phy4-YFP_C:HIP7, and phy4:YFP_N-YFP_C:HIP7 yielded significant split-YFP-signals mostly in nucleus and cytoplasm (column 1, YFP). In the latter case significant signals were only seen after R-pre-treatment (D and R conditions here being shown from independent experiments). Scale bars 30 μm.

HIP7 shows low but significant similarity to various *Arabidopsis* Ser/Thr-protein kinases such as HT1 (AT1G62400), a kinase with an armadillo repeat domain (AT5G18700), a MAP kinase (AT1G73660), and a PAS-domain-containing protein tyrosine kinase (AT3G06620) with ~40% query coverage and ~30% sequence identity). HIP7-based BLAST searches of *Viridiplantae* proteins yielded many with somewhat more concordant similarity, most of which have been judged to be HT1 relatives. *Arabidopsis* HT1 is a protein kinase involved in the regulation of stomatal aperture (Hashimoto et al., 2006). Although stomatal opening is generally seen as a phototropin-mediated B effect, R is effective in maintaining stomatal opening, a response specifically impaired in *ht1*⁻ (Matrosova et al., 2015). Although the effect can largely be explained by photosynthetic CO₂-depletion, a separate photoreceptor is probably also involved. Indeed, end-of-day FR treatment accelerated stomatal closure in *Phaseolus* (Holmes and Klein, 1985). Mosses develop stomata in the epidermis of the sporophyte: whereas they show responses to ABA similar to those of higher plants (Chater et al., 2011), light effects have not been studied. HIP7 might in any case be involved in other signaling pathways involving post-translational phosphorylation. Indeed, *Arabidopsis* HT1 is also expressed in stem, flower and even root tissues, implying diverse but still unknown cellular functions.

HIP8 (Pp3c11_25550V1.1)

Holophytochrome-interacting protein 8 (55.4 kDa) harbors a predicted protein kinase catalytic domain and several calcium-binding EF-hands and thus probably constitutes a CDPK. Y2H implied only weak interaction with phy4, but the signal was enhanced in Rc and stronger with BD:phy4 than with phy4:BD (Supplementary Figure S2). Expression of N- and C-terminal FP-fusions resulted in cytoplasmic and nuclear fluorescence signals approximately with equal signal intensities in both compartments with no apparent effect of light pre-treatment (Figure 7A). *In planta* interaction was verified for all combinations except YFP_C:phy4–HIP8:YFP_N (Figure 7B). Cytoplasmic interaction was clearly enhanced by R pre-treatments in phy4:YFP_N–YFP_C:HIP8, although in phy4:YFP_N–HIP8:YFP_C this was less pronounced. Although this implies that either the phy4–HIP8 complex is enhanced in or translocates to the cytoplasm in R, this behavior was only seen when the tag was fused to phy4 C-terminus: YFP_C:phy4–YFP_N:HIP8 interaction was exclusively cytoplasmic.

Holophytochrome-interacting protein 8 is apparently expressed under all conditions and in all cell types but shows strongly increased transcript levels during leaflet dedifferentiation (see Supplementary Materials). The Genevestigator data for treatments provide some evidence for synexpression with phy4 although this is not apparent for the other datasets.

The eukaryotic second messenger Ca²⁺ is involved in plant responses to various abiotic and biotic stimuli (Kudla et al., 2010), primary targets being Ca²⁺-channels and transporters (Wheeler and Brownlee, 2008). Ca²⁺ from the apoplast or the vacuole thereby briefly increases the local concentration near the plasma membrane or organelle membrane – a phenomenon known as

the calcium signature. Prior to the discovery of Pfr-dependent nuclear translocation and its interaction with the PIF family of transcription factors in the nucleus, Ca²⁺ had been suggested to be a key player in phytochrome signaling, showing crucial roles in chloroplast development and LHC expression, phyA-mediated light regulation of asparagine synthase, R-induced stem protoplast swelling and phyA- and PIF3-mediated regulation of succinate dehydrogenase (Neuhaus et al., 1993, 1997; Bowler et al., 1994; Long and Iino, 2001; Eprintsev et al., 2013). Indeed, there is evidence that phytochromes initiate Ca²⁺ and protein phosphorylation signals via CDPK's (Datta et al., 1985; Li et al., 1991). Furthermore, Ca²⁺ addition to *Cucumis* cotyledon extracts can mimic the R/FR effects on phosphorylation and at least two CDPK's were present whose activities were modulated by phytochromes (Vidal et al., 2007). CDPK's possess a N-terminal variable domain, a protein kinase-, an autoinhibitory and a C-terminal calmodulin-like domain, the latter typically consisting of two pairs of EF-hand motifs (Rutschmann et al., 2002; Christodoulou et al., 2004). CDPKs directly bind Ca²⁺, leading to a conformational change and kinase de-repression, thereby converting the Ca²⁺ signal into phosphorylation of metabolic enzymes, ion channels and transcription factors and thus into diverse downstream cellular responses (reviewed in Hamel et al., 2014). Kinase activity can also be regulated by (auto)phosphorylation/dephosphorylation, phospholipids and 14-3-3-proteins (reviewed in Cheng et al., 2002), however. Interestingly, the holo phy4:BD-screen also identified a 14-3-3 protein (see HIP14). Overall domain organization seems to be conserved in all CDPK's, and land plant CDPK's usually cluster in four distinct clades (Hamel et al., 2014). Comparison of the HIP8 amino acid sequence with the *Viridiplantae* protein database indeed revealed numerous orthologs with high levels of similarity, the *Arabidopsis* and *Physcomitrella* genomes encoding 34 and 25 CDPK's, respectively (Hamel et al., 2014). The only moss representative characterized to-date seems to be associated with nutrient starvation (Mitra and Johri, 2000). Several *Arabidopsis* CDPK's such as CPK1 – the most similar to HIP8 (covering 91% of the query with 65% identity) – are thought to be involved in ABA signaling, regulation of stomatal aperture, and stress responses (Mori et al., 2006; Ma and Wu, 2007; Zhu et al., 2007; Geiger et al., 2010; Zou et al., 2010; Brandt et al., 2012; Boudsocq and Sheen, 2013; Liese and Romeis, 2013). Although CPK1 is membrane-/peroxysome-localized and thus probably involved in oxidative stress and lipid metabolism (Dammann et al., 2003), this is not the case for HIP8 which thus probably has different functions. *Arabidopsis* CPK17 and perhaps other CDPKs on the other hand are involved in Ca²⁺-mediated pollen tube tip growth (Myers et al., 2009; Hamel et al., 2014). As *Physcomitrella* filament tip growth shows many similarities to that of pollen tubes including a tip-focused calcium gradient (Meske and Hartmann, 1995; Meske et al., 1996; see HIP13), we speculate that HIP8 might play a related role in phy4-steered phototropism. On the other hand, whereas *Arabidopsis* CPK17 is plasma membrane-localized (Hamel et al., 2014), this seems not to be the case for HIP8 (Figure 7). However, in both algae and higher plants several CDPKs are associated with the cortical actin cytoskeleton and cytoplasmic streaming (Putnam-Evans et al.,




FIGURE 7 | HIP8 (Pp3c11_25550V1.1) intracellular localization (A) and split-YFP-studies of HIP8–phy4 interaction (B) each without (D) and with R pre-treatment (R) using fluorescence- and confocal microscopy, respectively. Overall figure structure is analogous to **Figure 1**. **(A)** CFP:HIP8- (rows 1 and 2) and HIP8:CFP-fluorescence (rows 3 and 4) was observed in nucleus and cytoplasm (column 1, CFP). **(B)** All four split-YFP configurations were studied, however, in the case of YFP_N:phy4–YFP_C:HIP8 and YFP_N:phy4–HIP8:YFP_C the negative controls showed significant signals (thus these configurations were replaced by YFP_C:phy4–YFP_N:HIP8 and YFP_C:phy4–HIP8:YFP_N). Analyses of YFP_C:phy4–YFP_N:HIP8 (rows 1 and 2), YFP_C:phy4–HIP8:YFP_N (rows 3 and 4), phy4:YFP_N–YFP_C:HIP8 (rows 5–7; two different signal patterns in D) and phy4:YFP_N–HIP8:YFP_C (rows 8–10; two different signal patterns in R) resulted in either nucleo-cytoplasmic or exclusively cytoplasmic signals in all configurations except YFP_C:phy4–HIP8:YFP_N (column 1, YFP). Scale bars 30 μm.

1989; McCurdy and Harmon, 1992). Light-induced cytoplasmic streaming is also mediated by phytochrome in the angiosperm *Vallisneria*. This Ca^{2+} - and actin-dependent response is of particular interest regarding phytochrome cytoplasmic signaling as it is restricted to the area of irradiation and begins within 3 s of irradiation (Takagi et al., 2003), thereby ruling out a transcription-translation-based transduction mechanism. If the functions of actin-associated CDPKs and HIP8 are indeed related, it might represent the molecular link between the directional signal sensed by phytochrome and cytoskeleton-steered tip cell growth/bending response.

HIP9 (Pp1s254_10V6.2)

Holophytochrome-interacting protein 9 (63.9 kDa) harbors a predicted **rhomboid peptidase S54 domain** and predicted transmembrane regions. Y2H studies with phy4 showed weak but R-enhanced interaction with full length HIP9 in all configurations tested (Supplementary Figure S2). *In planta* localization studies showed cytoplasmic and perhaps plasma membrane (CFP:HIP9) or cytoplasmic and nuclear localization with no sign of membrane association (HIP9:CFP; **Figure 8A**). Although the predicted protein size (64 kDa + 27 kDa tag) exceeds the nuclear pore exclusion limit, the C-terminal FP-fusion was able to enter the nucleus. Indeed, two NLS in the N-terminal region are predicted by NucPred⁷ (Brameier et al., 2007), so nuclear import is not unreasonable. Interaction was confirmed *in planta* only in the phy4:YFP_N-HIP9:YFP_C configuration following R pre-treatment (in harmony with the Y2H data; **Figure 8B**). In this case, fluorescence was generally distributed between cytoplasm and nucleus although occasionally a signal at the plasma membrane was apparent.

Holophytochrome-interacting protein 5 is apparently expressed under all conditions and in all cell types (see Supplementary Materials). The data provided no evidence for synexpression with phy4, however.

Rhomboid domain proteases are ubiquitous intramembrane serine proteases (Urban et al., 2001; Koonin et al., 2003; Urban and Wolfe, 2005). According to crystal structures, RBLs possess seven transmembrane helices forming a conical cavity accessible to the aqueous surroundings, enabling cleavage of a transmembrane helix with the help of serine and histidine residues forming a catalytic dyad (Kanaoka et al., 2005; Baker et al., 2007; Ben-Shem et al., 2007). The *Arabidopsis* and *Physcomitrella* genomes encode 17 and 16 RBLs, respectively (Li et al., 2015), including the homolog (Pp3c22_8560V1.1, see Summary Table 1 in the Supplementary Material). Of those in *Arabidopsis*, four show poorly conserved active sites, two localizing to the Golgi apparatus suggesting a role in secretion, while others localize to mitochondria, plastids, and perhaps also the endoplasmic reticulum and the plasma membrane, implying different functions (Kanaoka et al., 2005; Lemberg and Freeman, 2007; Kmiec-Wisniewska et al., 2008; Knopf et al., 2012; Thompson et al., 2012). Transcription factors represent potential targets, ~10% of *Arabidopsis* transcription factors requiring release from membranes to become functional

(Kim and Park, 2007; Seo et al., 2008). RBLs have also been shown to be involved in Ca^{2+} -mediated stress responses, fertility and flower development, structural and functional maintenance of chloroplast and mitochondrion integrity and mitochondrial retrograde signaling (Aguilar-Hernandez et al., 2011; Thompson et al., 2012; Ng et al., 2013). HIP9 is most similar to RBL 9 (At5g38510), predicted to be inactive due to the lack both catalytic dyad amino acids but retaining the glycine within the GxSx rhomboid protease motif, perhaps representing a modified active site (Lemberg and Freeman, 2007). The substrates and functions of this RBL group are not known, however. Thus whether HIP9 is indeed an RBL membrane protein with switchable protease activity is doubtful. In particular, rather than membrane integration, our data implies at most a weak association with the plasma membrane. Whereas this would correlate with the phy4-phot interaction at the plasma membrane (Jaedicke et al., 2012), an involvement of HIP9 in that signaling complex remains no more than a possibility.

HIP11 (Pp3c20_16210V1.1)

Holophytochrome-interacting protein 11, (20.2 kDa) shows low similarity to the *Arabidopsis* pfkB-like carbohydrate kinase family protein AT5G19150. A likely homolog in *Physcomitrella* is Pp3c7_8840V1.1. Holo-BD:phy4 interacted weakly with ADg:HIP11 and HIP11:ADg in Y2H, whereas interactions of holo-phy4:BD with HIP11 were light-enhanced but fully FR-reversible only in case of ADg:HIP11 (Supplementary Figure S2). GFP:HIP11 and HIP11:GFP mostly showed cytoplasmic/perinuclear and rather unstructured cytoplasmic localization, respectively (**Figure 9A**), although in a subset of cells cytoplasmic and nuclear localization could be observed in both cases (not shown). *In planta* interaction was verified in the phy4:YFP_N-YFP_C:HIP11 configuration, the split-YFP signal being apparent in the cytoplasm and perinuclear region (**Figure 9B**) in accordance with the localization data.

Holophytochrome-interacting protein 11 is apparently expressed very differently under different conditions and particularly weakly at early developmental stages (such as in spores and protonemata). There is little evidence for synexpression with phy4.

Although HIP11 was predicted by InterPro to carry an N-terminal transmembrane region (residues 7–25), we found no direct evidence of membrane-association. On the other hand, its small size and an NLS (RRRR, residues 112–115, Brameier et al., 2007) predicted by NucPred are consistent with the occasional nuclear localization we observed. As BLAST searches revealed only 31% identity within the 36% query coverage with respect to *Arabidopsis* AT5G19150 (whose function is unknown), HIP11 and Pp3c7_8840V1.1 might be functionally unique to mosses.

HIP12 (Pp3c13_15620V1.1)

Holophytochrome-interacting protein 12 (17.5 kDa) is the evolutionarily and structurally conserved **eukaryotic translation initiation factor 5A** (eIF-5A). In Y2H studies, interaction of HIP12/eIF-5A with phy4 was relatively weak but nevertheless light-induced, strong and FR-reversible

⁷www.sbc.su.se/~maccallr/nucpred/





FIGURE 8 | HIP9 (Pp1s254_10V6.2) intracellular localization (A) and split-YFP-studies of HIP9–phy4 interaction (B) each without (D) and with R pre-treatment (R) using fluorescence- and confocal microscopy, respectively. Overall figure structure is analogous to **Figure 1**. **(A)** CFP:HIP9- (rows 1 and 2) and HIP9:CFP-fluorescence (rows 3 and 4) was observed in the cytoplasm or the nucleus and cytoplasm, respectively (column 1, CFP). **(B)** All possible split-YFP configurations were investigated: YFP_N:phy4–YFP_C:HIP9 (rows 1 and 2), YFP_N:phy4–HIP9:YFP_C (rows 3 and 4), phy4:YFP_N–YFP_C:HIP9 (rows 5 and 6) and phy4:YFP_N–HIP9:YFP_C (rows 7–9; two distinct patterns in R). However, two different signal patterns (nucleus and cytoplasm or perhaps nucleus and plasma membrane) were only observed for phy4:YFP_N–HIP9:YFP_C after R pre-treatment (column 1, YFP; D and R images here from separate experiments). Scale bars 30 μm.



and constitutive in cases of BD:phy4-ADg:HIP12/BD:phy4-HIP12:ADg, phy4:BD-ADg:HIP12, and phy4:BD-HIP12:ADg, respectively (Supplementary Figure S2). Irrespective of light pre-treatments or FP position, HIP12/eIF-5A was localized in the cytoplasm and nucleus (**Figure 10A**), presumably because the 18 kDa + 27 kDa FP chimera would be small enough to enter the nucleus passively, although in some HIP12:GFP-cells nuclear fluorescence was much weaker than in the cytoplasm (data not shown). Interaction was confirmed *in planta* although the site of localization was strongly dependent on the construct used (see **Figure 10B**). Whereas phy4:YFP_N-HIP12:YFP_C-interaction was seen in both nucleus and cytoplasm or cytoplasm only, YFP_N:phy4-YFP_C:HIP12 was apparent only in the cytoplasm. YFP_C:phy4-HIP12:YFP_N on the other hand failed to show significant interaction. phy4:YFP_N-YFP_C:HIP12 was omitted from the figure as the negative control :YFP_N-YFP_C:HIP12 showed substantial fluorescence signals, implying an artifactual interaction.

Holophytochrome-interacting protein 12 is apparently strongly expressed in under most conditions and in most cell types except leaflets (see Supplementary Materials). According to Genevestigator HIP12 expression is only weakly affected by different treatments but increases strongly on dedifferentiation of the leaflet cells (see Supplementary Materials). The Genevestigator treatment data alone provided evidence for synexpression with phy4.

Basic local alignment search tool analysis revealed a likely HIP12 homolog in *Physcomitrella* (Pp3c12_3540V1.1) and three in *Arabidopsis*. Despite its name, eIF-5A is now generally considered to be involved in translation elongation rather than initiation (Kemper et al., 1976; Gregio et al., 2009; Patel et al., 2009; Saini et al., 2009; Tong et al., 2009). The association of phy4 with eIF-5A and thus translation provides an interesting possible connection to the reported association of phyB Pfr with the cytoplasmic translational regulator PENTA1 and associated inhibition of PORA translation (Paik et al., 2012). Interestingly, in fission yeast eIF-5A has been associated with organization of the cytoskeleton, cell polarity and bud formation (Weir and Yaffe, 2004; Zanelli and Valentini, 2005; Chatterjee et al., 2006), providing a potential link to steering of *Physcomitrella* tip cell growth by phy4 (see also HIP13 below).

The complex localization behavior of HIP12 we observed *in planta* correlates with observations in other systems. Mammalian eIF-5A is localized in the cytoplasm and perinuclear region where it interacts with ribosomes attached to the endoplasmic reticulum, forming a network-like structure (Shi et al., 1996) resembling the pattern we observed (see **Figure 10**). Post-translational conversion of a lysine residue to hypusine (Chen and Liu, 1997; Park et al., 1997) regulates eIF5-A's activity and localization, however, the precursor being found in both cytoplasm and nucleus whereas the hypusinated eIF-5A localizes predominantly to the cytoplasm (Park et al., 1982; Chen, 1983; Torrelío et al., 1984; Park, 1989; Lee et al., 2009). In maize, phosphorylation of the terminal serine changes eIF-5A-2 localization, unphosphorylatable mutants localizing to nucleus and cytoplasm, while phospho-mimic mutants sequester to the nucleus (Lebska et al., 2010; Lewandowska-Gnatowska

et al., 2011). Indeed it was suggested that the phyA-associated phosphatase PP2A (see HIP14 below) is responsible for the dephosphorylation. In *Arabidopsis* several isoforms have been identified and associated with xylem formation, abiotic stress tolerance and responses to cytokinin including senescence (Chamot and Kuhlemeier, 1992; Wang et al., 2001, 2012; Thompson et al., 2004; Feng et al., 2007; Liu et al., 2008; Ren et al., 2013). HIP12 is most similar to eIF-5A-3, however, which is preferentially expressed in the phloem but whose function is not known (Ma et al., 2010).

HIP13 (Pp3c1_23670V1.1)

Holophytochrome-interacting protein 13 (49.3 kDa) is the eukaryotic/archaeal **translation elongation factor subunit EF1 α** . Y2H studies showed constitutive interaction with HIP13/EF1 α in case of BD:phy4 and rather weak but R-enhanced interaction with phy4:BD, respectively (Supplementary Figure S2). Irrespective of FP-fusion configuration or light treatment, HIP13/EF1 α localized to the cytoplasm (**Figure 11A**) and perinuclear region. Accordingly, most split-YFP signals also constitutively appeared in the cytoplasm and perinuclear region (YFP_C:phy4-YFP_N:HIP13 and phy4:YFP_N-YFP_C:HIP13, **Figure 11B**). In particular, YFP_C:phy4-YFP_N:HIP13 cytoplasmic signals resembled cytoskeleton-like substructures. phy4:YFP_C-HIP13:YFP_N signals were additionally seen in the nucleus, however. As the negative controls YFP_C: or YFP_N: with HIP13:YFP_N or HIP13:YFP_C, respectively, produced substantial signals in the cytoplasm and/or nucleus, the respective combinations with N-terminally fused phy4 were categorized as false positive signals and excluded from discussion.

Holophytochrome-interacting protein 13 is apparently expressed strongly under most conditions and in most cell types but weakly in spores and leaflets (see Supplementary Materials). There is little evidence for synexpression with phy4.

Translation elongation factor 1 α is an extremely conserved multifunctional protein harboring an N-terminal RAS-like GTPase (also called p-loop NTPase) and C-terminal GTP EFTU-D2 and -D3 domains. The RAS-like GTPase domain comprises several nucleotide-binding sites whereas the D3-domain is responsible for actin binding (Gross and Kinzy, 2005). As the name implies, EF1 α plays an essential role in translational elongation, thus interaction with phy4 might relate to phytochrome-regulated PENTA1-mediated phytochrome regulation of translation (Paik et al., 2012) as discussed for HIP12. On the other hand, EF1 α has been shown to bind and bundle actin filaments in various species (Demma et al., 1990; Dharmawardhane et al., 1991; Bassell et al., 1994; Collings et al., 1994). EF1 α constitutes 1–2% of total protein in normal growing cells (Slobin, 1980; Dharmawardhane et al., 1991), more than half of which is estimated to be associated with actin (Edmonds et al., 1995). Interaction with actin is dependent on both EF1 α -GTP binding and phosphorylation status (Dharmawardhane et al., 1991; Yang and Boss, 1994; Izawa et al., 2000). EF1 α activates phosphatidylinositol 4-kinase (Yang et al., 1993) thereby regulating the capping and severing of actin filaments proteins. It also serves as a substrate for Rho-associated and Ca²⁺-dependent protein kinases (Yang




FIGURE 10 | HIP12 (Pp3c13_15620V1.1) intracellular localization (A) and split-YFP-studies of HIP12–phy4 interaction (B) each without (D) and with R pre-treatment (R) using fluorescence- and confocal microscopy, respectively. Overall figure structure is analogous to **Figure 1**. **(A)** N- and C-terminally GFP-tagged HIP12 localized to nucleus and cytoplasm (rows 1 and 2; and 3 and 4, respectively; column 1, GFP). In some cases the HIP12::GFP signal was confined largely to the cytoplasm (not shown). **(B)** All four split-YFP configurations were tested, however, YFP_N::phy4–HIP12::YFP_C and phy4::YFP_N–YFP_C::HIP12 were excluded because the respective negative controls showed significant signals. Thus the following configurations are shown: YFP_N::phy4–YFP_C::HIP12 (rows 1 and 2), YFP_C::phy4–HIP12::YFP_N (rows 3 and 4) and phy4::YFP_N–HIP12::YFP_C (rows 5–8; two distinct patterns in both conditions). All configurations except YFP_C::phy4–HIP12::YFP_N yielded significant cytoplasmic or nucleo-cytoplasmic signals (column 1, YFP). Scale bars 30 μm.




FIGURE 11 | HIP13 (Pp3c1_23670V1.1) intracellular localization (A) and split-YFP studies of HIP13–phy4 interaction (B) each without (D) and with R pre-treatment (R) using fluorescence- and confocal microscopy, respectively. Overall figure structure is analogous to **Figure 1**. **(A)** Both CFP:HIP13 (rows 1 and 2) and HIP13:CFP (rows 3 and 4) were localized in the cytoplasm and perinuclear region (column 1, CFP). **(B)** All four split-YFP configurations were tested, however, YFP_C:phy4–HIP13:YFP_N and phy4:YFP_C–YFP_N:HIP13 were excluded because the respective negative controls showed significant signals. Thus the following configurations are shown: YFP_C:phy4–YFP_N:HIP13 (rows 1 and 2), phy4:YFP_N–YFP_C:HIP13 (rows 3 and 4) and phy4:YFP_C–HIP13:YFP_N (rows 5 and 6), all yielding significant cytoplasmic signals with the exception of phy4:YFP_C–HIP13:YFP_N the signal which could be observed in nucleus and cytoplasm (column 1, YFP). Scalebars 30 μ m.

and Boss, 1994; Izawa et al., 2000). Moreover, EF1 α binds and severs microtubules in a Ca²⁺/calmodulin-dependent manner in various organisms (Durso and Cyr, 1994; Kaur and Ruben, 1994; Shiina et al., 1994). Cytoskeleton-binding of EF1 α probably links the translation machinery to the spatial organization of the cell whereby EF1 α regulates transport and anchorage of mRNA and the compartmentalization of translation in eukaryotic cells (Gross and Kinzy, 2005).

Four and ~11 EF1 α -homologs exist in *Arabidopsis* and *Physcomitrella*, respectively (Ransom-Hodgkins, 2009). EF1 α 's promiscuous actions in the cell, in particular its association with the cytoskeleton and possibly Ca²⁺ signaling, make it an interesting candidate for a role in phy4-mediated phototropism in filament tip cells. In unilateral R the Pfr distribution at the plasma membrane somehow generates a Ca²⁺-gradient at the illuminated side of the cell, shifting the site of F-actin synthesis (Hartmann and Weber, 1988; Meske and Hartmann, 1995; Meske et al., 1996). As a result, a directionally translocated actin cap forms, defining the new center of outgrowth corresponding to polarized vesicle transport along the actin bundles (Schneepf, 1986; Meske et al., 1996). More recently, profilin 2 has been shown to be involved in actin turnover and in actin cytoskeleton-mediated vesicular trafficking together with EF1 α in *Arabidopsis* root cells (Takac et al., 2011). Indeed, profilin is essential for tip cell growth in *Physcomitrella* (Vidali et al., 2007). Profilin together with actin and CHUP1 are also central players in chloroplast movement responses in *Arabidopsis* (Schmidt von Braun, 2008; Schmidt von Braun and Schleiff, 2008). Chloroplast photorelocation is a cytoplasmic directional response mediated by phytochrome in mosses. Given that CHUP1 is also present in *Physcomitrella*, a machinery involving phytochrome, profilin, EF1 α and the actin-filament network can be envisaged. It should be recalled, however, that the equivalent response in *Arabidopsis* is mediated by the blue light receptor phototropin, phytochrome playing a modulating role (DeBlasio et al., 2003; Luesse et al., 2010).

HIP14 (Pp3c3_8540C1.1)


Holophytochrome-interacting protein 14 (29.1 kDa) is a 14-3-3 protein family member. Y2H analysis of phy4 interaction with the full-length protein model confirmed interaction only with the phy4:BD configuration (Supplementary Figure S2). *In planta* GFP:HIP14 localized to cytoplasm and nucleus, whereas the HIP14:GFP configuration was exclusively cytoplasmic showing a cytoskeleton-like pattern (Figure 12A) This pattern was not apparent in the GFP:HIP14 configuration. The HIP14-phy4 interaction was verified *in planta*: phy4:YFP_N-YFP_C:HIP14 on the one hand showed a strong nuclear and cytoplasmic signal, reflecting that of GFP:HIP14, and in addition showed an exclusively cytoplasmic, cytoskeleton-like pattern in a subset of cells (Figure 12B). The reconstituted phy4:YFP_N-HIP14:YFP_C signal, like that of HIP14:GFP, was exclusively cytoplasmic although we did not observe cytoskeleton-like patterns.

Holophytochrome-interacting protein 14 is apparently expressed very differently under different conditions and in different cell types with little evidence for synexpression with phy4 (see Supplementary Materials).

Ubiquitous in eukaryotic cells (Isobe et al., 1991; Yaffe, 2002), 14-3-3 proteins possess a conserved central region involved in dimerization and interaction with client proteins via an amphipathic groove. 14-3-3-targets are mostly phosphothreonine and phosphoserine proteins harboring (R/K)(XX/S)X(S/T)PXP motifs, the conformation of which is affected by 14-3-3 binding, leading to changes in the target's catalytic activity, binding to down-stream signaling components and/or intracellular localization (see de Boer et al., 2013). ~10 other 14-3-3 proteins are encoded in the *Physcomitrella* genome. Comparison with the 13 14-3-3 family members in *Arabidopsis* showed HIP14 to be most similar to isoforms GF14 ψ and ω which are localized predominantly in the trichome nucleus and cytoplasm with cytoskeleton association (Paul et al., 2005), similarly to our *in planta* localization data for HIP14:GFP and split-YFP-data for phy4:YFP_N-YFP_C:HIP14 (see Figures 12A,B). Despite their sequence similarity, plant 14-3-3 proteins execute specific functions on account of cell- or stimulus-specific expression, differential phosphorylation or intracellular localization (Paul et al., 2005). Altogether >750 14-3-3 protein targets have been identified in *Arabidopsis*, thus their functions are diverse, including regulation of hormone signaling and responses to external stimuli such as light (see Denison et al., 2011). Certain 14-3-3 isoforms mediate stomatal opening via interaction with light-activated phototropins and subsequently binding to the phosphorylated/activated plasma membrane H⁺-ATPase (Kinoshita et al., 2003; Ueno et al., 2005; Sullivan et al., 2009; Hayashi et al., 2010, 2011; Tseng et al., 2012), while others may regulate chloroplast movement perhaps also by interaction with phototropin (Sullivan et al., 2009; Tseng et al., 2012). Additionally, 14-3-3 proteins are involved in photoperiodic control of flowering, probably by direct interaction with key players in floral induction such as CONSTANS and FT (florigen; Pnueli et al., 2001; Mayfield et al., 2007; Purwestri et al., 2009). Numerous 14-3-3 clients are involved in light signaling including RPT2, phot2, PP2A, COP1 and -8, CONSTANS-LIKE 1, and PIF3, as well as several other gibberellic acid/phytochrome signaling proteins (Shin et al., 2011). 14-3-3 κ^- and χ^- single and double null mutants resembled PIF3-overexpressors (Mayfield et al., 2007; Adams et al., 2014), implying a role of certain 14-3-3's in plant phytochrome action perhaps by acting as antagonists/regulators of PIF3-dependent light-signaling. Further 14-3-3 clients included PRL1 (see HIP4 above) and phyB (Shin et al., 2011), providing intriguing corollaries to the present study. Moreover, whereas in Rc wild-type *Arabidopsis* seedlings show poor gravitropic orientation, 14-3-3 ν^- null mutants showed enhanced gravitropism, rather like those of *phyB^-* (Mayfield et al., 2007). Indeed, 14-3-3 protein association also with phototropin and cryptochrome signaling might imply involvement in different photoreceptor signaling complexes (see Paul et al., 2008), as was suggested for PKS1, NPH3, and PP2A (see Jaedicke et al., 2012). We speculate that the HIP14-phy4 interaction in *Physcomitrella* might be part of such a complex.

Comparison of phy4 and HIP Expression Data

Generally, partner proteins must be co-expressed (expressed in the same cells at the same times and under the same conditions)



unless the absence of one or other is physiologically relevant. Furthermore, the expression levels of functionally associated genes are often correlated (synexpression). We therefore collected available expression data from eFP, Phytozome and Genevestigator for *phy4* and HIP1-14 to investigate these aspects. *phy4* is expressed to some extent in all analyzed developmental stages and conditions (see Supplementary Materials). Light effects are fairly small (~twofold differences). Pairwise comparison of HIP and *phy4* expression showed correlation implying synexpression in several cases (HIP4–8 and -12). We also performed a hierarchical clustering analysis of all three datasets according to condition and gene similarities to identify HIP's with expression patterns similar to *phy4* (see Supplementary Figure S6). The Genevestigator treatment effects on *phy4* were most similar to HIP4 and -6 followed by -7 and -5. The most similar expression level patterns were for HIP12, -4, and -9 followed by -11 and -13 according to the Phytozome treatment data. The expression patterns of HIP4, -6, and -7 followed by -3 were most similar to *phy4* according to the eFP developmental data and to HIP3 and -6 followed by -4, -7, and -9 according to Genevestigator anatomy data. Taken together, these comparisons imply synexpression of *phy4* in particular with HIP4, -6, and -7, providing indirect support for their possible involvement in *phy4* action.

CONCLUSION

In summary, *phy4* steers directional growth in moss filaments, thereby necessarily signaling in the cytoplasm rather than via transcriptional regulation in the nucleus. How this cytoplasmic signal arises is not known, but it might derive from the physical interaction of *phy4* with phototropin at plasma membrane (Jaedicke et al., 2012). In the present study we identified and characterized a number of novel HIP's, all of which are at least partly located in the cytoplasm as are their interactions with *phy4*. Most are conserved in higher plants. Our findings and their possible significance are discussed in more detail below.

In addition to the *Arabidopsis* *phyA*-FHY1 positive control, most of the HIP's characterized in this work showed significant R-enhanced, FR-reversible binding in Y2H, indicating that at least the phytochrome-typical R/FR photochromicity and associated structural changes were significant in the growth readout. Notably, no FR-enhanced interaction was seen, in harmony with the established dogma that in plants Pr has no physiological function. *phy4*–*phy4* homodimerization, presumably resulting from interactions between DHP-like domains, was also demonstrated. We therefore consider it likely that the full-length holophytochrome hybrid baits employed in our screens were functional and folded similarly to the native photoreceptor. This is rather surprising in the case of the four candidates identified in the BD:*phy4* screen because it is unclear how the presumptive knot around the N-terminal PAS domain (Wagner et al., 2005; Essen et al., 2008) can be formed in the presence of the 17 kDa BD extension. Perhaps the BD and/or the loop through which it passes remain/s largely unfolded even after translation of the GAF domain has begun,

allowing the knot to form normally. Alternatively, the extended peptide chain might fold back on itself to form a hairpin which alone protrudes through the loop. It is also possible that the holoprotein remains functional even if knot formation is blocked.

Despite our earlier discovery of the interaction between *phy4* and phototropin at the plasma membrane *in planta*, we had quite expected PIF-like proteins to be found in the Y2H screens. This was not the case, however. One explanation might be that, in contrast to fragments, full-length holophytochrome bait is unattractive for such prey in the yeast nucleus. The selection for full-length cDNAs in the construction of the library we used might also be a significant factor (61% of the prey sequences from the initial screen represented the full-length CDS according to the corresponding gene model). In the case of *Arabidopsis* *phyB* for example, the interaction seen with the PHL N-terminus was absent for full-length PHL (Endo et al., 2013). Alternatively, PIF-like transcription factors might not be substrates for *phy4*. However, as different Y2H screens with *Arabidopsis* *phyA* fragments found different partners, it is not surprising that we also found others for *Physcomitrella* *phy4*. In any case, even though in the initial screen 10 of the 108 sequences identified were duplicated at least once, the library was not exhaustively screened, so rare *phy4* interactors are quite likely to have been missed.

Of the 14 putative HIP's (full-length CDS gene models showing robust Y2H interaction with *phy4* in at least one of the four possible hybrid configurations) only two failed to show split-YFP interaction *in planta*. This high rate of confirmation is probably partly because our Y2H screening protocol exploited mating rather than double transformation, significantly reducing background activation and thus the likelihood of false positives while enhancing sensitivity. We also suspect that our use of functional Pfr as bait in the initial screens increased the number of true interactors found. On the other hand, despite *phy4*–HIP interaction being apparent in both Y2H and *in planta*, false positives are still possible.

We emphasize that even a proven physical interaction does not necessarily represent a signaling route. Consequently, it is unlikely that all the HIP's identified in this work are involved in *phy4* signaling. None has been studied experimentally and most had not even been annotated. Summarizing our analyses of the HIP sequences:

- (1) Of the HIP's with possible associations with **general signaling** mechanisms, HIP1 is a likely pirin (a protein group suggested to associate *phyA* with G protein signaling), HIP4 and -5 probably form β -propeller structures and thus, like HIP3, might represent an E3 ubiquitin ligase, HIP7 is a likely signaling protein kinase and HIP8 is probably a Ca^{2+} -dependent protein kinase. HIP13 too might be associated with Ca^{2+} signaling.
- (2) HIP12 and -13 are likely to be involved in **translation** regulation. This correlates interestingly with *phyB* regulation of PORA translation by Pfr-dependent interaction with PENTA (Paik et al., 2012). Like HIP3, PENTA is a cytosolic zinc-finger protein.

- (3) Various HIP's might be associated with **posttranslational protein modification**. HIP7, -8, and perhaps -11 are likely Ser/Thr protein kinases, HIP6 is a likely sulfotransferase and HIP9 might represent an endopeptidase.
- (4) HIP4, -5, -12, -13, and -14 are associated in various ways with the **cytoskeleton**. This is of particular significance because phy4-mediated vectorial effects (photo- and polarotropism in filament tip cells as well as chloroplast translocation – see Mittmann et al., 2004) are associated with microfilaments and their organization (Meske and Hartmann, 1995; Meske et al., 1996). In particular the 14-3-3 protein HIP14 is interesting in this regard.

These and other characteristics of the various HIP's identified in this study are hypothetical, providing in many cases no more than hints as to their physiological roles. Obviously, the most appropriate way to clarify their true functions is to knockout the relevant loci by homologous recombination (Schaefer and Zrýd, 1997) as was successful for phy4 itself (Mittmann et al., 2004). This work has been initiated and we have established different physiological assays to study effects on vectorial responses the results of which will clarify the putative involvement of HIP's in phy4 signaling.

As discussed in the Section “Introduction,” whereas there is no doubt that the cytoplasmic signal retaining vectorial information derives from phytochromes in lower plants, it is unclear whether a related function exists in higher plants. Similarly, whereas phy4–phot interaction at the plasma membrane has been established and is likely to be associated with vectorial signaling in mosses, the situation in higher plants is less clear. We hope that useful clues regarding the latter and/or the evolution of phytochrome signaling will appear with elucidation of the moss system.

REFERENCES

- Adams, E., Diaz, C., Hong, J. P., and Shin, R. (2014). 14-3-3 proteins participate in light signaling through association with PHYTOCHROME INTERACTING FACTORS. *Int. J. Mol. Sci.* 15, 22801–22814. doi: 10.3390/ijms151222801
- Adams, M., and Jia, Z. (2005). Structural and biochemical analysis reveal pirins to possess quercetinase activity. *J. Biol. Chem.* 280, 28675–28682. doi: 10.1074/jbc.M501034200
- Agatep, R., Kirkpatrick, R. D., Parchaliuk, D. L., Woods, R. A., and Gietz, R. D. (1998). Transformation of *Saccharomyces cerevisiae* by the lithium acetate/single-stranded carrier DNA/polyethylene glycol protocol. *Tech. Tips Online* 3, 133–137. doi: 10.1016/S1366-2120(08)70121-1
- Aguilar-Hernandez, H. S., Santos, L., Leon-Galvan, F., Barrera-Pacheco, A., Espitia-Rangel, E., De Leon-Rodriguez, A., et al. (2011). Identification of calcium stress induced genes in amaranth leaves through suppression subtractive hybridization. *J. Plant Physiol.* 168, 2102–2109. doi: 10.1016/j.jplph.2011.06.006
- Baker, R. P., Young, K., Feng, L., Shi, Y., and Urban, S. (2007). Enzymatic analysis of a rhomboid intramembrane protease implicates transmembrane helix 5 as the lateral substrate gate. *Proc. Natl. Acad. Sci. U.S.A.* 104, 8257–8262. doi: 10.1073/pnas.0700814104
- Bassell, G. J., Powers, C. M., Taneja, K. L., and Singer, R. H. (1994). Single mRNAs visualized by ultrastructural in situ hybridization are principally localized at actin filament intersections in fibroblasts. *J. Cell Biol.* 126, 863–876. doi: 10.1083/jcb.126.4.863
- Ben-Shem, A., Fass, D., and Bibi, E. (2007). Structural basis for intramembrane proteolysis by rhomboid serine proteases. *Proc. Natl. Acad. Sci. U.S.A.* 104, 462–466. doi: 10.1073/pnas.0609773104
- Bhalerao, R. P., Salchert, K., Bakó, L., Ókrész, L., Szabados, L., Muranaka, T., et al. (1999). Regulatory interaction of PRL1 WD protein with *Arabidopsis* SNF1-like protein kinases. *Proc. Natl. Acad. Sci. U.S.A.* 96, 5322–5327. doi: 10.1073/pnas.96.9.5322
- Borden, K. L., and Freemont, P. S. (1996). The RING finger domain: a recent example of a sequence-structure family. *Curr. Opin. Struct. Biol.* 6, 395–401. doi: 10.1016/S0959-440X(96)80060-1
- Boudsocq, M., and Sheen, J. (2013). CDPKs in immune and stress signaling. *Trends Plant Sci.* 18, 30–40. doi: 10.1016/j.tplants.2012.08.008
- Bowler, C., Neuhaus, G., Yamagata, H., and Chua, N.-H. (1994). Cyclic GMP and calcium mediate phytochrome phototransduction [published erratum appears in *Cell* 1994 Nov 18;79(4):743]. *Cell* 77, 73–81. doi: 10.1016/0092-8674(94)90236-4
- Brameier, M., Krings, A., and Maccallum, R. M. (2007). NucPred—Predicting nuclear localization of proteins. *Bioinformatics* 23, 1159–1160. doi: 10.1093/bioinformatics/btm066
- Brandt, B., Brodsky, D. E., Xue, S., Negi, J., Iba, K., Kangasjarvi, J., et al. (2012). Reconstitution of abscisic acid activation of SLAC1 anion channel by CPK6 and OST1 kinases and branched ABI1 PP2C phosphatase action. *Proc. Natl. Acad. Sci. U.S.A.* 109, 10593–10598. doi: 10.1073/pnas.1116590109
- Buer, C. S., Kordbacheh, F., Truong, T. T., Hocart, C. H., and Djordjevic, M. A. (2013). Alteration of flavonoid accumulation patterns in transparent testa mutants disturbs auxin transport, gravity responses, and imparts

AUTHOR CONTRIBUTIONS

AE, KM, and JH designed research and; AE and KM performed experiments. AE and KM analyzed the data, AE and JH wrote and all authors critically read and corrected the manuscript.

FUNDING

We gratefully acknowledge the financial support of the *Deutsche Forschungsgemeinschaft* (Hu702/5 to JH).

ACKNOWLEDGMENTS

We thank Tanja Gans and Melanie Bingel for excellent technical assistance, Rabea Meyberg and Natalie Kempel for a subset of the localization data, Anne Holz and Adriaan Dorresteijn for use of the confocal microscope, Mathias Zeidler for numerous valuable suggestions and advice (all University of Gießen), Stefan Rensing and Pierre-François Perroud (University of Marburg, Germany) for helpful advice concerning expression data analysis, Hans Sommer (MPI for Plant Breeding Research, Cologne, Germany) for the pre-transformed Y2H cDNA library, Peter Uetz and Jürgen Haas (University of Karlsruhe, Germany) for the Gateway Y2H destination vectors, and Nebion (Zurich, Switzerland) for providing us with the Genevestigator data.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fpls.2016.00613>

- long-term effects on root and shoot architecture. *Planta* 238, 171–189. doi: 10.1007/s00425-013-1883-3
- Cardozo, T., and Pagano, M. (2004). The SCF ubiquitin ligase: insights into a molecular machine. *Nat. Rev. Mol. Cell Biol.* 5, 739–751. doi: 10.1038/nrm1471
- Chamot, D., and Kuhlmeier, C. (1992). Differential expression of genes encoding the hypusine-containing translation initiation factor, eIF-5A, in tobacco. *Nucleic Acids Res.* 20, 665–669. doi: 10.1093/nar/20.4.665
- Chater, C., Kamisugi, Y., Movahedi, M., Fleming, A., Cuming, A. C., Gray, J. E., et al. (2011). Regulatory mechanism controlling stomatal behavior conserved across 400 million years of land plant evolution. *Curr. Biol.* 21, 1025–1029. doi: 10.1016/j.cub.2011.04.032
- Chatterjee, I., Gross, S. R., Kinzy, T. G., and Chen, K. Y. (2006). Rapid depletion of mutant eukaryotic initiation factor 5A at restrictive temperature reveals connections to actin cytoskeleton and cell cycle progression. *Mol. Genet. Genomics* 275, 264–276. doi: 10.1007/s00438-005-0086-4
- Chen, K. Y. (1983). An 18000-dalton protein metabolically labeled by polyamines in various mammalian cell lines. *Biochim. Biophys. Acta* 756, 395–402. doi: 10.1016/0304-4165(83)90350-1
- Chen, K. Y., and Liu, A. Y. (1997). Biochemistry and function of hypusine formation on eukaryotic initiation factor 5A. *Biol. Signals* 6, 105–109. doi: 10.1159/000109115
- Chen, M., and Ni, M. (2006). RED AND FAR-RED INSENSITIVE 2, a RING-domain zinc finger protein, mediates phytochrome-controlled seedling deetiolation responses. *Plant Physiol.* 140, 457–465. doi: 10.1104/pp.105.073163
- Cheng, S.-H., Willmann, M. R., Chen, H.-C., and Sheen, J. (2002). Calcium signaling through protein kinases. The *Arabidopsis* calcium-dependent protein kinase gene family. *Plant Physiol.* 129, 469–485. doi: 10.1104/pp.005645
- Cheung, M. Y., Li, M. W., Yung, Y. L., Wen, C. Q., and Lam, H. M. (2013). The unconventional P-loop NTPase OsYchF1 and its regulator OsGAP1 play opposite roles in salinity stress tolerance. *Plant Cell Environ.* 36, 2008–2020. doi: 10.1111/pce.12108
- Christodoulou, J., Malmendal, A., Harper, J. F., and Chazin, W. J. (2004). Evidence for differing roles for each lobe of the calmodulin-like domain in a calcium-dependent protein kinase. *J. Biol. Chem.* 279, 29092–29100. doi: 10.1074/jbc.M401297200
- Citovsky, V., Lee, L. Y., Vyas, S., Glick, E., Chen, M. H., Vainstein, A., et al. (2006). Subcellular localization of interacting proteins by bimolecular fluorescence complementation in planta. *J. Mol. Biol.* 362, 1120–1131. doi: 10.1016/j.jmb.2006.08.017
- Collings, D. A., Wasteneys, G. O., Miyazaki, M., and Williamson, R. E. (1994). Elongation factor 1 alpha is a component of the subcortical actin bundles of characean algae. *Cell Biol. Int.* 18, 1019–1024. doi: 10.1006/cbir.1994.1025
- Dammann, C., Ichida, A., Hong, B., Romanowsky, S. M., Hrabak, E. M., Harmon, A. C., et al. (2003). Subcellular targeting of nine calcium-dependent protein kinase isoforms from *Arabidopsis*. *Plant Physiol.* 132, 1840–1848. doi: 10.1104/pp.103.020008
- Datta, N., Chen, Y. R., and Roux, S. J. (1985). Phytochrome and calcium stimulation of protein phosphorylation in isolated pea nuclei. *Biochem. Biophys. Res. Commun.* 128, 1403–1408. doi: 10.1016/0006-291X(85)91096-4
- de Boer, A. H., Van Kleeff, P. J., and Gao, J. (2013). Plant 14-3-3 proteins as spiders in a web of phosphorylation. *Protoplasma* 250, 425–440. doi: 10.1007/s00709-012-0437-z
- DeBlasio, S. L., Mullen, J. L., Luesse, D. R., and Hangarter, R. P. (2003). Phytochrome modulation of blue light-induced chloroplast movements in *Arabidopsis*. *Plant Physiol.* 133, 1471–1479. doi: 10.1104/pp.103.029116
- Decker, E. L., Frank, W., Sarnighausen, E., and Reski, R. (2006). Moss systems biology en route: phytohormones in *Physcomitrella* development. *Plant Biol. (Stuttg.)* 8, 397–405. doi: 10.1055/s-2006-923952
- Demarsy, E., and Fankhauser, C. (2009). Higher plants use LOV to perceive blue light. *Curr. Opin. Plant Biol.* 12, 69–74. doi: 10.1016/j.pbi.2008.09.002
- Demma, M., Warren, V., Hock, R., Dharmawardhane, S., and Condeelis, J. (1990). Isolation of an abundant 50,000-dalton actin filament bundling protein from *Dictyostelium* amoebae. *J. Biol. Chem.* 265, 2286–2291.
- Denison, F. C., Paul, A. L., Zupanska, A. K., and Ferl, R. J. (2011). 14-3-3 proteins in plant physiology. *Semin. Cell Dev. Biol.* 22, 720–727. doi: 10.1016/j.semcdb.2011.08.006
- Dharmawardhane, S., Demma, M., Yang, F., and Condeelis, J. (1991). Compartmentalization and actin binding properties of ABP-50: the elongation factor-1 alpha of *Dictyostelium*. *Cell Motil. Cytoskeleton* 20, 279–288. doi: 10.1002/cm.970200404
- Ding, S., Zhang, B., and Qin, F. (2015). *Arabidopsis* RZFP34/CHYR1, a Ubiquitin E3 ligase, regulates stomatal movement and drought tolerance via SnRK2.6-mediated phosphorylation. *Plant Cell* 27, 3228–3244. doi: 10.1105/tpc.15.00321
- Durso, N. A., and Cyr, R. J. (1994). A calmodulin-sensitive interaction between microtubules and a higher plant homolog of elongation factor-1 alpha. *Plant Cell* 6, 893–905. doi: 10.1105/tpc.6.6.893
- Edmonds, B. T., Murray, J., and Condeelis, J. (1995). pH regulation of the F-actin binding properties of *Dictyostelium* elongation factor 1 alpha. *J. Biol. Chem.* 270, 15222–15230. doi: 10.1074/jbc.270.25.15222
- Endo, M., Tanigawa, Y., Murakami, T., Araki, T., and Nagatani, A. (2013). PHYTOCHROME-DEPENDENT LATE-FLOWERING accelerates flowering through physical interactions with phytochrome B and CONSTANS. *Proc. Natl. Acad. Sci. U.S.A.* 110, 18017–18022. doi: 10.1073/pnas.1310631110
- Eprintsev, A. T., Fedorin, D. N., and Igamberdiev, A. U. (2013). Ca(2)(+) is involved in phytochrome A-dependent regulation of the succinate dehydrogenase gene *sdh1-2* in *Arabidopsis*. *J. Plant Physiol.* 170, 1349–1352. doi: 10.1016/j.jplph.2013.04.006
- Essen, L.-O., Mailliet, J., and Hughes, J. (2008). The structure of a complete phytochrome sensory module in the Pr ground state. *Proc. Natl. Acad. Sci. U.S.A.* 105, 14709–14714. doi: 10.1073/pnas.0806477105
- Fankhauser, C., Yeh, K. C., Lagarias, J. C., Zhang, H., Elich, T. D., and Chory, J. (1999). PKS1, a substrate phosphorylated by phytochrome that modulates light signaling in *Arabidopsis*. *Science* 284, 1539–1541. doi: 10.1126/science.284.5419.1539
- Faulkner, I. J., and Rubery, P. H. (1992). Flavonoids and flavonoid sulphates as probes of auxin-transport regulation in *Cucurbita pepo* hypocotyl segments and vesicles. *Planta* 186, 618–625. doi: 10.1007/BF00198044
- Feierbach, B., Verde, F., and Chang, F. (2004). Regulation of a formin complex by the microtubule plus end protein tea1p. *J. Cell Biol.* 165, 697–707. doi: 10.1083/jcb.200403090
- Feng, H., Chen, Q., Feng, J., Zhang, J., Yang, X., and Zuo, J. (2007). Functional characterization of the *Arabidopsis* eukaryotic translation initiation factor 5A-2 that plays a crucial role in plant growth and development by regulating cell division, cell growth, and cell death. *Plant Physiol.* 144, 1531–1545. doi: 10.1104/pp.107.098079
- Fenyk, S., Campillo Ade, S., Pohl, E., Hussey, P. J., and Cann, M. J. (2012). A nucleotide phosphatase activity in the nucleotide binding domain of an orphan resistance protein from rice. *J. Biol. Chem.* 287, 4023–4032. doi: 10.1074/jbc.M111.314450
- Fischer, C., Speth, V., Fleig-Eberenz, S., and Neuhaus, G. (1997). Induction of zygotic polyembryos in wheat: influence of auxin polar transport. *Plant Cell* 9, 1767–1780. doi: 10.1105/tpc.9.10.1767
- Flores-Perez, U., Perez-Gil, J., Closa, M., Wright, L. P., Botella-Pavia, P., Phillips, M. A., et al. (2010). Pleiotropic regulatory locus 1 (PRL1) integrates the regulation of sugar responses with isoprenoid metabolism in *Arabidopsis*. *Mol. Plant* 3, 101–112. doi: 10.1093/mp/ssp100
- Franciosini, A., Lombardi, B., Iafrate, S., Pecce, V., Mele, G., Lupacchini, L., et al. (2013). The *Arabidopsis* COP9 SIGNALOSOME INTERACTING F-BOX KELCH 1 protein forms an SCF ubiquitin ligase and regulates hypocotyl elongation. *Mol. Plant* 6, 1616–1629. doi: 10.1093/mp/ssp045
- Freemont, P. S. (1993). The RING finger. A novel protein sequence motif related to the zinc finger. *Ann. N. Y. Acad. Sci.* 684, 174–192. doi: 10.1111/j.1749-6632.1993.tb32280.x
- Geiger, D., Scherzer, S., Mumm, P., Marten, I., Ache, P., Matschi, S., et al. (2010). Guard cell anion channel SLAC1 is regulated by CDPK protein kinases with distinct Ca²⁺ affinities. *Proc. Natl. Acad. Sci. U.S.A.* 107, 8023–8028. doi: 10.1073/pnas.0912030107
- Gidda, S. K., Miersch, O., Levitin, A., Schmidt, J., Wasternack, C., and Varin, L. (2003). Biochemical and molecular characterization of a hydroxyjasmonate sulfotransferase from *Arabidopsis thaliana*. *J. Biol. Chem.* 278, 17895–17900. doi: 10.1074/jbc.M211943200

- Gietz, R. D., and Woods, R. A. (2002). Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Methods Enzymol.* 350, 87–96. doi: 10.1016/S0076-6879(02)50957-5
- Gregio, A. P., Cano, V. P., Avaca, J. S., Valentini, S. R., and Zanelli, C. F. (2009). eIF5A has a function in the elongation step of translation in yeast. *Biochem. Biophys. Res. Commun.* 380, 785–790. doi: 10.1016/j.bbrc.2009.01.148
- Gross, S. R., and Kinzy, T. G. (2005). Translation elongation factor 1A is essential for regulation of the actin cytoskeleton and cell morphology. *Nat. Struct. Mol. Biol.* 12, 772–778. doi: 10.1038/nsmb979
- Hamel, L. P., Sheen, J., and Seguin, A. (2014). Ancient signals: comparative genomics of green plant CDPKs. *Trends Plant Sci.* 19, 79–89. doi: 10.1016/j.tplants.2013.10.009
- Harmon, F. G., and Kay, S. A. (2003). The F box protein AFR is a positive regulator of phytochrome A-mediated light signaling. *Curr. Biol.* 13, 2091–2096. doi: 10.1016/j.cub.2003.11.019
- Hartmann, E., and Weber, M. (1988). Storage of the phytochrome-mediated phototropic stimulus of moss protonematal cells. *Planta* 175, 39–49. doi: 10.1007/BF00402880
- Hashimoto, M., Negi, J., Young, J., Israelsson, M., Schroeder, J. I., and Iba, K. (2006). *Arabidopsis* HT1 kinase controls stomatal movements in response to CO₂. *Nat. Cell Biol.* 8, 391–397. doi: 10.1038/ncb1387
- Hayashi, M., Inoue, S., Takahashi, K., and Kinoshita, T. (2011). Immunohistochemical detection of blue light-induced phosphorylation of the plasma membrane H⁺-ATPase in stomatal guard cells. *Plant Cell Physiol.* 52, 1238–1248. doi: 10.1093/pcp/pcr072
- Hayashi, Y., Nakamura, S., Takemiya, A., Takahashi, Y., Shimazaki, K., and Kinoshita, T. (2010). Biochemical characterization of in vitro phosphorylation and dephosphorylation of the plasma membrane H⁺-ATPase. *Plant Cell Physiol.* 51, 1186–1196. doi: 10.1093/pcp/pcq078
- Hectors, K., Van Oevelen, S., Geuns, J., Guisez, Y., Jansen, M. A., and Prinsen, E. (2014). Dynamic changes in plant secondary metabolites during UV acclimation in *Arabidopsis thaliana*. *Physiol. Plant.* 152, 219–230. doi: 10.1111/pp.12168
- Hiltbrunner, A., Viczian, A., Bury, E., Tscheuschler, A., Kircher, S., Toth, R., et al. (2005). Nuclear accumulation of the phytochrome A photoreceptor requires PHY1. *Curr. Biol.* 15, 2125–2130. doi: 10.1016/j.cub.2005.10.042
- Hirschmann, F., Krause, F., and Papenbrock, J. (2014). The multi-protein family of sulfotransferases in plants: composition, occurrence, substrate specificity, and functions. *Front. Plant Sci.* 5:556. doi: 10.3389/fpls.2014.00556
- Holmes, M. G., and Klein, W. H. (1985). Evidence for phytochrome involvement in light-mediated stomatal movement in *Phaseolus vulgaris* L. *Planta* 166, 348–353. doi: 10.1007/BF00401172
- Hua, Z., and Vierstra, R. D. (2011). The cullin-RING ubiquitin-protein ligases. *Annu. Rev. Plant Biol.* 62, 299–334. doi: 10.1146/annurev-arplant-042809-112256
- Hughes, J. (2013). Phytochrome cytoplasmic signaling. *Annu. Rev. Plant Biol.* 64, 377–402. doi: 10.1146/annurev-arplant-050312-120045
- Huseby, S., Koprivova, A., Lee, B. R., Saha, S., Mithen, R., Wold, A. B., et al. (2013). Diurnal and light regulation of sulphur assimilation and glucosinolate biosynthesis in *Arabidopsis*. *J. Exp. Bot.* 64, 1039–1048. doi: 10.1093/jxb/ers378
- Isobe, T., Ichimura, T., Sunaya, T., Okuyama, T., Takahashi, N., Kuwano, R., et al. (1991). Distinct forms of the protein kinase-dependent activator of tyrosine and tryptophan hydroxylases. *J. Mol. Biol.* 217, 125–132. doi: 10.1016/0022-2836(91)90616-E
- Izawa, T., Fukata, Y., Kimura, T., Iwamatsu, A., Dohi, K., and Kaibuchi, K. (2000). Elongation factor-1 alpha is a novel substrate of rho-associated kinase. *Biochem. Biophys. Res. Commun.* 278, 72–78. doi: 10.1006/bbrc.2000.3772
- Jaedicke, K., Lichtenthaler, A. L., Meyberg, R., Zeidler, M., and Hughes, J. (2012). A phytochrome-phototropin light signaling complex at the plasma membrane. *Proc. Natl. Acad. Sci. U.S.A.* 109, 12231–12236. doi: 10.1073/pnas.1120203109
- Ji, H., Wang, S., Li, K., Szakonyi, D., Koncz, C., and Li, X. (2015). PRL1 modulates root stem cell niche activity and meristem size through WOX5 and PLTs in *Arabidopsis*. *Plant J.* 81, 399–412. doi: 10.1111/tpj.12733
- Jones, A. M., Ecker, J. R., and Chen, J. G. (2003). A reevaluation of the role of the heterotrimeric G protein in coupling light responses in *Arabidopsis*. *Plant Physiol.* 131, 1623–1627. doi: 10.1104/pp.102.017624
- Kadota, A., Sato, Y., and Wada, M. (2000). Intracellular chloroplast photorelocation in the moss *Physcomitrella patens* is mediated by phytochrome as well as by a blue-light receptor. *Planta* 210, 932–937. doi: 10.1007/s004250050700
- Kami, C., Hersch, M., Trevisan, M., Genoud, T., Hiltbrunner, A., Bergmann, S., et al. (2012). Nuclear phytochrome A signaling promotes phototropism in *Arabidopsis*. *Plant Cell* 24, 566–576. doi: 10.1105/tpc.111.095083
- Kanaoka, M. M., Urban, S., Freeman, M., and Okada, K. (2005). An *Arabidopsis* Rhomboid homolog is an intramembrane protease in plants. *FEBS Lett.* 579, 5723–5728. doi: 10.1016/j.febslet.2005.09.049
- Kanegae, T., Hayashida, E., Kuramoto, C., and Wada, M. (2006). A single chromoprotein with triple chromophores acts as both a phytochrome and a phototropin. *Proc. Natl. Acad. Sci. U.S.A.* 103, 17997–18001. doi: 10.1073/pnas.0603569103
- Kanegae, T., and Kimura, I. (2015). A phytochrome/phototropin chimeric photoreceptor of fern functions as a blue/far-red light-dependent photoreceptor for phototropism in *Arabidopsis*. *Plant J.* 83, 480–488. doi: 10.1111/tpj.12903
- Karimi, M., Inzé, D., and Depicker, A. (2002). GATEWAY (TM) vectors for *Agrobacterium*-mediated plant transformation. *Trends Plant Sci.* 7, 193–195. doi: 10.1016/S1360-1385(02)02251-3
- Kaur, K. J., and Ruben, L. (1994). Protein translation elongation factor-1 alpha from *Trypanosoma brucei* binds calmodulin. *J. Biol. Chem.* 269, 23045–23050.
- Kemper, W. M., Merrick, W. C., Redfield, B., Liu, C. K., and Weissbach, H. (1976). Purification and properties of rabbit reticulocyte elongation factor 1. *Arch. Biochem. Biophys.* 174, 603–612. doi: 10.1016/0003-9861(76)90389-1
- Kim, S. G., and Park, C. M. (2007). Membrane-mediated salt stress signaling in flowering time control. *Plant Signal. Behav.* 2, 517–518. doi: 10.4161/psb.2.6.4645
- Kinoshita, T., Emi, T., Tominaga, M., Sakamoto, K., Shigenaga, A., Doi, M., et al. (2003). Blue-light- and phosphorylation-dependent binding of a 14-3-3 protein to phototropins in stomatal guard cells of broad bean. *Plant Physiol.* 133, 1453–1463. doi: 10.1104/pp.103.029629
- Klein, M., and Papenbrock, J. (2004). The multi-protein family of *Arabidopsis* sulphotransferases and their relatives in other plant species. *J. Exp. Bot.* 55, 1809–1820. doi: 10.1093/jxb/erh183
- Klein, M., and Papenbrock, J. (2009). Kinetics and substrate specificities of desulfoglucosinolate sulfotransferases in *Arabidopsis thaliana*. *Physiol. Plant.* 135, 140–149. doi: 10.1111/j.1399-3054.2008.01182.x
- Kmieć-Wisniewska, B., Krumpe, K., Urantowka, A., Sakamoto, W., Pratić, E., and Janska, H. (2008). Plant mitochondrial rhomboid, ATRBL12, has different substrate specificity from its yeast counterpart. *Plant Mol. Biol.* 68, 159–171. doi: 10.1007/s11103-008-9359-8
- Knopf, R. R., Feder, A., Mayer, K., Lin, A., Rozenberg, M., Schaller, A., et al. (2012). Rhomboid proteins in the chloroplast envelope affect the level of allene oxide synthase in *Arabidopsis thaliana*. *Plant J.* 72, 559–571. doi: 10.1111/j.1365-313X.2012.05090.x
- Komori, R., Amano, Y., Ogawa-Ohnishi, M., and Matsubayashi, Y. (2009). Identification of tyrosylprotein sulfotransferase in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 106, 15067–15072. doi: 10.1073/pnas.0902801106
- Koonin, E. V., Makarova, K. S., Rogozin, I. B., Davidovic, L., Letellier, M. C., and Pellegrini, L. (2003). The rhomboids: a nearly ubiquitous family of intramembrane serine proteases that probably evolved by multiple ancient horizontal gene transfers. *Genome Biol.* 4:R19.
- Koonin, E. V., Wolf, Y. I., and Aravind, L. (2000). Protein fold recognition using sequence profiles and its application in structural genomics. *Adv. Protein Chem.* 54, 245–275. doi: 10.1016/S0065-3233(00)54008-X
- Kosarev, P., Mayer, K. F., and Hardtke, C. S. (2002). Evaluation and classification of RING-finger domains encoded by the *Arabidopsis* genome. *Genome Biol.* 3, RESEARCH0016. doi: 10.1186/gb-2002-3-4-research0016
- Kudla, J., Batistic, O., and Hashimoto, K. (2010). Calcium signals: the lead currency of plant information processing. *Plant Cell* 22, 541–563. doi: 10.1105/tpc.109.072686
- Kuhn, B. M., Geisler, M., Bigler, L., and Ringli, C. (2011). Flavonols accumulate asymmetrically and affect auxin transport in *Arabidopsis*. *Plant Physiol.* 156, 585–595. doi: 10.1104/pp.111.175976
- Laity, J. H., Lee, B. M., and Wright, P. E. (2001). Zinc finger proteins: new insights into structural and functional diversity. *Curr. Opin. Struct. Biol.* 11, 39–46. doi: 10.1016/S0959-440X(00)00167-6

- Lapik, Y. R., and Kaufman, L. S. (2003). The *Arabidopsis* cupin domain protein AtPirin1 interacts with the G protein alpha-subunit GPA1 and regulates seed germination and early seedling development. *Plant Cell* 15, 1578–1590. doi: 10.1105/tpc.011890
- Lebska, M., Ciesielski, A., Szymona, L., Godecka, L., Lewandowska-Gnatowska, E., Szczegieliński, J., et al. (2010). Phosphorylation of maize eukaryotic translation initiation factor 5A (eIF5A) by casein kinase 2: identification of phosphorylated residue and influence on intracellular localization of eIF5A. *J. Biol. Chem.* 285, 6217–6226. doi: 10.1074/jbc.M109.018770
- Lee, J. H., Terzaghi, W., Gusmaroli, G., Charron, J. B., Yoon, H. J., Chen, H., et al. (2008). Characterization of *Arabidopsis* and rice DWD proteins and their roles as substrate receptors for CUL4-RING E3 ubiquitin ligases. *Plant Cell* 20, 152–167. doi: 10.1105/tpc.107.055418
- Lee, S. B., Park, J. H., Kaevel, J., Sramkova, M., Weigert, R., and Park, M. H. (2009). The effect of hypusine modification on the intracellular localization of eIF5A. *Biochem. Biophys. Res. Commun.* 383, 497–502. doi: 10.1016/j.bbrc.2009.04.049
- Lemberg, M. K., and Freeman, M. (2007). Functional and evolutionary implications of enhanced genomic analysis of rhomboid intramembrane proteases. *Genome Res.* 17, 1634–1646. doi: 10.1101/gr.6425307
- Leng, R. P., Lin, Y., Ma, W., Wu, H., Lemmers, B., Chung, S., et al. (2003). Pirh2, a p53-induced ubiquitin-protein ligase, promotes p53 degradation. *Cell* 112, 779–791. doi: 10.1016/S0092-8674(03)00193-4
- Lewandowska-Gnatowska, E., Szymona, L., Lebska, M., Szczegieliński, J., and Muszynska, G. (2011). Phosphorylation of maize eukaryotic translation initiation factor on Ser2 by catalytic subunit CK2. *Mol. Cell. Biochem.* 356, 241–244. doi: 10.1007/s11010-011-0952-9
- Lewis, D. R., Ramirez, M. V., Miller, N. D., Vallabhaneni, P., Ray, W. K., Helm, R. F., et al. (2011). Auxin and ethylene induce flavonol accumulation through distinct transcriptional networks. *Plant Physiol.* 156, 144–164. doi: 10.1104/pp.111.172502
- Li, H., Dauwalder, M., and Roux, S. J. (1991). Partial purification and characterization of a Ca(2+)-dependent protein kinase from pea nuclei. *Plant Physiol.* 96, 720–727. doi: 10.1104/pp.96.3.720
- Li, Q., Zhang, N., Zhang, L., and Ma, H. (2015). Differential evolution of members of the rhomboid gene family with conservative and divergent patterns. *New Phytol.* 206, 368–380. doi: 10.1111/nph.13174
- Liese, A., and Romeis, T. (2013). Biochemical regulation of in vivo function of plant calcium-dependent protein kinases (CDPK). *Biochim. Biophys. Acta* 1833, 1582–1589. doi: 10.1016/j.bbamcr.2012.10.024
- Liu, Z., Duguay, J., Ma, F., Wang, T. W., Tshin, R., Hopkins, M. T., et al. (2008). Modulation of eIF5A1 expression alters xylem abundance in *Arabidopsis thaliana*. *J. Exp. Bot.* 59, 939–950. doi: 10.1093/jxb/ern017
- Long, C., and Iino, M. (2001). Light-dependent osmoregulation in pea stem protoplasts. photoreceptors, tissue specificity, ion relationships, and physiological implications. *Plant Physiol.* 125, 1854–1869. doi: 10.1104/pp.125.4.1854
- Luesse, D. R., Deblasio, S. L., and Hangarter, R. P. (2010). Integration of Phot1, Phot2, and PhyB signalling in light-induced chloroplast movements. *J. Exp. Bot.* 61, 4387–4397. doi: 10.1093/jxb/erq242
- Ma, F., Liu, Z., Wang, T. W., Hopkins, M. T., Peterson, C. A., and Thompson, J. E. (2010). *Arabidopsis* eIF5A3 influences growth and the response to osmotic and nutrient stress. *Plant Cell Environ.* 33, 1682–1696. doi: 10.1111/j.1365-3040.2010.02173.x
- Ma, S. Y., and Wu, W. H. (2007). AtCPK23 functions in *Arabidopsis* responses to drought and salt stresses. *Plant Mol. Biol.* 65, 511–518. doi: 10.1007/s11103-007-9187-2
- Marino, D., Froidure, S., Canonne, J., Ben Khaled, S., Khafif, M., Pouzet, C., et al. (2013). *Arabidopsis* ubiquitin ligase MIEL1 mediates degradation of the transcription factor MYB30 weakening plant defence. *Nat. Commun.* 4, 1476. doi: 10.1038/ncomms2479
- Marsolais, F., Boyd, J., Paredes, Y., Schinas, A. M., Garcia, M., Elzein, S., et al. (2007). Molecular and biochemical characterization of two brassinosteroid sulfotransferases from *Arabidopsis*, AtST4a (At2g14920) and AtST1 (At2g03760). *Planta* 225, 1233–1244. doi: 10.1007/s00425-006-0413-y
- Mata, J., and Nurse, P. (1997). Tea1 and the microtubular cytoskeleton are important for generating global spatial order within the fission yeast cell. *Cell* 89, 939–949. doi: 10.1016/S0092-8674(00)80279-2
- Matrosova, A., Bogireddi, H., Mateo-Penas, A., Hashimoto-Sugimoto, M., Iba, K., Schroeder, J. I., et al. (2015). The HT1 protein kinase is essential for red light-induced stomatal opening and genetically interacts with OST1 in red light and CO₂-induced stomatal movement responses. *New Phytol.* 208, 1126–1137. doi: 10.1111/nph.13566
- Matthews, J. M., and Sunde, M. (2002). Zinc fingers—folds for many occasions. *IUBMB Life* 54, 351–355. doi: 10.1080/15216540216035
- Mayfield, J. D., Folta, K. M., Paul, A. L., and Ferl, R. J. (2007). The 14-3-3 Proteins mu and upsilon influence transition to flowering and early phytochrome response. *Plant Physiol.* 145, 1692–1702. doi: 10.1104/pp.107.108654
- McCurdy, D. W., and Harmon, A. C. (1992). Calcium-dependent protein kinase in the green alga *Chara*. *Planta* 188, 54–61. doi: 10.1007/BF00198939
- Meske, V., and Hartmann, E. (1995). Reorganisation of microfilaments in protonemal tip cells of the moss *Ceratodon purpureus* during the phototropic response. *Protoplasma* 188, 59–69. doi: 10.1007/BF01276796
- Meske, V., Rupert, V., and Hartmann, E. (1996). Structural basis for the red light induced repolarisation of tip growth in caulonemal cells of *Ceratodon purpureus*. *Protoplasma* 192, 189–198. doi: 10.1007/BF01273891
- Milner-White, E. J., Coggins, J. R., and Anton, I. A. (1991). Evidence for an ancestral core structure in nucleotide-binding proteins with the type A motif. *J. Mol. Biol.* 221, 751–754. doi: 10.1016/0022-2836(91)80170-Y
- Mitra, D., and Johri, M. M. (2000). Enhanced expression of a calcium-dependent protein kinase from the moss *Funaria hygrometrica* under nutritional starvation. *J. Biosci.* 25, 331–338. doi: 10.1007/BF02703786
- Mittmann, F., Brücker, G., Zeidler, M., Repp, A., Abts, T., Hartmann, E., et al. (2004). Targeted knockout in *Physcomitrella* reveals direct actions of phytochrome in the cytoplasm. *Proc. Natl. Acad. Sci. U.S.A.* 101, 13939–13944. doi: 10.1073/pnas.0403140101
- Mittmann, F., Dienstbach, S., Weisert, A., and Forreiter, C. (2009). Analysis of the phytochrome gene family in *Ceratodon purpureus* by gene targeting reveals the primary phytochrome responsible for photo- and polarotropism. *Planta* 230, 27–37. doi: 10.1007/s00425-009-0922-6
- Mori, I. C., Murata, Y., Yang, Y., Munemasa, S., Wang, Y. F., Andreoli, S., et al. (2006). CDPKs CPK6 and CPK3 function in ABA regulation of guard cell S-type anion- and Ca(2+)-permeable channels and stomatal closure. *PLoS Biol.* 4:e327. doi: 10.1371/journal.pbio.0040327
- Myers, C., Romanowsky, S. M., Barron, Y. D., Garg, S., Azuse, C. L., Curran, A., et al. (2009). Calcium-dependent protein kinases regulate polarized tip growth in pollen tubes. *Plant J.* 59, 528–539. doi: 10.1111/j.1365-313X.2009.03894.x
- Nemeth, K., Salchert, K., Putnok, P., Bhalerao, R., Koncz-Kalman, Z., Stankovic-Stangeland, B., et al. (1998). Pleiotropic control of glucose and hormone responses by PRL1, a nuclear WD protein, in *Arabidopsis*. *Genes Dev.* 12, 3059–3073. doi: 10.1101/gad.12.19.3059
- Neuhaus, G., Bowler, C., Hiratsuka, K., Yamagata, H., and Chua, N.-H. (1997). Phytochrome-regulated repression of gene expression requires calcium and cGMP. *EMBO J.* 16, 2554–2564. doi: 10.1093/emboj/16.10.2554
- Neuhaus, G., Bowler, C., Kern, R., and Chua, N.-H. (1993). Calcium/calmodulin-dependent and -independent phytochrome signal transduction pathways. *Cell* 73, 937–952. doi: 10.1016/0092-8674(93)90272-R
- Ng, S., Ivanova, A., Duncan, O., Law, S. R., Van Aken, O., De Clercq, I., et al. (2013). A membrane-bound NAC transcription factor, ANAC017, mediates mitochondrial retrograde signaling in *Arabidopsis*. *Plant Cell* 25, 3450–3471. doi: 10.1105/tpc.113.113985
- Nozue, K., Kanegae, T., Imaizumi, T., Fukuda, S., Okamoto, H., Yeh, K. C., et al. (1998). A phytochrome from the fern *Adiantum* with features of the putative photoreceptor NPH1. *Proc. Natl. Acad. Sci. U.S.A.* 95, 15826–15830. doi: 10.1073/pnas.95.26.15826
- Okamoto, H., Matsui, M., and Deng, X. W. (2001). Overexpression of the heterotrimeric G-protein alpha-subunit enhances phytochrome-mediated inhibition of hypocotyl elongation in *Arabidopsis*. *Plant Cell* 13, 1639–1652. doi: 10.2307/3871391
- Orozco-Nunnelly, D. A., Muhammad, D., Mezzich, R., Lee, B. S., Jayathilaka, L., Kaufman, L. S., et al. (2014). Pirin1 (PRN1) is a multifunctional protein that regulates quercetin, and impacts specific light and UV responses in the seed-to-seedling transition of *Arabidopsis thaliana*. *PLoS ONE* 9:e93371. doi: 10.1371/journal.pone.0093371

- Orzaez, D., De Jong, A. J., and Woltering, E. J. (2001). A tomato homologue of the human protein PIRIN is induced during programmed cell death. *Plant Mol. Biol.* 46, 459–468. doi: 10.1023/A:1010618515051
- Paik, I., Yang, S., and Choi, G. (2012). Phytochrome regulates translation of mRNA in the cytosol. *Proc. Natl. Acad. Sci. U.S.A.* 109, 1335–1340. doi: 10.1073/pnas.1109683109
- Pang, H., Bartlam, M., Zeng, Q., Miyatake, H., Hisano, T., Miki, K., et al. (2004). Crystal structure of human pirin: an iron-binding nuclear protein and transcription cofactor. *J. Biol. Chem.* 279, 1491–1498. doi: 10.1074/jbc.M310022200
- Park, M. H. (1989). The essential role of hypusine in eukaryotic translation initiation factor 4D (eIF-4D). Purification of eIF-4D and its precursors and comparison of their activities. *J. Biol. Chem.* 264, 18531–18535.
- Park, M. H., Cooper, H. L., and Folk, J. E. (1982). The biosynthesis of protein-bound hypusine (N epsilon -(4-amino-2-hydroxybutyl)lysine). Lysine as the amino acid precursor and the intermediate role of deoxyhypusine (N epsilon -(4-aminobutyl)lysine). *J. Biol. Chem.* 257, 7217–7222.
- Park, M. H., Lee, Y. B., and Joe, Y. A. (1997). Hypusine is essential for eukaryotic cell proliferation. *Biol. Signals* 6, 115–123. doi: 10.1159/000109117
- Patel, P. H., Costa-Mattioli, M., Schulze, K. L., and Bellen, H. J. (2009). The *Drosophila* deoxyhypusine hydroxylase homologue nero and its target eIF5A are required for cell growth and the regulation of autophagy. *J. Cell Biol.* 185, 1181–1194. doi: 10.1083/jcb.200904161
- Pathak, E., Atri, N., and Mishra, R. (2014). Analysis of P-loop and its flanking region subsequence of diverse NTPases reveals evolutionary selected residues. *Bioinformatics* 10, 216–220. doi: 10.6026/97320630010216
- Paul, A. L., Folta, K. M., and Ferl, R. J. (2008). 14-3-3 proteins, red light and photoperiodic flowering: a point of connection? *Plant Signal. Behav.* 3, 511–515. doi: 10.4161/psb.3.8.5717
- Paul, A. L., Sehnke, P. C., and Ferl, R. J. (2005). Isoform-specific subcellular localization among 14-3-3 proteins in *Arabidopsis* seems to be driven by client interactions. *Mol. Biol. Cell* 16, 1735–1743. doi: 10.1091/mbc.E04-09-0839
- Peer, W. A., Brown, D. E., Tague, B. W., Muday, G. K., Taiz, L., and Murphy, A. S. (2001). Flavonoid accumulation patterns of transparent testa mutants of *Arabidopsis*. *Plant Physiol.* 126, 536–548. doi: 10.1104/pp.126.2.536
- Pnueli, L., Gutfinger, T., Hareven, D., Ben-Naim, O., Ron, N., Adir, N., et al. (2001). Tomato SP-interacting proteins define a conserved signaling system that regulates shoot architecture and flowering. *Plant Cell* 13, 2687–2702. doi: 10.2307/3871528
- Purwestri, Y. A., Ogaki, Y., Tamaki, S., Tsuji, H., and Shimamoto, K. (2009). The 14-3-3 protein GF14c acts as a negative regulator of flowering in rice by interacting with the florigen Hd3a. *Plant Cell Physiol.* 50, 429–438. doi: 10.1093/pcp/pcp012
- Putnam-Evans, C., Harmon, A. C., Palevitz, B. A., Fechheimer, M., and Cormier, M. J. (1989). Calcium-dependent protein kinase is localized with F-actin in plant cells. *Cell Motil. Cytoskeleton* 12, 12–22. doi: 10.1002/cm.970120103
- Ramakrishnan, C., Dani, V. S., and Ramasarma, T. (2002). A conformational analysis of Walker motif A [GXXXXGKT (S)] in nucleotide-binding and other proteins. *Protein Eng.* 15, 783–798. doi: 10.1093/protein/15.10.783
- Ransom-Hodgkins, W. D. (2009). The application of expression analysis in elucidating the eukaryotic elongation factor one alpha gene family in *Arabidopsis thaliana*. *Mol. Genet. Genomics* 281, 391–405. doi: 10.1007/s00438-008-0418-2
- Ren, B., Chen, Q., Hong, S., Zhao, W., Feng, J., Feng, H., et al. (2013). The *Arabidopsis* eukaryotic translation initiation factor eIF5A-2 regulates root protoxylem development by modulating cytokinin signaling. *Plant Cell* 25, 3841–3857. doi: 10.1105/tpc.113.11.6236
- Roberts, D., Pedmale, U. V., Morrow, J., Sachdev, S., Lechner, E., Tang, X., et al. (2011). Modulation of phototropic responsiveness in *Arabidopsis* through ubiquitination of phototropin 1 by the CUL3-Ring E3 ubiquitin ligase CRL3(NPH3). *Plant Cell* 23, 3627–3640. doi: 10.1105/tpc.111.087999
- Rösler, J., Klein, I., and Zeidler, M. (2007). *Arabidopsis* fhl/fhy1 double mutant reveals a distinct cytoplasmic action of phytochrome A. *Proc. Natl. Acad. Sci. U.S.A.* 104, 10737–10742. doi: 10.1073/pnas.0703855104
- Rouleau, M., Marsolais, F., Richard, M., Nicolle, L., Voigt, B., Adam, G., et al. (1999). Inactivation of brassinosteroid biological activity by a salicylate-inducible steroid sulfotransferase from *Brassica napus*. *J. Biol. Chem.* 274, 20925–20930. doi: 10.1074/jbc.274.30.20925
- Rutschmann, F., Stalder, U., Piotrowski, M., Oecking, C., and Schaller, A. (2002). LeCPK1, a calcium-dependent protein kinase from tomato. Plasma membrane targeting and biochemical characterization. *Plant Physiol.* 129, 156–168. doi: 10.1104/pp.000869
- Saini, P., Eyler, D. E., Green, R., and Dever, T. E. (2009). Hypusine-containing protein eIF5A promotes translation elongation. *Nature* 459, 118–121. doi: 10.1038/nature08034
- Saraste, M., Sibbald, P. R., and Wittinghofer, A. (1990). The P-loop—a common motif in ATP- and GTP-binding proteins. *Trends Biochem. Sci.* 15, 430–434. doi: 10.1016/0968-0004(90)90281-F
- Saurin, A. J., Borden, K. L., Boddy, M. N., and Freemont, P. S. (1996). Does this have a familiar RING? *Trends Biochem. Sci.* 21, 208–214. doi: 10.1016/S0968-0004(96)80017-X
- Sawa, M., Nusinow, D. A., Kay, S. A., and Imaizumi, T. (2007). FKF1 and GIGANTEA complex formation is required for day-length measurement in *Arabidopsis*. *Science* 318, 261–265. doi: 10.1126/science.1146994
- Schaefer, D. G., and Zryd, J.-P. (1997). Efficient gene targeting in the moss *Physcomitrella patens*. *Plant J.* 11, 1195–1206. doi: 10.1046/j.1365-313X.1997.11061195.x
- Schmidt von Braun, S. (2008). *Chup1 – A Chloroplast Movement Protein and its Interactions*. Ph.D. dissertation, Ludwig Maximilian University of Munich, Munich.
- Schmidt von Braun, S., and Schleiff, E. (2008). The chloroplast outer membrane protein CHUP1 interacts with actin and profilin. *Planta* 227, 1151–1159. doi: 10.1007/s00425-007-0688-7
- Schnepf, E. (1986). Cellular polarity. *Ann. Rev. Plant Physiol.* 37, 23–47. doi: 10.1146/annurev.pp.37.060186.000323
- Schumann, N., Navarro-Quezada, A., Ullrich, K., Kuhl, C., and Quint, M. (2011). Molecular evolution and selection patterns of plant F-box proteins with C-terminal kelch repeats. *Plant Physiol.* 155, 835–850. doi: 10.1104/pp.110.166579
- Seo, P. J., Kim, S. G., and Park, C. M. (2008). Membrane-bound transcription factors in plants. *Trends Plant Sci.* 13, 550–556. doi: 10.1016/j.tplants.2008.06.008
- Shi, X. P., Yin, K. C., Zimolo, Z. A., Stern, A. M., and Waxman, L. (1996). The subcellular distribution of eukaryotic translation initiation factor, eIF-5A, in cultured cells. *Exp. Cell Res.* 225, 348–356. doi: 10.1006/excr.1996.0185
- Shiina, N., Gotoh, Y., Kubomura, N., Iwamatsu, A., and Nishida, E. (1994). Microtubule severing by elongation factor 1 alpha. *Science* 266, 282–285. doi: 10.1126/science.7939665
- Shin, R., Jez, J. M., Basra, A., Zhang, B., and Schachtman, D. P. (2011). 14-3-3 proteins fine-tune plant nutrient metabolism. *FEBS Lett.* 585, 143–147. doi: 10.1016/j.febslet.2010.11.025
- Slobin, L. I. (1980). The role of eucaryotic factor Tu in protein synthesis. The measurement of the elongation factor Tu content of rabbit reticulocytes and other mammalian cells by a sensitive radioimmunoassay. *Eur. J. Biochem.* 110, 555–563. doi: 10.1111/j.1432-1033.1980.tb04898.x
- Stellberger, T., Hauser, R., Baiker, A., Pothineni, V. R., Haas, J., and Uetz, P. (2010). Improving the yeast two-hybrid system with permutated fusions proteins: the Varicella Zoster Virus interactome. *Proteome Sci.* 8, 8. doi: 10.1186/1477-5956-8-8
- Sullivan, S., Thomson, C. E., Kaiserli, E., and Christie, J. M. (2009). Interaction specificity of *Arabidopsis* 14-3-3 proteins with phototropin receptor kinases. *FEBS Lett.* 583, 2187–2193. doi: 10.1016/j.febslet.2009.06.011
- Szekeres, M., Nemeth, K., Koncz-Kalman, Z., Mathur, J., Kauschmann, A., Altmann, T., et al. (1996). Brassinosteroids rescue the deficiency of CYP90, a cytochrome P450, controlling cell elongation and de-etiolation in *Arabidopsis*. *Cell* 85, 171–182. doi: 10.1016/S0092-8674(00)81094-6
- Takac, T., Pechan, T., Richter, H., Muller, J., Eck, C., Bohm, N., et al. (2011). Proteomics on brefeldin A-treated *Arabidopsis* roots reveals profilin 2 as a new protein involved in the cross-talk between vesicular trafficking and the actin cytoskeleton. *J. Proteome Res.* 10, 488–501. doi: 10.1021/pr100690f
- Takagi, S., Kong, S. G., Mineyuki, Y., and Furuya, M. (2003). Regulation of actin-dependent cytoplasmic motility by type II phytochrome occurs within seconds in *Vallisneria gigantea* epidermal cells. *Plant Cell* 15, 331–345. doi: 10.1105/tpc.007237
- Takase, T., Nishiyama, Y., Tanihigashi, H., Ogura, Y., Miyazaki, Y., Yamada, Y., et al. (2011). LOV KELCH PROTEIN2 and ZEITLUPE repress *Arabidopsis*

- photoperiodic flowering under non-inductive conditions, dependent on FLAVIN-BINDING KELCH REPEAT F-BOX1. *Plant J.* 67, 608–621. doi: 10.1111/j.1365-3113X.2011.04618.x
- Takekoshi, N., Higashitsuji, Y., Konzack, S., and Fischer, R. (2008). Apical sterol-rich membranes are essential for localizing cell end markers that determine growth directionality in the filamentous fungus *Aspergillus nidulans*. *Mol. Biol. Cell* 19, 339–351. doi: 10.1091/mbc.E07-06-0523
- Tanada, T. (1968). A rapid photoreversible response of barley root tips in the presence of 3-indoleacetic acid. *Proc. Natl. Acad. Sci. U.S.A.* 59, 376–380. doi: 10.1073/pnas.59.2.376
- Thelander, M., Nilsson, A., Olsson, T., Johansson, M., Girod, P. A., Schaefer, D. G., et al. (2007). The moss genes PpSKI1 and PpSKI2 encode nuclear SnRK1 interacting proteins with homologues in vascular plants. *Plant Mol. Biol.* 64, 559–573. doi: 10.1007/s11103-007-9176-5
- Thelander, M., Olsson, T., and Ronne, H. (2004). Snf1-related protein kinase 1 is needed for growth in a normal day-night light cycle. *EMBO J.* 23, 1900–1910. doi: 10.1038/sj.emboj.7600182
- Thompson, E. P., Smith, S. G., and Glover, B. J. (2012). An *Arabidopsis* rhomboid protease has roles in the chloroplast and in flower development. *J. Exp. Bot.* 63, 3559–3570. doi: 10.1093/jxb/ers012
- Thompson, J. E., Hopkins, M. T., Taylor, C., and Wang, T. W. (2004). Regulation of senescence by eukaryotic translation initiation factor 5A: implications for plant growth and development. *Trends Plant Sci.* 9, 174–179. doi: 10.1016/j.tplants.2004.02.008
- Tong, Y., Park, I., Hong, B. S., Nedyalkova, L., Tempel, W., and Park, H. W. (2009). Crystal structure of human eIF5A1: insight into functional similarity of human eIF5A1 and eIF5A2. *Proteins* 75, 1040–1045. doi: 10.1002/prot.22378
- Torrelío, B. M., Paz, M. A., and Gallop, P. M. (1984). Cellular proliferation and hypusine synthesis. *Exp. Cell Res.* 154, 454–463. doi: 10.1016/0014-4827(84)90169-1
- Tseng, T. S., Whippo, C., Hangarter, R. P., and Briggs, W. R. (2012). The role of a 14-3-3 protein in stomatal opening mediated by PHOT2 in *Arabidopsis*. *Plant Cell* 24, 1114–1126. doi: 10.1105/tpc.111.092130
- Ueno, K., Kinoshita, T., Inoue, S., Emi, T., and Shimazaki, K. (2005). Biochemical characterization of plasma membrane H⁺-ATPase activation in guard cell protoplasts of *Arabidopsis thaliana* in response to blue light. *Plant Cell Physiol.* 46, 955–963. doi: 10.1093/pcp/pci104
- Uetz, P., and Grigoriev, A. (2005). “The yeast interactome,” in *Encyclopedia of Genetics, Genomics, Proteomics and Bioinformatics*, Vol. 5, eds L. B. Jorde, P. F. R. Little, M. J. Dunn, and S. Subramaniam (Chichester: John Wiley & Sons Ltd), 2033–2051.
- Urano, D., Chen, J.-G., Botella, J. R., and Jones, A. M. (2013). Heterotrimeric G protein signalling in the plant kingdom. *Open Biol.* 3, 120186. doi: 10.1098/rsob.120186
- Urban, S., Lee, J. R., and Freeman, M. (2001). *Drosophila* rhomboid-1 defines a family of putative intramembrane serine proteases. *Cell* 107, 173–182. doi: 10.1016/S0092-8674(01)00525-6
- Urban, S., and Wolfe, M. S. (2005). Reconstitution of intramembrane proteolysis in vitro reveals that pure rhomboid is sufficient for catalysis and specificity. *Proc. Natl. Acad. Sci. U.S.A.* 102, 1883–1888. doi: 10.1073/pnas.0408306102
- Varin, L., Deluca, V., Ibrahim, R. K., and Brisson, N. (1992). Molecular characterization of two plant flavonol sulfotransferases. *Proc. Natl. Acad. Sci. U.S.A.* 89, 1286–1290. doi: 10.1073/pnas.89.4.1286
- Vetter, I. R., and Wittinghofer, A. (1999). Nucleoside triphosphate-binding proteins: different scaffolds to achieve phosphoryl transfer. *Q. Rev. Biophys.* 32, 1–56. doi: 10.1017/S0033583599003480
- Vidal, D., Gil, M. T., Alvarez-Florez, F., Moysset, L., and Simon, E. (2007). Protein kinase activity in *Cucumis sativus* cotyledons: effect of calcium and light. *Phytochemistry* 68, 438–445. doi: 10.1016/j.phytochem.2006.11.001
- Vidal, L., Augustine, R. C., Kleinman, K. P., and Bezanilla, M. (2007). Profilin is essential for tip growth in the moss *Physcomitrella patens*. *Plant Cell* 19, 3705–3722. doi: 10.1105/tpc.107.053413
- von Schwartzenberg, K. (2006). Moss biology and phytohormones – cytokinins in *Physcomitrella*. *Plant Biol. (Stuttg.)* 8, 382–388. doi: 10.1055/s-2006-923962
- Wagner, J. R., Brunzelle, J. S., Forest, K. T., and Vierstra, R. D. (2005). A light-sensing knot revealed by the structure of the chromophore-binding domain of phytochrome. *Nature* 438, 325–331. doi: 10.1038/nature04118
- Walker, J. E., Saraste, M., Runswick, M. J., and Gay, N. J. (1982). Distantly related sequences in the alpha- and beta-subunits of ATP synthase, myosin, kinases and other ATP-requiring enzymes and a common nucleotide binding fold. *EMBO J.* 1, 945–951.
- Wang, L., Xu, C., Wang, C., and Wang, Y. (2012). Characterization of a eukaryotic translation initiation factor 5A homolog from *Tamarix androssowii* involved in plant abiotic stress tolerance. *BMC Plant Biol.* 12:118. doi: 10.1186/1471-2229-12-118
- Wang, T. W., Lu, L., Wang, D., and Thompson, J. E. (2001). Isolation and characterization of senescence-induced cDNAs encoding deoxyhypusine synthase and eukaryotic translation initiation factor 5A from tomato. *J. Biol. Chem.* 276, 17541–17549. doi: 10.1074/jbc.M008544200
- Warpeha, K. M., Upadhyay, S., Yeh, J., Adamiak, J., Hawkins, S. I., Lapik, Y. R., et al. (2007). The GCR1, GPA1, PRN1, NF-Y signal chain mediates both blue light and abscisic acid responses in *Arabidopsis*. *Plant Physiol.* 143, 1590–1600. doi: 10.1104/pp.106.089904
- Wei, N., Chamovitz, D. A., and Deng, X. W. (1994a). *Arabidopsis* COP9 is a component of a novel signaling complex mediating light control of development [published erratum appears in *Cell* 1994 Oct 7;79(1):following 179]. *Cell* 78, 117–124. doi: 10.1016/0092-8674(94)90578-9
- Wei, N., Kwok, S. F., Von Arnim, A. G., Lee, A., Mcnellis, T. W., Piekos, B., et al. (1994b). *Arabidopsis* COP8, COP10, and COP11 genes are involved in repression of photomorphogenic development in darkness. *Plant Cell* 6, 629–643. doi: 10.2307/3869868
- Wei, N., Tsuge, T., Serino, G., Dohmae, N., Takio, K., Matsui, M., et al. (1998). The COP9 complex is conserved between plants and mammals and is related to the 26S proteasome regulatory complex. *Curr. Biol.* 8, 919–922. doi: 10.1016/S0960-9822(07)00372-7
- Weir, B. A., and Yaffe, M. P. (2004). Mmd1p, a novel, conserved protein essential for normal mitochondrial morphology and distribution in the fission yeast *Schizosaccharomyces pombe*. *Mol. Biol. Cell* 15, 1656–1665. doi: 10.1091/mbc.E03-06-0371
- Wheeler, G. L., and Brownlee, C. (2008). Ca²⁺ signalling in plants and green algae-changing channels. *Trends Plant Sci.* 13, 506–514. doi: 10.1016/j.tplants.2008.06.004
- Xia, G., Ramachandran, S., Hong, Y., Chan, Y. S., Simanis, V., and Chua, N. H. (1996). Identification of plant cytoskeletal, cell cycle-related and polarity-related proteins using *Schizosaccharomyces pombe*. *Plant J.* 10, 761–769. doi: 10.1046/j.1365-3113X.1996.10040761.x
- Yaffe, M. B. (2002). How do 14-3-3 proteins work?– Gatekeeper phosphorylation and the molecular anvil hypothesis. *FEBS Lett.* 513, 53–57. doi: 10.1016/S0014-5793(01)03288-4
- Yamashino, T., Kitayama, M., and Mizuno, T. (2013). Transcription of ST2A encoding a sulfotransferase family protein that is involved in jasmonic acid metabolism is controlled according to the circadian clock- and PIF4/PIF5-mediated external coincidence mechanism in *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* 77, 2454–2460. doi: 10.1271/bbb.130559
- Yang, W., and Boss, W. F. (1994). Regulation of phosphatidylinositol 4-kinase by the protein activator PIK-A49. Activation requires phosphorylation of PIK-A49. *J. Biol. Chem.* 269, 3852–3857.
- Yang, W., Burkhart, W., Cavallius, J., Merrick, W. C., and Boss, W. F. (1993). Purification and characterization of a phosphatidylinositol 4-kinase activator in carrot cells. *J. Biol. Chem.* 268, 392–398.
- Yasui, Y., Mukougawa, K., Uemoto, M., Yokofuji, A., Suzuri, R., Nishitani, A., et al. (2012). The phytochrome-interacting vascular plant one-zinc finger1 and VOZ2 redundantly regulate flowering in *Arabidopsis*. *Plant Cell* 24, 3248–3263. doi: 10.1105/tpc.112.101915
- Zanelli, C. F., and Valentini, S. R. (2005). Pkc1 acts through Zds1 and Gic1 to suppress growth and cell polarity defects of a yeast eIF5A mutant. *Genetics* 171, 1571–1581. doi: 10.1534/genetics.105.048082
- Zhang, B., Tremousaygue, D., Denance, N., Van Esse, H. P., Horger, A. C., Dabos, P., et al. (2014). PIRIN2 stabilizes cysteine protease XCP2 and increases susceptibility to the vascular pathogen *Ralstonia solanacearum* in *Arabidopsis*. *Plant J.* 79, 1009–1019. doi: 10.1111/tpj.12602
- Zhou, W., Wei, L., Xu, J., Zhai, Q., Jiang, H., Chen, R., et al. (2010). *Arabidopsis* Tyrosylprotein sulfotransferase acts in the auxin/PLETHORA pathway in

- regulating postembryonic maintenance of the root stem cell niche. *Plant Cell* 22, 3692–3709. doi: 10.1105/tpc.110.075721
- Zhu, S. Y., Yu, X. C., Wang, X. J., Zhao, R., Li, Y., Fan, R. C., et al. (2007). Two calcium-dependent protein kinases, CPK4 and CPK11, regulate abscisic acid signal transduction in *Arabidopsis*. *Plant Cell* 19, 3019–3036. doi: 10.1105/tpc.107.050666
- Zou, J. J., Wei, F. J., Wang, C., Wu, J. J., Ratnasekera, D., Liu, W. X., et al. (2010). *Arabidopsis* calcium-dependent protein kinase CPK10 functions in abscisic acid- and Ca²⁺-mediated stomatal regulation in response to drought stress. *Plant Physiol.* 154, 1232–1243. doi: 10.1104/pp.110.157545

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Ermert, Mailliet and Hughes. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.