Parasitology Research (2026) 125:41
https://doi.org/10.1007/s00436-026-08658-6

RESEARCH

Endoparasites of the Imperial Cormorant (*Leucocarbo atriceps*) in the Falkland Islands: investigation of secondary hosts and detection methods

Hannah Métaireau[1] · **Carlos Hermosilla**[2] · **Petra Quillfeldt**[1]

Received: 15 September 2025 / Accepted: 25 February 2026
© The Author(s) 2026

Abstract
Seabirds are important apex predators and valuable sentinels of marine ecosystem health. Because they integrate signals from multiple trophic levels, their diet provides important insights into marine food-web dynamics, gastrointestinal parasite fauna, and the impacts of human activities. Changes in diet composition can affect seabird reproductive success, health status and parasite load. Many protozoan and helminth parasites depend on predator-prey interactions to complete their life cycles, making dietary analysis a useful tool for understanding parasite transmission and ecology. Here, 82 pellets were collected from an Imperial Cormorant (*Leucocarbo atriceps*) colony in the Falkland Islands/Malvinas Islands (51°43'S, 61°19'W) in 2020 and 2022. Preys were identified using hard remains and metazoan endoparasites using morphological and genetic identification. Parasitological analysis of pellets was performed using the standard sodium-acetate-formaldehyde method (SAF). The diet primarily consisted of fish, particularly ray-thinned fish species (*Patagonotothen* spp.), crustaceans [squat lobsters; (*Munida* spp.)], and to a lesser extent polychaetes and cephalopods. The presence of nematodes of the family Anisakidae, including the genus *Contracaecum*, was detected using both methods (PCR and SAF). The comparison with the diet suggests that ray-thinned fish *(Patagonotothen* spp.) likely act as paratenic hosts for *Contracaecum* (Anisakidae) infections in the Imperial Cormorant. Further, a fish-borne trematode (Heterophyidae) was detected by SAF, which is the first reported in free-ranging Imperial Cormorants. However, no relationship could be demonstrated between this parasite and prey. Further studies are necessary to fill the gaps in our knowledge of parasites' life cycles and their effects on Imperial Cormorant's health in the Falklands.

Keywords Parasite-host associations · Anisakid nematodes · Fish-borne trematodes · Seabird parasites · Non-invasive sampling · Molecular analysis

Introduction

Knowledge of helminth associations with their hosts is crucial in understanding their complex life cycles, particularly in the marine ecosystem. However, detecting endoparasites in each host and their role during their life cycle is challenging (Garbin et al. 2019a). Several studies have been conducted to establish the life cycle of *Contracaecum* species experimentally (Huizinga 1966; Semenova 1979; Køie and Fagerholm 1995). Nevertheless, studying the life cycle in situ is particularly difficult due to its complexity, involving paratenic and intermediate hosts and the significant morphological similarity between the larval stages of the different species. Methods such as cadaver examination or pellet analysis are valuable for addressing such questions, as they enable the detection of adults, larval stages and/or prey items (Garbin et al. 2008, 2019a, b). Endoparasites, including protozoan and metazoan parasites, have also been detected as cysts and eggs using the standard sodium-acetate-formaldehyde (SAF) technique (Kleinertz et al. 2014).

Section Editor: Guillermo Salgado-Maldonado

✉ Hannah Métaireau
hmetaireau@icloud.com

[1] Institute of Zoology, Justus Liebig University Giessen, Heinrich-Buff-Ring 26, Giessen D-35392, Germany

[2] Institute of Parasitology, Biomedical Research Centre Seltersberg (BFS), Justus Liebig University Giessen, Giessen, Germany

Published online: 11 March 2026

This method is commonly employed for scat sample analysis. In this study, we aimed to apply the SAF method to the remains of the pellets to determine whether this type of sample was compatible with this method for detecting endoparasites. Furthermore, we sought to investigate whether this approach could enhance the detection results of the conventional use of pellets for identifying larvae and eggs.

The Imperial Cormorant or Imperial Shag (*Leucocarbo atriceps*), known locally as the King Shag (order Pelecaniformes, family Phalacrocoracidae) is endemically distributed in South America, from central Chile and Argentina to their coasts and the Falklands Islands (Johnsgard 1993). Imperial Cormorants consume various marine prey, primarily feeding from the benthic zone. However, it has been shown that while females forage in coastal waters, males move into pelagic waters to feed on larger prey (Quillfeldt et al. 2011). Three groups can be identified as the primary food sources: fish, crustaceans, and cephalopods, with fish being the most important prey (Thompson 1989; Punta et al. 1993, 2003; Malacalza et al. 1994; Gosztonyi and Kuba 1998; Ferrari et al. 2004; Bulgarella et al. 2008; Michalik et al. 2010; Yorio et al. 2010; Ibarra et al. 2018; Morgenthaler et al. 2022). They are prone to endoparasites as their diet is mainly composed of fish. The Natural History Museum Host-Parasite database (Gibson et al. 2022) records helminths found parasitising Imperial Cormorants, and various studies have documented other helminths in different regions of South America (Gutiérrez 1943; Torres et al. 1990; Malacalza et al. 1998; Garbin et al. 2008, 2013, 2019a; Garbin 2009; Diaz et al. 2009; Oyarzún-Ruiz and Muñoz-Alvarado 2015). However, to our knowledge, no study has specifically investigated the parasites of the Imperial Cormorant in the Falkland Islands.

Our work aimed to assess the diet and associated endoparasites of the Imperial Cormorant colony on New Island in the Falkland Islands. This assessment was conducted by analysing pellets [regurgitated casts, primarily composed of hard food items (Michalik et al. 2010)]. Specifically, we aimed to characterise the prey and endoparasites of the Imperial Cormorant, assess whether the SAF technique can be applied to pellet remains and determine parasite-prey associations.

Materials and methods

Study area

The study was conducted at New Island, Falkland/Malvinas Islands (51°43'S, 61°19'W) in the South Atlantic, east of southern Argentina (Fig. 1). The Falkland Islands comprise 778 islands and islets, with two major islands, West and East Falkland, covering 12,200 km^2 (Woods 2001). New Island is located in the southwest of the Falkland Islands and has been managed by Falklands Conservation since 2020. The island hosts 39 breeding bird species, accounting for 65% of the total breeding species in the Falkland Islands, and most are seabirds. Several endangered species breed on New Island, such as the Magallanic Penguin (*Spheniscus magellanicus*), Gentoo (*Pygoscelis papua*) and Rockhopper Penguins (*Eudyptes chrysocome*) and Black-browed Albatross (*Thalassarche melanophris*), which all share their rookeries with the Imperial Cormorant (Falklands Conservation 2023). The Falkland Island ecosystem is highly productive due to nutrient upwelling from the Falkland Current and the Antarctic Circumpolar Current (Agnew 2002; Van Der Grient et al. 2023), making this area attractive to many species, including seabirds. On New Island, approximately 3,000 breeding pairs of Imperial Cormorants are present in the colony. They arrive at the colony in early October for courtship and nest-building (Michalik et al. 2010). They lay eggs between early November and late December, and nestlings are present from December to March (Quillfeldt et al. 2011).

Pellet sampling

Samples were collected during the austral summer (January-February) of two breeding seasons, 2020 and 2022. The collection involved slowly walking around the colony and searching for intact pellets without major disturbance of seabirds. When a pellet was found, a protective board was used to collect pellets close to the birds, preventing bites safely. The pellets were placed in labelled tubes and stored at −20 °C until examination. A total of 82 pellets were collected, with 42 from 2020 to 40 from 2022.

Diet analysis

Once in the laboratory, the pellets were removed from the freezer and left to thaw for dissection. Each pellet was placed in a Petri dish and dissected under a binocular magnifying glass. Hard remnants, such as otoliths, squid beaks, fish bones, and polychaete jaws were removed, sorted, and placed inside tubes. The remaining material from the pellets was returned to the containers with ethanol (70%) for preservation and further analysis. The identification of the hard prey remnants was carried out under the stereomicroscope with reference guides. Otoliths were sorted by size and shape and then paired when possible. For pellets containing only fish bones, they accounted for the presence of fish. For the identification of the otoliths, the Otoliths Reference Collection available on the Global Biodiversity Information Facility (gbif.org) was used as well as some identification books (Hecht 1987; Reid 1996; Volpedo and



Fig. 1 Map of the Falkland Islands. The study colony is shown by a red cross

Echeverría 2000) and help from experts (cf. Acknowledgments). Cephalopod beaks were identified using a reference guide (Xavier and Cherel 2021) and the help of experts (cf. Acknowledgments). Crustaceans were identified using an identification guide (Xavier et al. 2020). Other prey items (e.g. polychaete jaws, gastropods, and bivalves) found were identified with the help of experts (cf. Acknowledgments).

Parasite analysis

Morphological methods

Parasite larvae found inside the pellets, during the dissection, were put inside a labelled tube with ethanol (70%) to preserve them for identification. Pictures of the larvae, the anterior part, and the tail were taken, and the length of all individuals was measured using the stereomicroscope (Leica MZ 7.5). Photos were taken using an Olympus SC30 camera and the software cellSens Dimension, and can be found in the Supplementary file. Identification of the larvae was done using identification guides and articles (Anderson 2000; Garbin 2009; Garbin et al. 2013, 2019b, 2023; González-Acuña et al. 2020). Because identification based on morphology cannot provide precise information at the species level (Bhadury et al. 2006), we decided to use DNA barcoding for final species identification.

DNA amplification and sequencing

Larvae preserved in 70% ethanol were first dried to remove residual ethanol prior to DNA extraction. When multiple small larvae were recovered from a single pellet, they were pooled for a single extraction to obtain sufficient DNA for sequencing. All other samples contained a single larva, resulting in a total of 46 extraction samples. Molecular analyses were restricted to nematodes, and no protozoan-specific PCR assays were performed.

DNA extraction from the larvae was then performed using the QIAwave DNA Blood and Tissue Kit according to the manufacturer's instructions. The extracted DNA was stored at −18 °C, and Nanodrop measurements (NanoDrop 2000, Thermo Scientific) were used to determine DNA concentration (optimal DNA concentration: 20ng/µl). A negative control was used in all PCR tests.

The mitochondrial cytochrome c oxidase subunit I (COI) region was amplified using the primer pair JB3/JB4.5 (Bowles et al. 1992). The ITS1 and ITS2 regions were amplified using the primer pairs SS1/NC13R and SS2/NC2,

respectively (Shamsi et al. 2008). These primers are widely used for molecular identification of anisakid/ascaridoid nematodes. For the JB3/JB4.5 primers, PCR was performed in 25 µl containing 12.5 µl of Dream Taq Buffer (DreamTaq DNA Polymerase (ThermoScientific), 5.5 µl of water, 2.5 µl of each primer and 2 µl of DNA. The PCR conditions for the JB3/JB4.5 primer pair were as follows: initial denaturation (94 °C for 3 min), 40 cycles at 94 °C for 45 s (denaturation), 53 °C for 45 s (annealing), and 72 °C for 45 s (extension), and 72 °C for 10 min (final extension). For the SS1/NC13R and SS2/NC2 primers, the PCR mixture of 25 µl contained 12.5 µl of Dream Taq Buffer (DreamTaq DNA Polymerase (ThermoScientific), 9 µl of water, 1.25 µl of each primer and 2 µl of DNA. The PCR conditions were as follows: initial denaturation (94 °C for 5 min), 30 cycles at 94 °C for 30 s (denaturation), 55 °C for 30 s (annealing), and 72 °C for 30 s (extension), and 72 °C for 5 min (final extension). A QIAxcel Advanced (Qiagen, Switzerland) high-resolution capillary gel electrophoresis was used to visualize band heights. The PCR amplicons of positive samples were sent to Seqlab (Göttingen, Germany) for bi-directional Sanger sequencing after being purified. The results were read and assembled with CLC Main Workbench 7.6.4 (CLC Bio, Aarhus, Denmark). The resulting consensus sequences were checked, and any conflicting bases were assigned manually as far as possible. Using the NCBI Blast (GenBank), every sequence was checked for its closest match to determine the parasite genus.

SAF method

The remains of the pellets were then taken to the Institute of Parasitology of the Justus Liebig University for examination of intestinal parasites. The SAF technique was used to detect parasitic stages of helminths. Under a fume hood, a small quantity of the sample was taken (bean size) and placed in a tube containing 10 ml of the SAF solution [sodium acetate: 15 g, glacial acetic acid: 20 ml, formaldehyde (37%): 40 ml, water: 925 ml]. The sample was mixed with the liquid to make it homogeneous and then filtered over a funnel containing double-layer gauze. The tubes were centrifuged at 2000 rpm for 3 min at room temperature (RT). The supernatant was then drained. A sodium chloride solution (7 ml) was added, and the deposit was mixed using a Pasteurizer pipette. Acid ethyl ester (2.5–3 ml) was added, and the tube was shaken using a stopper. The tubes were centrifuged again at 2000 rpm for 3 min at RT, and the supernatant was drained. To identify the intestinal parasites, 2–3 drops of the sample were allocated on top of a microscope slide and examined under an Olympus BX43 light microscope (x20 – x40 magnification). Pictures were taken using an Olympus SC30 digital camera and the software cellSens Dimension, and can be found in the Supplementary file. The eggs were identified using images from (Pennycott 2016a, b).

Data analysis

The frequency of occurrence was calculated, expressed as a percentage, for the prey items and can be defined as the proportion of pellets containing a certain prey type per pellet. Prevalence expressed as percentage (i.e. the number of hosts infected with 1 or more individuals of a particular parasite species divided by the number of hosts examined for that parasite species) and mean intensity (i.e. the total number of parasites of a particular species found in a sample divided by the number of hosts infected with that parasite) were calculated for the parasites (Bush et al. 1997). Only larvae were taken into account for the mean intensity. Confidence intervals were calculated using the binom package (Dorai-Raj 2022).

To test if the detection of parasites was similar between the larvae found in pellets and the SAF, Cohen's kappa was calculated (Cohen 1960) with the DescTools package (Signorell 2024). The results were interpreted as follows: no agreement for values ≤ 0, none to slight for values between 0.01 and 0.20, fair for values between 0.21 and 0.40, moderate for values between 0.41 and 0.60, substantial for values between 0.61 and 0.80, and almost perfect agreement for values between 0.81 and 1.00. The Cohen's kappa is reported, along with its 95% CI. Chi-squared tests were used to compare the prevalence of parasites, and Kruskal-Wallis tests were performed to compare the intensities of the different parasites, followed by a Pairwise Wilcoxon Rank Sum Test. A correspondence analysis was performed for prey with an occurrence above 5% to visualise parasite-prey association. Chi-squared tests were performed using observed frequencies of prey and parasites to investigate associations. The level of significance was set at $\alpha = 0.05$ throughout this study.

Results

Analyses of Imperial Cormorant pellets revealed 14 prey items belonging to 6 different taxa: fish, cephalopod, crustacea, gastropod, bivalve, and polychaete (Table 1).

Based on the morphological analyses, isolated larvae corresponded to L3 according to Anderson et al. (1974). Pellet dissection detected larval stages, whereas the SAF method detected parasite eggs. Only *Contracaecum* spp. were detected with both methods, whereas the other parasites were detected only as larvae and trematode eggs (Heterophyidae) were detected only with SAF (Fig. 2). Considering all parasites, agreement between the two methods was

Table 1 Frequency of occurrence (FO%) of prey items found inside the pellets ($n=82$) of the Imperial Cormorant *Leucocarbo atriceps* from the Falkland Islands

		FO (%)
Fish		95.1
Notothenidae		
	Patagonotothen sp	57.3
	Eleginops maclovinus	1.2
Moridae		
	Salilota australis	2.4
Unidentified[a]		67.1
Cephalopoda		**12.2**
Gonatidae		
	Gonatus antarticus	1.2
Loligonidae		
	Doryteuthis gahi	4.9
Octopodidae		
	Enteroctopus megalocyathus	4.9
Sepiolidae		1.2
Crustacea		**59.8**
Galatheidae		
	Munida sp	59.8
Talitridae		1.2
Gastropoda		**7.3**
Buccinidae		
	Pareuthria atrata	3.7
	Pareuthria fuscata	2.4
Unidentified		3.7
Bivalvia		**2.4**
Veneridae		
	Ameghinomya antiqua	2.4
Unidentified		1.2
Polychaeta		**14.6**
Nereididae		11.0
Polynoidae		6.1
Unidentified		2.4

[a]Based on otoliths and fish bones

considered fair with a *k* value of 0.31 (0.09–0.53). For *Contracaecum*, the agreement was moderate, with a *k* value of 0.45 (0.23–0.67).

PCR amplification was successful for 43 out of 46 parasite larvae samples. The JB3/JB4.5 primers yielded a success rate of 82.6%, followed by SS2/NC2 (76.1%) and SS1/NC13R (71.4%). For three samples, only one sequence was obtained using the SS1/NC13R primers. The three samples that failed to amplify were identified as Anisakidae based on morphological examination, while all other samples were identified using genetic analyses. Sequence quality was sufficient for reliable taxonomic assignment in 45 out of 46 samples, with sequence similarity values above 90%, and one sample showing 89.8% similarity. No protozoan was detected using the SAF method, and only helminth larvae were genetically identified. Among the identified larvae, two nematode families were detected: Anisakidae and Raphidascarididae. For Anisakidae, species composition within each genus (based on successfully sequenced larvae) was as follows: Contracaecum (*C. chubutensis* and *C. osculatum*; 77.8% and 3.7%, respectively), Anisakis (*A. simplex* and *A. pegreffii*; 50% and 25%, respectively), and Pseudoterranova (*P. decipiens* and *P. cattani*; 81.8% and 18.2%, respectively). The Raphidascarididae were represented by *Hysterothylacium aduncum* (100%), with nine larvae detected in a single sample. In addition, eggs belonging to the trematode family Heterophyidae were detected using the SAF method, but no genetic analyses were performed on these eggs.

Anisakidae and Raphidascarididae were more prevalent than Heterophyidae ($X^2= 21.8$, $df=1$, $p<0.001$). The genus *Contracaecum* had the highest prevalence, followed by *Anisakis*, *Pseudoterranova* and *Hysterothylacium* (Fig. 2). The difference in prevalence between the Anisakidae and Raphidascarididae parasites was significantly different ($X^2= 57.6$, $df=3$, $p<0.001$).

In terms of mean intensities, *Hysterothylacium* had the highest mean intensity, although a large range was attributed to only two samples with different intensities (Fig. 3). The range of intensities for *Pseudoterranova* was lower, with an average of 3 larvae per pellet for this parasite genus. *Contracaecum* and *Anisakis* intensities were the lowest, with most pellets containing only 1 larva of these parasites (Fig. 3). The difference in intensities was significant (Kruskal-Wallis - $X^2=9.9$, $df=3$, $p=0.019$). However, the only significant difference was between *Contracaecum* and *Pseudoterranova* (Pairwise Wilcoxon Rank Sum Test – $p=0.041$).

The Correspondence Analysis (CA) showed an association between three anisakid parasites (i.e. *Contracaecum*, *Anisakis* and *Pseudoterranova*), the trematode (Heterophyidae) and ray-thinned fish (*Patagonotothen* sp.; Fig. 4). *Hysterothylacium* was not associated with any prey. Only the association between *Contracaecum* and *Patagonotothen* sp. was significant ($X^2=5.9$, $df=1$, $p=0.015$).

Discussion

The Imperial Cormorant diet in the Falkland Islands consisted mostly of fish, followed by crustaceans, polychaetes, and cephalopods. This finding is consistent with other studies in this location and colonies along the Argentinian coast (Casaux et al. 1997; Punta et al. 2003; Bulgarella et al. 2008; Michalik et al. 2010; Yorio et al. 2010, 2017; Quillfeldt et al. 2011; Quintana et al. 2011; Harris et al. 2016; Ibarra et al. 2018; Morgenthaler et al. 2022). Among the fish prey, *Patagonotothen* sp. was found to be the most common, with some biases due to the exclusive use of otoliths




Fig. 2 Prevalence of the different parasites inside pellets depending on the detection method. The bars correspond to the 95% Confidence Intervals. Numbers correspond to the prevalence for each parasite depending on the detection method.

for fish identification (Jobling and Breiby 1986; Johnstone et al. 1990).

Endoparasites found in the Imperial Cormorant population on the Falkland Islands were primarily nematodes from the Anisakidae family. Previous research also documented anisakid nematodes in Imperial Cormorants at various locations (Gutiérrez 1943; Torres et al. 1990; Malacalza et al. 1998; Garbin et al. 2008, 2013, 2019a; Diaz et al. 2009; Oyarzún-Ruiz and Muñoz-Alvarado 2015), many of which belong to the genus *Contracaecum*. The three genera of Anisakidae found in our study (*Contracaecum*, *Anisakis*, *Pseudoterranova*) were also found in Imperial Cormorant at the Chubut province, and *Hysterothylacium* was found in Red-legged cormorant (*Poikilocarbo gaimardi*) in the Santa Cruz province (Garbin et al. 2019a).

This study found lower mean intensity compared to others. The mean intensity of infection by *Contracaecum* sp. was 7.1 and 6.6 in Punta Leon for the years 1991/1992 and 1992/1993, respectively (Malacalza et al. 1998). A more recent study at the same location found similar mean intensities for three Anisakidae genera, except for *Pseudoterranova*, which had the highest intensity (Garbin et al. 2019a). The differences in prevalence and intensity of parasites in our study compared to others may be attributed to disparities in location. In fact, Fonteneau and Cook (2013) reported different prevalences and intensities of *Contracaecum rudolphii* in Kerguelen Shags (*Phalacrocorax verrucosus*) breeding at two colonies 25 km apart. Our findings suggest that the diet of Imperial Cormorants at our location may be less diverse than in other locations, which may be associated with lower parasite diversity.

The SAF method has enabled us to detect fish-borne trematode eggs of the Heterophyidae family. This is the first record of this intestinal fluke family parasitising the Imperial Cormorant. However, the prevalence of this endoparasite is lower than for Anisakidae. This family is frequently found in fish-eating birds (Sepulveda et al. 1994, 1996, 1999). Santos and Borges (2020) review records of Heterophyidae in South America. In Argentina, records of this family showed a predominance of the genus *Ascocotyle*, followed by *Cryptocotyle* and *Pygidiopsis*. However, no cormorant in Argentina was found to be infected by Heterophyidae;




Fig. 3 Mean intensities of parasites found as larvae inside pellets. Each dot represents the number of parasite larvae found in one pellet

such infections were observed only in Brazil, Ecuador, Venezuela, and Chile (González-Acuña et al. 2020; Santos and Borges 2020). A more precise identification was impossible because Heterophyidae were only found as eggs. Other methods could be useful to gain better insight into the trematode parasites, such as nested PCR used by Hermosilla et al. (2018).

The two methods employed to evaluate the prevalence of parasites in the Imperial Cormorant produced divergent results in detecting the presence of endoparasites. Anisakids were the most prevalent parasites found with both methods. However, only larvae or eggs were found for some pellets, resulting in a lower agreement between the methods. Only *Contracaecum* from the Anisakidae family was found as larvae inside pellets and as eggs with the SAF technique. The other Anisakidae genera were not found by using the SAF method. This may reflect differences in host suitability among parasite genera. Heterophyid eggs were detected solely through the SAF method, with neither juvenile nor adult trematodes found in pellets. This result is surprising because we would have expected to find adult stages in the pellet when eggs are found.

Pellets are a non-invasive method that are simple to collect and have already been used in previous studies on parasites of the Imperial Cormorant (Malacalza et al. 1998; Garbin et al. 2019a). Moreover, dead Imperial Cormorants have also been examined (Garbin et al. 2008). The SAF method was applied for the first time, to our knowledge, with remnants of dissected pellets, which has been used for faecal samples originating from marine mammals and seabirds (Kleinertz et al. 2014; Ebmer et al. 2020; Fusaro et al. 2024). However, our results indicate that this method can also be used with pellets. Fusaro et al. (2024) compared three sampling techniques (eviscerated hosts, faecal samples, and regurgitates) to study endoparasites in the Southern Giant Petrel *(Macronectes giganteus*) in Antarctica. Viscera and faecal matter yielded superior results compared to regurgitates, but they are a valuable addition as it is challenging to access recently deceased specimens. Non-invasive techniques in parasitology have gained popularity in recent years (Hermosilla et al.




Fig. 4 Results of the correspondence analysis showing the parasite-prey associations. Anisk: *Anisakis sp*; Contr: *Contracaecum sp*; Heter: Heterophyidae; Hyst: *Hysterothylacium sp*; Pseudo: *Pseudoterranova sp*; Dor: *Doryteuthis gahi*; Enter: *Enteroctopus megalocyathus*; Mun: *Munida sp*; Nerei: *Nereididae sp*; Patag: *Patagonotothen sp*; Poly: *Polynoidae sp*

2018; Ebmer et al. 2020) compared to invasive methods like endoscopy, which necessitates a comprehensive evaluation of ethical considerations (Burthe et al. 2013). The absence of parasitic protozoa in the remaining pellets after SAF treatment may be due to the pellet material, which may not preserve protozoan cysts effectively. The SAF method could be used with scat samples from the Imperial Cormorant to confirm this hypothesis, or protozoan-specific PCR assays could be performed.

Among the parasites, the results showed a visual association between Rock cods and three parasite genera (*Contracaecum*, *Anisakis*, *Pseudoterranova*) and one family (Heterophyidae), but this association was significant only for *Contracaecum*. The lack of association between *Hysterothylacium and* other species is probably due to its low prevalence, which reduces the likelihood of association. In fact, studies carried out around the Falkland Islands on Rock cods have shown a wide diversity of parasites, including the nematodes found here (González et al. 2007; Brickle et al. 2021).

Among the Anisakidae, the genus *Contracaecum* is the only one maturing into adults in birds (Garbin et al. 2019a). In the case of this genus, several studies on its life cycle showed that copepods are the main intermediate host of *Contracaecum* sp., with euphausiids and amphipods playing a lesser role (Huizinga 1966; Bartlett 1996; Anderson 2000; Moravec 2009). These prey items were found in the diet of rock cods (Laptikhovsky 2004) and rock cods (*Patagonotothen ramsayi*) from around the Falkland Islands were found to have a high parasite diversity (González et al. 2007; Brickle et al.

2021). Together with our association results, this is consistent with rock cod likely acting as paratenic hosts for parasites infecting Imperial Cormorants. Studies on intermediate hosts around the Falkland Islands should be carried out to gain a better understanding of the life cycle of these parasites.

Even if the occurrence of fish-borne trematode (Heterophyidae) was visually associated with rock cods, it was not significant. Surveys have been conducted in the past around the Falkland Islands to assess parasites of several fish and squid species (MacKenzie and Longshaw 1995; Brickle et al. 2001, 2006; Brickle and MacKenzie 2007; González et al. 2007; Brown et al. 2013; MacKenzie et al. 2013). Further surveys are needed to better characterise trematode diversity in seabirds from this region. Although parasites belonging to the Trematoda class were found inside them, none of the species belonged so far to the Heterophyidae family. In the case of Heterophyidae, the trematode parasite is acquired through the consumption of infected fish where the metacercariae encyst in the organs (Yamaguti 1971, 1975). It is by eating a mollusc (gastropod) that the fish becomes infected. Species from this family were found as adults in seals and in penguins from San Matias Gulf, Argentina. Heterophyid trematodes were also found in Cormorant species such as the Neotropic Cormorant (*Phalacrocorax brasilianus*) in coastal lagoons of Mexico (Violante-González et al. 2011), the Flightless Cormorant (*Phalacrocorax harrisi*) in the Galapagos Islands (Carrera-Játiva et al. 2014), and the Double-Crested Cormorant (*Phalacrocorax auritus*) in the Mississippi Delta region (O'Hear et al. 2014). This represents the first record of this heterophyid trematode in the Imperial Cormorant. However, the identity of the intermediate hosts involved in its transmission in the Falkland Islands remains unresolved.

In addition to morphological identification, the molecular analyses performed in this study provided an important complementary line of evidence for species identification. It enabled the detection and taxonomic discrimination of anisakid taxa (Contracaecum, Anisakis, and Pseudoterranova) and the species *Hysterothylacium aduncum* (Raphidascarididae) from degraded pellet material, where morphological identification alone was insufficient. This taxonomic resolution allowed evaluation of parasite-prey associations, since accurate identification is necessary to link parasites with specific prey sources. The integration of molecular and morphological approaches, therefore, increases confidence in these associations and provides a more robust framework for reconstructing trophic and transmission pathways. However, the use of pellets may have biased estimates of both prey and parasite diversity towards taxa that are more resistant to digestion. Although molecular analyses improved taxonomic resolution, PCR amplification was unsuccessful for some samples, and trematode eggs were not genetically characterised, limiting species-level identification.

Supplementary Information The online version contains supplementary material available at https://doi.org/10.1007/s00436-026-08658-6.

Acknowledgements The authors would like to thank the experts who helped with the identification of prey items: P. Brickle, Y. Cherel and V. Laptikhovsky, B. Schöne, G. Pastorino, T. Darbyshire, L. Hughes.

Author contributions Conceptualization: H.M., P.Q.; Data collection: P.Q.; Analysis: H.M.; Writing-original draft: H.M.; Writing, review, editing: H.M., C.H., P.Q.

Funding Open Access funding enabled and organized by Projekt DEAL. No funding was received for conducting this study.

Data availability The dataset generated and analysed during the current study is available from the corresponding author on reasonable request.

Declarations

Ethics declaration Not applicable.

Competing interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

Agnew DJ (2002) Critical aspects of the Falkland Islands pelagic ecosystem: distribution, spawning and migration of pelagic animals in relation to oil exploration. Aquat Conserv 12:39–50. https://doi.org/10.1002/aqc.474

Anderson RC (2000) Nematode parasites of vertebrates: Their development and transmission, 2nd edn. CABI Publishing, Wallingford, Oxon, UK. https://doi.org/10.1079/9780851994215.0000

Anderson RC, Chabaud AG, Willmott S (1974) Keys to the nematode parasites of vertebrates. Commonwealth Agricultural Bureaux, Farnham Royal, Bucks, England

Bartlett CM (1996) Morphogenesis of *Contracaecum rudolphii* (Nematoda: Ascaridoidea), a parasite of fish-eating birds, in its copepod precursor and fish intermediate hosts. Parasite 3:367–376. https://doi.org/10.1051/parasite/1996034367

Bhadury P, Austen M, Bilton D, Lambshead P, Rogers A, Smerdon G (2006) Development and evaluation of a DNA-barcoding approach for the rapid identification of nematodes. Mar Ecol Prog Ser 320:1–9. https://doi.org/10.3354/meps320001

Bowles J, Blair D, McManus DP (1992) Genetic variants within the genus *Echinococcus* identified by mitochondrial DNA sequencing. Mol Biochem Parasitol 54:165–173. https://doi.org/10.1016/0166-6851(92)90109-W

Brickle P, MacKenzie K (2007) Parasites as biological tags for *Eleginops maclovinus* (Teleostei: Eleginopidae) around the Falkland Islands. J Helminthol 81:147–153. https://doi.org/10.1017/S0022149X07750514

Brickle P, Olson PD, Littlewood DTJ, Bishop A, Arkhipkin AI (2001) Parasites of *Loligo gahi* from waters off the Falkland Islands, with a phylogenetically based identification of their cestode larvae. Can J Zool 79:2289–2296. https://doi.org/10.1139/cjz-79-12-2289

Brickle P, MacKenzie K, Pike A (2006) Variations in the parasite fauna of the Patagonian Toothfish (*Dissostichus eleginoides* Smitt, 1898), with length, season and depth of habitat around the Falkland Islands. J Parasitol 92:282–291. https://doi.org/10.1645/GE-539R.1

Brickle P, Randhawa HS, Reid MR, Lee B, Shcherbich Z, Arkhipkin AI (2021) Otolith trace elemental analyses and parasites provide useful tools for the stock discrimination of *Patagonotothen ramsayi* (Regan, 1913) (Nototheniidae) on the southern Patagonian Shelf. Fish Res 244. https://doi.org/10.1016/j.fishres.2021.106129

Brown J, Brickle P, Scott BE (2013) The parasite fauna of the Patagonian toothfish *Dissostichus eleginoides* off the Falkland Islands. J Helminthol 87:501–509. https://doi.org/10.1017/S0022149X12000636

Bulgarella M, Pizarro LC, Quintana F, Gosztonyi A, Kuba L (2008) Diet of Imperial Cormorants (*Phalacrocorax atriceps*) and Rock Shags (*P. magellanicus*) breeding sympatrically in Patagonia, Argentina. Ornitol Neotrop 19:553–563

Burthe S, Newell MA, Goodman G, Butler A, Bregnballe T, Harris E, Wanless S, Cunningham EJA, Daunt F (2013) Endoscopy as a novel method for assessing endoparasite burdens in free-ranging European shags (*Phalacrocorax aristotelis*). Methods Ecol Evol 4:207–216. https://doi.org/10.1111/2041-210x.12015

Bush AO, Lafferty KD, Lotz JM, Shostak AW (1997) Parasitology meets ecology on its own terms: Margolis et al. revisited. J Parasitol 83(4):575. https://doi.org/10.2307/3284227

Carrera-Játiva PD, Rodríguez-Hidalgo R, Sevilla C, Jiménez-Uzcátegui G (2014) Gastrointestinal parasites in the Galapagos Penguin *Spheniscus mendiculus* and the flightless cormorant *Phalacrocorax harrisi* in the Galapagos Islands. Mar Ornithol 42:77–80. https://doi.org/10.5038/2074-1235.42.1.1066

Casaux RJ, Favero M, Coria N, Silva P (1997) Diet of the Imperial Cormorant *Phalacrocorax atriceps*: comparison of pellets and stomach contents. Mar Ornithol 25:1–4. https://doi.org/10.5038/2074-1235.25.1.373

Cohen J (1960) A coefficient of agreement for nominal scales. Educ Psychol Meas 20:37–46. https://doi.org/10.1177/001316446002000104

Diaz JI, Fonteneau F, Panisse G, Cremonte F, Navone GT (2009) Redescription of <Emphasis Type="Italic">Ingliseria cirrohamata</Emphasis> (Linstow, 1888) (Nematoda: Acuariidae), including new hosts and geographical records. J Parasitol 95:396–402. https://doi.org/10.1645/GE-1758.1

Dorai-Raj S (2022) Package binom. Version 1.1–1.1. Binomial confidence intervals for several parameterizations. CRAN

Ebmer D, Navarrete MJ, Muñoz P, Flores LM, Gärtner U, Brabec J, Poppert S, Taubert A, Hermosilla C (2020) Anthropozoonotic parasites circulating in synanthropic and pacific colonies of South American sea lions (*Otaria flavescens*): non-invasive techniques data and a review of the literature. Front Mar Sci 7. https://doi.org/10.3389/fmars.2020.543829

Falklands Conservation (2023) New Island Wildlife. https://www.newislandtrust.com/wildlife/. Accessed 14 March 2024

Ferrari S, Alegre B, Gandini P (2004) Dieta del cormoran imperial (*Phalacrocorax atriceps*) en el sur de Santa Cruz (Patagonia, Argentina). Ornitol Neotrop 15:103–110

Fonteneau F, Cook TR (2013) New data on gastrointestinal helminths in shags (*Phalacrocorax verrucosus*) at Kerguelen Archipelago. Polar Biol 36:1839–1843. https://doi.org/10.1007/s00300-013-1391-8

Fusaro B, Capasso S, Lorenti E, Panisse G, Fuentes L, Fornillo MV, Libertelli M, Ansaldo M, Diaz JI (2024) Assessing helminth infection in some reproductive colonies of Southern Giant Petrel (*Macronectes giganteus*) from Antarctica. Polar Biol 47:1065–1076. https://doi.org/10.1007/s00300-024-03281-6

Garbin L (2009) Taxonomia y evaluacion de la especificidad hospedatoria de nematodos Anisakidae parasitos de aves marinas en el aera de Peninsula Valdés, Chubut, Argentina. Universidad Nac de La Plata Facultad de Ciencias Naturales y Museo. https://doi.org/10.35537/10915/4323

Garbin L, Diaz JI, Cremonte F, Navone GT (2008) A new Anisakid species parasitizing the Imperial cormorant *Phalacrocorax atriceps* from the North Patagonian Coast, Argentina. J Parasitol 94:852–859. https://doi.org/10.1645/GE-1369.1

Garbin L, Mattiucci S, Paoletti M, Diaz JI, Nascetti G, Navone GT (2013) Molecular identification and larval morphological description of *Contracaecum pelagicum* (Nematoda: Anisakidae) from the anchovy *Engraulis anchoita* (Engraulidae) and fish-eating birds from the Argentine North Patagonian Sea. Parasitol Int 62:309–319. https://doi.org/10.1016/j.parint.2013.03.001

Garbin L, Diaz JI, Morgenthaler A, Millones A, Kuba L, Fuchs D, Navone GT (2019aa) Cormorant pellets as a tool for the knowledge of parasite-intermediate host associations and nematode diversity in the environment. Helminthologia 56:296–302. https://doi.org/10.2478/helm-2019a-0027

Garbin L, Diaz JI, Navone GT (2019) Species of Contracaecum parasitizing the Magellanic Penguin *Spheniscus magellanicus* (Spheniscidae) from the Argentinean Coast. J Parasitol 105(2):222. https://doi.org/10.1645/17-91

Garbin L, Diaz JI, Servián A, Fusaro B, Navone GT (2023) The genus Contracaecum Raillet and Henry (Nematoda: Anisakidae): host and geographical distribution on Neotropical and Antarctic species. Zootaxa 5256:43–62. https://doi.org/10.11646/zootaxa.5256.1.3

Gibson DI, Harris E, Bray R, Hussey C (2022) Host-Parasite database [. https://doi.org/10.5519/nplk7c5u. Dataset] Natural History Museum

González MJ, Gallardo JM, Brickle P, Medina I (2007) Nutritional composition and safety of *Patagonotothen ramsayi*, a discard species from Patagonian Shelf. Int J Food Sci Technol 42:1240–1248. https://doi.org/10.1111/j.1365-2621.2006.01472.x

González-Acuña D, Llanos-Soto S, Oyarzún-Ruiz P, Kinsella JM, Barrientos C, Thomas R, Cicchino A, Moreno L (2020) Parasites of the Neotropic cormorant *Nannopterum (Phalacrocorax) brasilianus* (Aves, Phalacrocoracidae) in Chile. Rev Bras Parasitol Vet 29. https://doi.org/10.1590/s1984-29612020049

Gosztonyi AE, Kuba L (1998) Fishes in the diet of the Imperial cormorant *Phalacrocorax atriceps* at Punta Loberia, Chubut, Argentina. Mar Ornithol 26:59–61. https://doi.org/10.5038/2074-1235.26.1.401

Gutiérrez RO (1943) Sobre la morfología de una nueva especie de Contracaecum (Nematoda: Ascariroidea). Rev Bras Biol 3:159–172

Harris S, Samaniego RAS, Rey AR (2016) Insights into diet and foraging behavior of Imperial Shags (*Phalacrocorax atriceps*) breeding at Staten and Becasses Islands, Tierra Del Fuego, Argentina. Wilson J Ornithol 128:811–820. https://doi.org/10.1676/15-141.1

Hecht T (1987) A guide to the otoliths of Southern Ocean fishes. S Afr J Antarct Res 17:1–87

Hermosilla C, Hirzmann J, Silva LMR, Brotons JM, Cerdà M, Prenger-Berninghoff E, Ewers C, Taubert A (2018) Occurrence of anthropozoonotic parasitic infections and faecal microbes in free-ranging sperm whales (*Physeter macrocephalus*) from the

Mediterranean Sea. Parasitol Res 117:2531–2541. https://doi.org/10.1007/s00436-018-5942-3
Huizinga H (1966) Studies on the life cycles and development of *Contracaecum spiculigerum* (Rudolphi, 1809) (Ascaridoidea: Heterocheilidae) from marine piscivorous birds. J Elisha Mitchell Sci Soc 82:181–195
Ibarra C, Marinao C, Suárez N, Yorio P (2018) Differences between colonies and chick-rearing stages in Imperial Cormorant (*Phalacrocorax atriceps*) diet composition: implications for trophic studies and monitoring. Wilson J Ornithol 130:224–234. https://doi.org/10.1676/16-184.1
Jobling M, Breiby A (1986) The use and abuse of fish otoliths in studies of feeding habits of marine piscivores. Sarsia 71:265–274. https://doi.org/10.1080/00364827.1986.10419696
Johnsgard PA (1993) Cormorants, Darters, and Pelicans of the World. Smithsonian Institution Press, Washington, D.C., USA
Johnstone IG, Harris MP, Wanless S, Graves JA (1990) The usefulness of pellets for assessing the diet of adult Shags *Phalacrocorax aristotelis*. Bird Study 37:5–11. https://doi.org/10.1080/00063659009477030
Kleinertz S, Christmann S, Silva LMR, Hirzmann J, Hermosilla C, Taubert A (2014) Gastrointestinal parasite fauna of Emperor Penguins (*Aptenodytes forsteri*) at the Atka Bay, Antarctica. Parasitol Res 113:4133–4139. https://doi.org/10.1007/s00436-014-4085-4
Køie M, Fagerholm HP (1995) The life cycle of *Contracaecum osculatum* (Rudolphi, 1802) *sensu stricto* (Nematoda, Ascaridoidea, Anisakidae) in view of experimental infections. Parasitol Res 81:481–489
Laptikhovsky VV (2004) A comparative study of diet in three sympatric populations of *Patagonotothen* species (Pisces: Nototheniidae). Polar Biol 27:202–205. https://doi.org/10.1007/s00300-003-0573-1
MacKenzie K, Longshaw M (1995) Parasites of the hakes *Merluccius australis* and *M. Hubbsi* in the waters around the Falkland Islands, southern Chile, and Argentina, with an assessment of their potential value as biological tags. Can J Fish Aquat Sci 52:213–224. https://doi.org/10.1139/f95-529
MacKenzie K, Brickle P, Hemmingsen W, George-Nascimento M (2013) Parasites of hoki, *Macruronus magellanicus*, in the Southwest Atlantic and Southeast Pacific Oceans, with an assessment of their potential value as biological tags. Fish Res 145:1–5. https://doi.org/10.1016/j.fishres.2013.03.008
Malacalza E, Poretti I, Marcelo N (1994) La dieta de *Phalacrocorax albiventer* en Punta Leon (Chubut, Argentina) durante la temporada reproductiva. Ornitol Neotrop 5:91–97
Malacalza E, Bertellotti M, Poretti T (1998) Presencia de nematodes (*Contracaecum sp.*) en 'pellets' del cormoran real (*Phalacrocorax albiventer*) en Punta Leon, Chubut, Argentina. Nat Patag 6:29–34
Michalik A, Van Noordwijk HJ, Brickle P, Eggers T, Quillfeldt P (2010) The diet of the Imperial Shag *Phalacrocorax atriceps* at a colony on New Island, Falkland/Malvinas Islands combining different sampling techniques. Polar Biol 33:1537–1546. https://doi.org/10.1007/s00300-010-0843-7
Moravec F (2009) Experimental studies on the development of *Contracaecum rudolphii* (Nematoda: Anisakidae) in copepod and fish paratenic hosts. Folia Parasitol 56:185–193. https://doi.org/10.14411/fp.2009.023
Morgenthaler A, Millones A, Gandini P, Frere E (2022) Diet of adult and immature Imperial cormorants, *Leucocarbo atriceps*, from southern Patagonia. A combined dietary approach and an exploratory analysis of stable isotopes of pellet membrane. Polar Biol 45:1529–1539. https://doi.org/10.1007/s00300-022-03086-5
O'Hear M, Pote L, Yost M, Doffitt C, King T, Panuska C (2014) Morphologic and molecular identifications of digenetic trematodes in double-crested cormorants (*Phalacrocorax auritus*) from the Mississippi Delta, USA. J Wildl Dis 50:42–49. https://doi.org/10.7589/2012-10-249
Oyarzún-Ruiz P, Muñoz-Alvarado P (2015) Cormorán imperial, Phalacrocorax atriceps (Aves, Phalacrocoracidae): Nuevo hospedero para un parásito respiratorio, Cyathostoma (Cyathostoma) phenisci (Nematoda, Syngamidae). Rev Biol Mar Oceanogr 50:353–358. https://doi.org/10.4067/S0718-19572015000300013
Pennycott T (2016a) Seabirds. Images of helminths (nematodes, cestodes, trematodes and thorny-headed worms) and pentastomid 'tongue-worms'., 1994–2013 [image]. University of Edinburgh. Royal (Dick) School of Veterinary Studies. https://doi.org/10.7488/ds/1566
Pennycott T (2016b) Seabirds. Images of pathology (miscellaneous diseases 1)., 1994–2013 [image]. University of Edinburgh. Royal (Dick) School of Veterinary Studies. https://doi.org/10.7488/ds/1564
Punta G, Saravia JRC, Yorio PM (1993) The diet and foraging behaviour of two Patagonian cormorants. Mar Ornithol 21:27–36. https://doi.org/10.5038/2074-1235.21.1.292
Punta G, Yorio P, Herrera G (2003) Temporal patterns in the diet and food partitioning in Imperial Cormorants (*Phalacrocorax atriceps*) and Rock Shags (*P. magellanicus*) breeding at Bahía Bustamante, Argentina. Wilson Bull 115:307–315
Quillfeldt P, Schroff S, Van Noordwijk H, Michalik A, Ludynia K, Masello J (2011) Flexible foraging behaviour of a sexually dimorphic seabird: Large males do not always dive deep. Mar Ecol Prog Ser 428:271–287. https://doi.org/10.3354/meps09058
Quintana F, Wilson R, Dell'Arciprete P, Shepard E, Laich AG (2011) Women from Venus, men from Mars: inter-sex foraging differences in the Imperial cormorant *Phalacrocorax atriceps* a colonial seabird. Oikos 120:350–358. https://doi.org/10.1111/j.1600-0706.2010.18387.x
Reid K (1996) A guide to the use of otoliths in the study of predators at South Georgia. British Antarctic Survey, Cambridge
Santos CP, Borges JN (2020) Current knowledge of small flukes (Digenea: Heterophyidae) from South America. Korean J Parasitol 58:373–386. https://doi.org/10.3347/kjp.2020.58.4.373
Semenova MK (1979) The role of copepods in the life-cycle of *Contracaecum micropapillatum* (Ascaridata, Anisakidae). Trudy Gel'mintologicheskoi Laboratorii. (Gel'minty Zhivotnykh i Rastenii) 29:126–129
Sepulveda MS, Spalding MG, Kinsella JM, Bjork RD, Mclaughlin GS (1994) Helminths of the Roseate Spoonbill, *Ajaia ajaja*, in Southern Florida. J Helminthol Soc Wash 61:179–189
Sepulveda MS, Spalding MG, Kinsella JM, Forrester DJ (1996) Parasitic helminths of the Little Blue Heron, *Egretta caerulea*, in Southern Florida. J Helminthol Soc Wash 63:136–140
Sepulveda MS, Spalding MG, Kinsella JM, Forrester DJ (1999) Parasites of the Great Egret (*Ardea albus*) in Florida and a review of the Helminths reported for the species. J Helminthol Soc Wash 66:7–13
Shamsi S, Gasser R, Beveridge I, Alizadeh Shabani A (2008) *Contracaecum pyripapillatum* n. sp. (Nematoda: Anisakidae) and a description of *C. multipapillatum* (von Drasche, 1882) from the Australian pelican, *Pelecanus conspicillatus*. Parasitol Res 103:1031–1039. https://doi.org/10.1007/s00436-008-1088-z
Signorell A (2024) DescTools: Tools for Descriptive Statistics. Version 0.99.54. CRAN
Thompson K (1989) An assessment of the potential for competition between seabirds and fisheries in the Falkland Islands. Falkland Islands Foundation Project Report
Torres P, Ruíz E, Rebolledo C, Mira A, Cubillos V, Navarrete N, Gesche W, Montefusco A, Valdés L, Alberdi A (1990) Parasitismo en peces y comunidades humanas ribereñas de los lagos Huillinco y Natri: Isla Grande de Chiloé: Chile. Bol Chil Parasitol 45:47–55

Van Der Grient J, Morley S, Arkhipkin A, Bates J, Baylis A, Brewin P, Harte M, White JW, Brickle P (2023) The Falkland Islands marine ecosystem: A review of the seasonal dynamics and trophic interactions across the food web. Adv Mar Biol 94:1–68. https://doi.org/10.1016/bs.amb.2023.01.001

Violante-González J, Monks S, Gil-Guerrero S, Rojas-Herrera A, Flores-Garza R, Larumbe-Morán E (2011) Parasite communities of the neotropical cormorant *Phalacrocorax brasilianus* (Gmelin) (Aves, Phalacrocoracidae) from two coastal lagoons in Guerrero state, Mexico. Parasitol Res 109:1303–1309. https://doi.org/10.1007/s00436-011-2377-5

Volpedo AV, Echeverría DD (2000) Catálogo y claves de otolitos para la identificación de peces del mar argentino. 1: Peces de importancia económica. Dunken, Buenos Aires

Woods RW (2001) A survey of the number, size and distribution of islands in the Falklands archipelago. Falkl Isl J 7:1–25

Xavier JC, Cherel Y (2021) Cephalopod beak guide for the Southern Ocean: An update on taxonomy. British Antarctic Survey, Cambridge, UK

Xavier JC, Cherel Y, Boxshall GA, Brandt A, Coffer T, Forman J, Havermans C, Jażdżewska AM, Kouwenberg J, Schiaparelli S, Schnabel K, Siegel V, Tarling GA, Thatje S, Ward P, Gutt J, with Scientific Committee on Antarctic Research) (2020) (Crustacean guide for predator studies in the Southern Ocean. Original edition. Scientific Committee on Antarctic Research (SCAR), Cambridge, UK

Yamaguti S (1971) Synopsis of digenetic trematodes of vertebrates. Vols I and II. Keigaku Publishing Co., Tokyo

Yamaguti S (1975) A synoptical review of life histories of digenetic trematodes of vertebrates: With special reference to the morphology of their larval forms, 1st ed. Yugakusha, Tokyo

Yorio P, Copello S, Kuba L, Gosztonyi A, Quintana F (2010) Diet of Imperial Cormorants *Phalacrocorax atriceps* breeding at Central Patagonia, Argentina. Waterbirds 33:70–78. https://doi.org/10.1675/063.033.0108

Yorio P, Ibarra C, Marinao C (2017) Induced regurgitation versus stomach sampling: assessing their value for the characterization of Imperial Cormorant (*Phalacrocorax atriceps*) diet. Waterbirds 40:162–167. https://doi.org/10.1675/063.040.0208

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.