Corrections to

"Bifurcation from a saddle connection in functional differential equations: An approach with inclination lemmas"

(Dissertationes Math. 291 (1990))

by Hans-Otto Walther (München)

Page 21, line 16: Delete "and is continuous".

Include "Each map $X(t,\cdot,a):C\to C$ is of class C^1 ."

Page 31, line 10: Add a line:

"(vi) The maps DG, DG^-, D_1G, D_1G^- are bounded."

Page 32, lines 14 and 15: Delete "and are continuous".

Add "Each map $Y(t,\cdot,a), R(t,\cdot,a)$ is of class C^1 ."

Page 32, lines 24–26: Delete "The assignments . . . into $L_c(C,C)$."

Page 33, lines 10-14: Replace these lines by the following text.

"Proposition 5.! There exists a constant const ≥ 0 such that we have

(5.6)
$$|D_2 R_{p_a}(t, \psi)| + |D_2 R_{q_a}(t, \psi)| < const$$

for all $(t, \psi, a) \in [0, N] \times D^1 \times A_7$.

Proof. Let
$$(t,\psi,a)\in [0,N] imes D^1 imes A_7$$
 be given. We have

$$|D_2R(t,\psi,a)| \leq |D_2Y(t,\psi,a)| + |T(t,\cdot,a)|$$

and

$$|D_2Y(t,\psi,a)| \le \sup |D_1G||D_2X(t,G^-(\psi,a),a)| \sup |D^1G^-|$$

$$\le \sup |D_1G| \sup |D_1G^-|(1+\max |h'|)^{N+1},$$

by Proposition 5.1(vi) and Corollary 3.1. There is a constant $k_{00} \ge 1$ such that

$$|T(t,\cdot,a)| \leq k_{00}e^{-\lambda t}$$
 for all $t \geq 0$, $a \in A_7 \subset A_3$.

Now the desired estimate becomes obvious.

From Proposition 5.3(i) we infer that there exist an open ball $D^{2.1} \subset D^1$, centered at $0 \in C$, and an open interval A_8 (with $\operatorname{cl} A_8 \subset A_7$) such that we have"

 $(5.7) \dots$

Page 34, lines 25 and 27: Replace "c" by "const".

Page 34, line 28: Replace " $(1+c+k_0)e^{-\lambda_1 N}$ " by " $(1+const+k_0)e^{-\lambda_1 N}$ ".

Page 35, lines 10 and 11: Write

"
$$|p_a \circ Y_a(3, \psi)| \le (const + 1)e^{3\mu_2}|p_a\psi|$$
".

Page 35, line 21: Write

$$c_4 := \frac{c_3}{(const + 1)e^{3\mu_2}}.$$

Page 36, lines 25 and 26: Replace "c" by "const".

MATHEMATISCHES INSTITUT LUDWIG-MAXIMILIANS -UNIVERSITÄT THERESIENSTR. 39 D-8000 MÜNCHEN 2, GERMANY