

having decision-making power, which will be crucial for lasting impact and meaningful progress.

We declare no competing interests.

*Hrushitaa Murali, Cecilia Ashaley Fofo, *Awiwe Funani, Sireesha J Bobbili, Melani O'Leary, Byron Bitanihirwe, Antonis A Kousoulis, Sarah Kline*
awiwe@unitedgmh.org

Being Initiative People with Lived Experience Advisory Group, Toronto, ON, Canada (HM, CAF); Huesofthemind Mental Wellbeing Foundation, Gurugram, India (HM); Basic Needs Ghana, Accra, Ghana (CAF); United for Global Mental Health, London E1 6RA, UK (AF, AAK, SK); Grand Challenges Canada, Toronto, ON, Canada (SJB, MO); Science for Africa Foundation, Nairobi, Kenya (BB); Global Mental Health Action Network, London, UK (AAK)

- 1 McGorry PD, Mei C, Dalal N, et al. The *Lancet Psychiatry* Commission on youth mental health. *Lancet Psychiatry* 2024; **11**: 731–44.
- 2 Amarasekera S, Ugo V, Florez V, Patalay P. Research task shifting and youth as drivers of a more global mental health science. *Lancet Psychiatry* 2021; **8**: 1027–29.
- 3 UNICEF. Young people's participation and mental health. August, 2022. <https://www.unicef.org/reports/young-peoples-participation-and-mental-health> (accessed June 4, 2024).
- 4 Partnership for Maternal Newborn & Child Health. Make adolescent well-being a priority. Oct 10, 2020. <https://pmnch.who.int/resources/publications/m/item/make-adolescent-well-being-a-priority> (accessed June 4, 2024).
- 5 Bell J, Lim A, Williams R, Girdler S, Milbourn B, Black M. 'Nothing about us without us': co-production ingredients for working alongside stakeholders to develop mental health interventions. *Adv Ment Health* 2023; **21**: 4–16.
- 6 The Being Initiative. Mapping youth mental health landscapes: local insights from 13 countries. April 18, 2024. https://being-initiative.org/wp-content/uploads/2024/04/Mapping-Youth-Mental-Health-Landscapes_April-18-2024.pdf (accessed June 13, 2024).
- 7 Global Mental Health Action Network. Collaborative resource: guiding principles and recommendations for effective lived experience youth engagement practices. August, 2023. <https://gmhan.org/news/youth-engagement-guidelines> (accessed June 13, 2024).
- 8 African Union Office of the Youth Envoy. Youth mental health toolkit on 5 emerging issues in Africa. February, 2024. https://au.int/sites/default/files/documents/43549-doc-AU_Youth_Mental_Health_Toolkit_A4.v5.1.pdf (accessed June 17, 2024).
- 9 Channel Kindness. Welcome to our 2023 advisory board! Jan 30, 2023. <https://www.channelkindness.org/2023-advisory-board/> (accessed June 13, 2024).
- 10 Knowles S, Sharma V, Fortune S, Wadman R, Churchill R, Hetrick S. Adapting a codesign process with young people to prioritize outcomes for a systematic review of interventions to prevent self-harm and suicide. *Health Expect* 2022; **25**: 1393–404.



WHO treatment guideline for mental disorders

Published Online
 July 24, 2024
[https://doi.org/10.1016/S2215-0366\(24\)00169-X](https://doi.org/10.1016/S2215-0366(24)00169-X)
 See [Comment](#) page 678

In 2024, WHO issued updated guidelines for treating mental health conditions, emphasising evidence-based manual-guided psychotherapeutic treatments.¹ We applaud WHO's effort to broaden access to evidence-based psychotherapy. Yet, we are concerned that the recommendations predominantly endorse behavioural therapy and cognitive behavioural therapy for both adults and young people. This selection overlooks the substantial body of evidence supporting other therapeutic approaches, including but not limited to psychodynamic therapy, except for a brief acknowledgment of this method's use in treating depression. High-quality research reviews published in leading scientific journals² have shown that manual-guided psychodynamic therapy meets the updated American Psychological Association's criteria for empirically supported treatments,³ based on several comprehensive meta-analyses.² This endorsement covers the psychodynamic treatment of depressive, anxiety, somatoform, and personality disorders, with clinically meaningful effect sizes compared with controls, no meaningful differences in efficacy compared with other evidence-based treatments, and confidence intervals comparable with, for example, cognitive behavioural therapy.² The quality (certainty) of evidence was similar to that on which WHO based their recommendations (eg, for depressive disorders, anxiety disorders, or self-harm and suicide).⁴ In line with the American Psychological

Association's criteria for evidence-based treatments, this substantial body of evidence was shown to warrant a strong recommendation of psychodynamic therapy in the conditions listed.²

Further evidence for psychodynamic therapy's efficacy exists for the treatment of many other mental conditions including somatoform disorders,^{5,6} eating disorders,⁵ post-traumatic stress disorder,⁵ substance-related disorders (opiate addiction),^{5,6} and treatment of children and adolescents with depression.⁷ Psychodynamic psychotherapy is included in the UK guidelines on childhood depression developed by the National Institute for Health and Care Excellence. Additionally, evidence exists that psychodynamic therapy is efficacious in patients whose symptoms are unresponsive to other treatments, including individuals with treatment-resistant or chronic somatoform disorders⁵ or treatment-resistant chronic depression. Psychodynamic therapy has also been shown to help reduce suicide attempts and self-harm in both adults and adolescents.⁸ For the latter, however, WHO recommended only digital stand-alone methods based on cognitive behavioural therapy, dialectical behaviour therapy, problem-solving therapy, and mindfulness, with only a conditional recommendation based on low-certainty of evidence.¹

Evidence shows that behavioural therapy and cognitive behavioural therapy cannot be viewed

as the sole solutions for mental health care. The response rates, for example, of cognitive behavioural therapy in depressive and anxiety disorders, including obsessive-compulsive and post-traumatic stress disorders, range between 34% and 44%,⁹ with even lower remission rates. These data underscore that no single psychotherapy approach can be regarded as the definitive treatment for all patients. A substantial number of patients who do not respond to behavioural therapy or cognitive behavioural therapy might benefit from alternative evidence-based psychotherapeutic approaches and vice versa. In clinical medicine, no one would recommend one drug with a response rate of 40% as the only treatment for all patients when other drugs with similar efficacy were available.

Our concern is heightened by the WHO's recommendation of primarily a singular psychotherapy approach,¹⁴ possibly leaving many patients without access to more suitable treatments that could offer them substantial benefits. For example, for anxiety disorders, WHO noted that further research is needed to clarify whether other therapies than cognitive behavioural therapy can offer similar benefits.⁴ We highlight such evidence for psychodynamic therapy, not only for anxiety disorders but also for various other mental conditions.^{2,5,6}

With regard to implementation, there is evidence that can be obtained from the first author that psychodynamic therapies are adaptable and can be effectively taught to new practitioners from diverse theoretical and professional backgrounds. Substantial evidence points to the broad applicability and efficacy of psychodynamic methods.^{2,5} Benefits of these methods outweigh the costs and harms.² In addition, psychodynamic therapies can be delivered in guided and online formats making them accessible and distributable on a wide scale.^{5,6}

For these reasons, we advocate the incorporation of psychodynamic therapies, among other evidence-based psychotherapeutic approaches, into the WHO guidelines. We also recommend the involvement of experts in the WHO guideline development group who have knowledge on the outcomes of other approaches than cognitive behavioural therapy, such as psychodynamic or interpersonal therapy. This procedure might help to improve transparency and to avoid biased recommendations. This proposal is consistent with recommendations for advancing guideline development

in health care. Although evidence exists that reporting quality of WHO guidelines has generally improved over the years, the reporting quality of guidelines can be enhanced in several areas.¹⁰ By embracing a broader array of empirically supported therapeutic methods, the goal is to raise the overall quality and efficacy of global mental health care.

All authors have been involved in the development and evaluation of psychodynamic treatments. FL has received grants from the German Ministry of research and royalties from Hogrefe publisher. PL has been involved in the development, evaluation, and dissemination of psychodynamic treatments; and has received royalties and grants in these contexts. NM has received grants from the Association of Child Psychotherapists (UK), and the International Psychoanalytic Association. BM has received grants from American Psychoanalytic Association Fund for Psychoanalytic Research, Einstein-Rockefeller-CUNY CFAR (grant number P30 AI124414); received the grant of Jonathan Alpert MD PhD, University Chair, Department of Psychiatry, Albert Einstein College of Medicine and Multiple; and honoraria for psychiatric talks. All other authors declare no competing interest.

**Falk Leichsenring, Allan Abbass, Peter Fonagy, Kenneth N Levy, Peter Lillengren, Patrick Luyten, Nick Midgley, Barbara Milrod, Christiane Steinert*
falk.leichsenring@psycho.med.uni-giessen.de

Department of Psychosomatics and Psychotherapy, University of Giessen, Giessen 35392, Germany (FL,CS); University of Rostock, Department of Psychosomatics and Psychotherapy, Rostock, Germany (FL); Department of Psychology, College of the Liberal Arts, Penn State University, University Park, PA, USA (KNL); Department of Psychiatry, Dalhousie University, Halifax, NS, Canada (AA); Research Department of Clinical, Educational and Health Psychology, University College London, London, UK (PF, PLu, NM); Department of Psychology, University of Stockholm, Sweden (PLi); Faculty of Psychology and Educational Sciences, University of Leuven, Leuven, Belgium (PLu); Albert Einstein College of Medicine, New York, NY, USA (BM); International Psychoanalytic University, Berlin, Germany (CS)

- 1 WHO. Implementation manual for psychological interventions: integrating evidence-based psychological interventions into existing services. 2024. <https://www.who.int/publications/i/item/9789240087149>, (accessed May 29, 2024).
- 2 Leichsenring F, Abbass A, Heim N, et al. The status of psychodynamic psychotherapy as an empirically supported treatment for common mental disorders—an umbrella review based on updated criteria. *World Psychiatry* 2023; **22**: 286–304.
- 3 Tolin D, McKay D, Forman EM, Klonsky ED, Thombs BD. Empirically supported treatments: recommendations for a new model. *Clin Psychol* 2015; **22**: 317–38.
- 4 WHO. Mental health gap action programme (mhGAP) guideline for mental, neurological and substance use disorders. 2023. <https://www.who.int/publications/i/item/9789240084278> (accessed May 29, 2024).
- 5 Leichsenring F, Luyten P, Hilsenroth MJ, et al. Psychodynamic therapy meets evidence-based medicine: a systematic review using updated criteria. *Lancet Psychiatry* 2015; **2**: 648–60.
- 6 Lillengren P. A comprehensive overview of randomized controlled trials of psychodynamic psychotherapies. *Psychoanal Psychother* 2023; **37**: 117–40.
- 7 Midgley N, Mortimer R, Bhatra P, Cirasola A, Kennedy E. The evidence-base for psychodynamic psychotherapy with children and adolescents: a systematic review and narrative synthesis. *Front Psychol* 2021; **12**: 662671.
- 8 Briggs S, Netuveli G, Gould N, et al. The effectiveness of psychoanalytic/psychodynamic psychotherapy for reducing suicide attempts and self-harm: systematic review and meta-analysis. *Br J Psychiatry* 2019; **214**: 320–28.
- 9 Cuijpers P, Miguel C, Ciharova M, et al. Absolute and relative outcomes of psychotherapies for eight mental disorders: a systematic review and meta-analysis. *World Psychiatry* 2024; **23**: 267–75.
- 10 Wang X, Zhou Q, Chen Y, et al. Using RIGHT (Reporting Items for Practice Guidelines in Healthcare) to evaluate the reporting quality of WHO guidelines. *Health Res Policy Syst* 2020; **18**: 75.