



Complete Genome Sequence of *bla*_{CTX-M-27}⁻ Encoding *Escherichia coli* Strain H105 of Sequence Type 131 Lineage C1/H30R

Hiren Ghosh,^a Boyke Bunk,^b Swapnil Doijad,^a Judith Schmiedel,^a
Linda Falgenhauer,^a Cathrin Spröer,^b Can Imirzalioglu,^a Jörg Overmann,^b
Trinad Chakraborty^a

Institute of Medical Microbiology, Justus Liebig University, Giessen and German Center for Infection Research (DZIF), Partner site Giessen-Marburg-Langen, Giessen, Germany^a; Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures, German Center for Infection Research (DZIF), Partner site Hannover-Braunschweig, Braunschweig, Germany^b

ABSTRACT *Escherichia coli* sequence type 131 (ST131) is the most frequent antimicrobial-resistant lineage of *E. coli*, propagating extended-spectrum β -lactamases (ESBL) worldwide. Recently, an alarming rate of increase in isolates of the sublineage C1/H30R-*bla*_{CTX-M-27} of ST131 in geographically distant countries was reported. Here, we present the complete genome sequence of the ST131 sublineage C1/H30R *E. coli* isolate harboring *bla*_{CTX-M-27} from Germany.

We sequenced the *Escherichia coli* isolate H105, a sequence type 131 (ST131) C1/H30R extended-spectrum β -lactamase (ESBL)-producing isolate obtained in 2010 from a vaginal swab sample of an individual from Germany. The isolate harbors a *bla*_{CTX-M-27} allele and represents an early isolate with this genotype combination in our collection. As reported in Japan (1) and in France (2), there is currently a strong surge in the number of ST131-C1/H30R-*bla*_{CTX-M-27} isolates detected in Germany, particularly in the past 3 years, suggesting an ongoing shift in CTX-M alleles associated with ST131 infections.

For the whole-genome sequencing, genomic DNA was isolated using a PureLink genomic DNA kit (Invitrogen, Darmstadt, Germany) from an overnight culture grown at 37°C in LB medium. Short-read sequencing was performed on an Illumina MiSeq machine (Illumina, the Netherlands) using a Nextera XT library with MiSeq v3 (2 × 300 bp) reagent kit. Single-molecule real-time sequencing (SMRT) was conducted using the PacBio RSII system (Pacific Biosciences, USA).

De novo genome assembly of 59,447 PacBio reads with an average read length of 10,355 bp was performed using the “RS_HGAP_Assembly.3” included in the SMRT Portal version 2.3.0. Subsequently, Illumina short reads were mapped onto the assembled sequences in order to obtain a highly accurate genome with QV60 final quality. The chromosome was adjusted to *dnaA* as the first gene. Annotation was performed both using Prokka 1.10 (3) and the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (<http://www.ncbi.nlm.nih.gov/genomes/static/Pipeline.html>). Antimicrobial resistance genes and prophages were identified using Resfinder and PHAST, respectively (4, 5).

The assembly resulted in a closed, circular chromosomal contig of 4,978,342 bp and a plasmid of 134,499 bp, designated the plasmid pH 105. The chromosome of *E. coli* H105 exhibited a 50.7% G+C content, and harbored 4,635 open reading frames, 87 tRNAs, 15 rRNAs, and 6 intact prophage regions. *In silico* analysis revealed that *E. coli* H105 belongs to the sequence type ST131, serotype O25:H4, and phylogroup B2. The genome of H105 includes multiple chromosomally encoded virulence genes such as for

Received 13 June 2017 Accepted 19 June 2017 Published 3 August 2017

Citation Ghosh H, Bunk B, Doijad S, Schmiedel J, Falgenhauer L, Spröer C, Imirzalioglu C, Overmann J, Chakraborty T. 2017. Complete genome sequence of *bla*_{CTX-M-27}-encoding *Escherichia coli* strain H105 of sequence type 131 lineage C1/H30R. *Genome Announc* 5: e00736-17. <https://doi.org/10.1128/genomeA.00736-17>.

Copyright © 2017 Ghosh et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Trinad Chakraborty, trinad.chakraborty@mikrobio.med.uni-giessen.de.

iron uptake (yersiniabactin, aerobactin, *sitABC*), serum resistance (*iss*, *traT*), autotransporter proteases (*sat*, *pic*), and the postsegregational killing system *ccdA/ccdB*.

The plasmid pH105 is a multireplicon plasmid, depicting IncFIA, FIB, and FII replicons with the pMLST type F1:A2:B20 (6). The ESBL gene *bla*_{CTX-M-27} encoded on pH105 confers resistance to extended-spectrum β -lactams. In addition, pH105 also carries genes conferring resistance to aminoglycosides (*aadA5*, *strA*, *strB*), macrolides [*mph*(A)], tetracyclines [*tet*(B)], sulfonamide (*sul1*, *sul2*), and trimethoprim (*dfrA17*).

The sequence of H105 represents the first complete genome of the *bla*_{CTX-M-27}-encoding *E. coli* ST131 of the lineage C1/H30R. The high-quality genome of H105 will serve as a valuable resource for comparative studies on epidemiology of globally emerging ST131-C1/H30R-*bla*_{CTX-M-27}.

Accession number(s). The complete genome sequence of *E. coli* H105 has been deposited in GenBank under GenBank accession no. [CP021454](#) (chromosome) and [CP021871](#) (plasmid).

ACKNOWLEDGMENTS

This study was supported by grants from the Federal Ministry of Education and Research (BMBF, Germany) within the framework of the RESET research network (contracts 01KI1013G and 01KI1313G) and the German Center for Infection Research (DZIF/grants 8000 701-3 [HZI] to T.C. and C.I. and TI06.001 and 8032808811 to T.C.).

We thank Christina Gerstmann, Natalia Lest, Simone Severitt, and Nicole Mrotzek for excellent technical assistance.

REFERENCES

1. Matsumura Y, Pitout JDD, Gomi R, Matsuda T, Noguchi T, Yamamoto M, Peirano G, DeVinney R, Bradford PA, Motyl MR, Tanaka M, Nagao M, Takakura S, Ichihama S. 2016. Global *Escherichia coli* sequence type 131 clade with *bla*_{CTX-M-27} gene. *Emerg Infect Dis* 22:1900–1907. <https://doi.org/10.3201/eid2211.160519>.
2. Birgy A, Bidet P, Levy C, Sobral E, Cohen R, Bonacorsi S. 2017. CTX-M-27-producing *Escherichia coli* of sequence type 131 and clade C1-M27, France. *Emerg Infect Dis* 23:885. <https://doi.org/10.3201/eid2305.161865>.
3. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
4. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
5. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
6. Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller Aarestrup F, Hasman H. 2014. *In silico* detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. <https://doi.org/10.1128/AAC.02412-14>.